

POLITICAL DEPARTMENT

RAJPUTANA AGENCY

OFFICE MANUAL



FOR OFFICIAL USE ONLY.

SIMLA
MANAGER GOVERNMENT OF INDIA PRESS
1927

IX.—ARRANGEMENT OF A CASE—

Single case	45
Binding of files	46
Linking cases	47

X.—FIRST ACTION IN THE BRANCH—

New subjects	48—51
Pending subject	52
Letters from Princes	53
Printed copies of letters received	54

XI.—REFERENCING—

"Referencing" defined	55
New file	56
Addition to file	57

XII.—NOTES—

Object	58
Contents	59
Incorrect notes	60
Routine notes	61
Notes on P U C	62
Form of ordinary notes	63
Marginal notes	64
Notes to be initialled	65
Branch clerk's note	66
Head clerk's note	67
Further duties of branch clerk	68 & 69

XIII.—SUBMISSION FOR ORDERS—

Submission to Secretary	70
Books, etc	71
Case affecting two branches	72
Head clerk's duties	73
Return by the Agent to the Governor-General	74

XIV.—DRAFTS—

Object	75
When draft is unnecessary	76
Forms of communications	77
Letters	78
Memorandums	79
Endorsements	80
Telegrams	81
Spare Copies of a General Communication	82
Demi-official letters	83
Unofficial references	84
Preparation of the draft	85
Form of draft	86—88
Enclosures	89
Copies of enclosures	90
Addressees	91
First reminder	92
Reference in reply	93
Place in case	94

TABLE OF CONTENTS

I—Preliminary—	Para
Extent of use	1
Definitions	2
II—Attendance Holidays and Casual Leave—	
Office hours	3
Leave	4
Gazetted holidays	5
Casual leave	6
Temporary absence from office	7
Register of leave	8
III—General Discipline Etc—	
Official information to be treated as confidential	9
Removal of official papers	10
Character rolls	11
Certificates	12
Notes affecting discipline or conduct	13
IV—Office Organisation and System—	
Object	14
Distribution of work	15
Head Clerk's responsibilities	16
General outline of system	17
V—Duties of the Registrar—	
General supervisory	18 & 19
Character rolls	20
Dak and route papers	21
Statutory and forms	22
VI—Opening the Dak	23
VII—Registration of Letters—	
Objects	24 & 25
Registers	26 & 27
Numbering	28
Abbreviations	29
When registration is unnecessary	30
Register of miscellaneous correspondence	31
Register of files	32
Periodical returns	33
Opening new registers	34
Central record room	35
VIII—Indexing—	
Object	36
Card or loose leaf index	37
Major and minor heads	38
Cross references	39
Cards	40
Branch index	41
Working cases	42
Subsidiary indices	43
When indexing is unnecessary	44

IX — ARRANGEMENT OF A CASE —	
Single case	45
Binding of files	46
Linking cases	47
X — FIRST ACTION IN THE BRANCH —	
New subjects	48—51
Pending subject	52
Letters from Princes	53
Printed copies of letters received	54
XI — REFERENCING —	
Referencing defined	55
New file	56
Addition to file	57
XII — NOTES —	
Object	58
Contents	59
Incorrect notes	60
Rout ne notes	61
Notes on P U C	62
Form of ordinary notes	63
Marginal notes	64
Notes to be initialed	65
Branch clerk's note	66
Head clerk's note	67
Further duties of branch clerk	68 & 69
XIII — SUBMISSION FOR ORDERS —	
Submission to Secretary	70
Books etc	71
Case affecting two branches	72
Head clerk's duties	73
Return by the Agent to the Governor General	74
XIV — DRAFTS —	
Object	75
When draft is unnecessary	76
Forms of communications	77
Letters	78
Memorandums	79
Endorsements	80
Telegrams	81
Spare Copies of a General Communication	82
Semi-official letters	83
Unofficial references	84
Preparation of the draft	85
Form of draft	86—88
Enclosures	89
Copies of enclosures	90
Addressees	91
First reminder	92
Reference in reply	93
Index in case	94

XV—FAIR COPYING—

Definition	95
When fair copy to be made in branch	96
When draft sent to copying branch	97
Date	98
Number	99
Comparing	100
Signature	101

XVI—DESPATCH—

Papers to be sent to despatcher	102
Return of office copies, etc.	103
Telegrams, etc.	104
Hours of closing Dak to be fixed	105—107
Accumulation of papers	108
Closing Dak	109
Economy labels	110
Registered post	111
Stamps	112 & 113
Further action by branch clerk	114

XVII—REMINERS—

Diary	115
Issue of reminders	116
Signature and despatch	117
Subsequent reminders	118
Reminders to Government of India	119
Further uses of diary	120
Periodical returns	121

XVIII—CLOSING FILES—

When file should be closed	122
Orders to close	123
Procedure in closing	124 & 125

XIX—RECORD KEEPER—

Responsibilities	126
Pending cases	127
Closed files	128
Files taken out	129
Opening a file	130
Closing a file	131
Re-opening closed file prohibited	132
Destruction of records	133

XX—DUTIES OF CONFIDENTIAL CLERK

.. .. .	134 & 135
---------	-----------

XXI—MISCELLANEOUS—

Check on disposal of business	136—137
Interviews	138
Corrections	140
Signature and initials to be dated	141
Stationery	142
Furniture	143
Tracts	144

XXII —TOWN ARRANGEMENTS—

Modifications in procedure	165
Camp register	166
New file	167
List of files	168

APPENDIX I —Form A —Register of miscellaneous correspondence

Form B.—Register of files (and alternative forms to Form B.)

Form C.—Register of periodicals.

Form D.—File cover (front page).

Form E.—Stamp register.

APPENDIX II —Abbreviations.

APPENDIX III.—Directions regarding Indexing and Opening Files (with the list of the Major and Minor Heads)

APPENDIX IV —Specimen Forms.

APPENDIX V —Rules for the Destruction of Records and Periods for which they are to be retained.

APPENDIX VI —Instructions regarding forms in Appendix I.

MANUAL OF OFFICE PROCEDURE

I—PRELIMINARY

Extent of use—This Manual is primarily intended for use in the office of the Agent to the Governor General and Chief Commissioner. The same procedure should be adopted, as far as may be suitable, in the offices of the Political Agents and Residents, but, as their offices are smaller, the procedure will, in certain respects, be automatically simplified, in applying the Manual to the offices of Political Agents or Residents, necessary changes should be made, *e g*, for "Agent to the Governor-General" or "Secretary" the words 'Resident' or "Political Agent" should be read, and for "Registrar" or 'Superintendent' the words 'Head Clerk' should be read.

2 Definitions—(a) The term "Secretary" includes "Under Secretary" with reference to branches of the office under the Under Secretary, unless the contrary appears from the context.

(b) The term "Head Clerk" in these rules means the Superintendent, Head Clerk or Senior Clerk in charge of a branch, unless the contrary appears from the context.

(c) The term "branch clerk" means a clerk under a head clerk as defined above.

(d) The "P U C" means the first or latest receipt in the course of a correspondence upon which action has to be taken.

(e) A "file" consists of papers which are or have been under consideration and any notes written in connection with them.

(f) A "current file" or "pending file" is a file which has not been closed.

(g) A "case" consists of a current file, and any files, books, etc., put up for reference with it.

(h) The "subject" of a file or case is the major, minor and sub heads under which it is registered and indexed.

(i) A "receipt" means a communication received from outside the office or branch, and may include a note from the Agent to the Governor General or Secretary which has to be added to a pending file or on which a new file has to be opened.

II—ATTENDANCE, HOLIDAYS AND CASUAL LEAVE.

3 Office hours—The prescribed office hours are from 10-30 A.M. to 5-0 P.M. but a clerk may be required by the Registrar to work beyond those hours if necessary.

4 Leave—Applications for leave must be submitted through the Registrar to the Secretary in sufficient time to enable the latter to issue orders granting or refusing it before the leave is to begin. Any one absent for

that two days on the plea of illness must submit a medical certificate. If a man in a family will not ordinarily be considered a sufficient reason for granting leave, unless the sickness is infectious. In the latter case the Registrar should at once notify the medical officer so that he can visit the house and verify the information if he has not already done so.

5 **Gazetted Holidays**—Gazetted holidays will ordinarily be allowed to the classes mentioned in the notification issued each year but clerks may be required to work on gazetted holidays by the Secretary if necessary. The Secretary may grant a 'compensation' holiday to any clerk required to

I.

15-1

051

20.

two

If true,

the c

orders.

NOT:-

complaint

his wife,

his leave.

regular 100%

F

10

●

1179

88

nd

the

FIOR

he

to

tins

10 Removal of official papers -- No official paper out of the office or to his home without the permission of the Registrar

No clerk may take a copy of any portion of an official document which comes before him in his official capacity or to which he is allowed to have access without the previous permission of the Registrar, this rule applies also to copies of notes and orders on applications or memorials.

11 Character Rolls.—The character rolls and service books relating to members of the clerical and menial establishments will be kept by the Registrar in the prescribed form with an alphabetical index of their names. Any censure punishment or commendation passed on a clerk or menial should be entered in his character roll and in the service book. The character rolls will be submitted when there is any change in the personnel of the Secretary for any remarks which the outgoing Secretary may wish to record regarding those who worked under him. The character rolls and service books will be submitted when any proposals are made for promotion or the grant of an increment of pay.

12 Certificates—No certificate may be given to any member of the office establishment unless it is signed by the Secretary

13 Notes affecting Discipline or Conduct—Notes affecting office discipline or the conduct of clerks which arise in an ordinary file should not be kept with that file but should be sent to the Registrar

IV—OFFICE ORGANIZATION AND SYSTEM

14 Object—The general object of the office organization and system is to enable the clerical establishment to render due assistance to the Agent to the Governor-General and the Secretaries in the prompt and correct disposal of their official business. The Agent to the Governor-General and the Secretary should be relieved of unimportant work as far as possible and the work to be done should be distributed among the members of the staff according to their abilities and experience each having his own particular share which should not be done a second time by another member of the staff. It is often more troublesome and more exasperating to check, correct and redo the work of a clerk who has attempted something beyond his competence than to do the work in the first instance, and it causes more delay as well as waste of the clerk's time and energy which might have been employed otherwise with advantage. Reasonable checks and safeguards against delays, shirking of work and the accidental loss of papers must be adopted. Care must be taken that, while these are efficient, they are worth the labour expended on them.

15 Distribution of Work—The Agency office is divided into branches, and the work is distributed among the various branches according to subjects in accordance with the orders passed from time to time by the Secretary. Each branch will be in the charge of the Registrar, the Superintendent, a head clerk, or a senior clerk appointed for the purpose. In a small office it may be unnecessary to divide it into branches.

16 Head Clerk's responsibilities—The head clerk will be personally responsible for the work of the other clerks in the branch, the latter are appointed to assist him and to relieve him of comparatively unimportant work, but he must not expect them to do his share of the work of the branch, he must constantly pay due attention to training them, and time and trouble spent by him in this respect will be amply repaid by the additional relief which they will ultimately give him, the nature and extent of the work which he can give a particular branch clerk to do and the closeness of his supervision over the latter's work should depend on the abilities and experience of the branch clerk. The head clerk should not however call upon a branch clerk to do important work which he must ultimately do himself. The head clerk should in like manner realize that, while it is his duty to assist and relieve the Secretary, he is not expected to discharge all the duties of the Secretary. He should freely consult the Secretary personally if he has any doubt as to the action he should take in respect of a particular case.

When there is more than one clerk in a branch the head clerk will see in that branch that the work is distributed among the clerks by the best method. The distribution may be altered from time to time if necessary, to meet the work is fairly distributed according to the abilities of the clerks.

various clerks, but, as far as possible, a clerk should be left in charge of the same subjects continuously

17 General outline of system—The Dāk will be delivered to the addressee, initialled by him and sent to the Registrar. Each paper will then be sent to the branch concerned and registered. If it relates to a subject of general correspondence on which there is no pending file, a fresh file will be opened and indexed and previous correspondence likely to be useful in the disposal of the paper will be obtained from the records. The branch clerk will reference the papers and 'flag' the files put up for reference and, if necessary, make a précis under the instructions of the head clerk. The head clerk will note on the file. It will be open to him to suggest the method of disposal. The case will then be submitted to the Secretary unless it can be disposed of by the Registrar under his powers. The Secretary will either pass orders or obtain the orders of the Agent to the Governor-General. A draft will then be prepared by the head clerk or by the branch clerk under his instructions and will be submitted to the Secretary, who will, if necessary, submit the draft to the Agent to the Governor-General. In important or intricate cases the Secretary will prepare the draft himself. After the draft is passed, it will be sent to the copying branch where the fair copy will be made and sent to the officer concerned for signature, after signature it will be sent to the despatcher for despatch. The case will then be sent from the branch concerned to the record keeper who will either complete the closure of the file, if its closure has been ordered, or will keep the case till the reply expected is received or other action has to be taken.

In certain cases it will be seen from what follows that this general procedure is shortened or modified.

V—DUTIES OF THE REGISTRAR

18 General Supervision—The Registrar, in addition to being in charge of work of any particular branch assigned to him will generally supervise the despatch of business throughout the office. He should exercise his control by the inspection of the registers of files, miscellaneous correspondence and periodical returns and should occasionally examine a long pending case to see that no undue delay has occurred at any stage.

He should pay particular attention to the despatch of business in the copying, despatch and record branches.

on
of t
point

20 Character Rolls—The Registrar will keep, in such form as may be prescribed by the Secretary (character roll—see paragraph 11) of all applications from eligible persons for submitting and grant of leave to members of the clerical and menial staffs.

21 Dak and Routine Papers.—The Registrar will see to the distribution of the Dak and will dispose of routine papers on behalf of the Secretary and the Under Secretary in accordance with such orders as may be issued from time to time.

22 Stationery, Forms, etc.—The Registrar will maintain stock books of stationery, and will see that economy is exercised in the use of stationery, forms and service stamps (see paragraph 113)

VI—OPENING THE DAK.

23 All communications will be delivered to the officer addressed, who will initial them and note the date of receipt. Confidential communications will be sent direct to the Confidential Clerk. Other communications will be sent to the Registrar, who will sort them according to branches.

VII—REGISTRATION OF PAPERS

24 Objects—The main objects of registration are,—

- (a) to provide a check on delays in the despatch of business ;
- (b) to render difficult the deliberate suppression of correspondence in the office ,
- (c) to enable any papers lost to be readily replaced by reference to the persons with whom the correspondence has been carried on ,
- (d) to facilitate the tracing of papers in the records or elsewhere in the office , and
- (e) to facilitate the selection of records for destruction when no longer required.

25 Registration should never be entrusted to a junior clerk, responsible for dealing with the papers to be registered, except under the direct personal supervision of a Head Clerk, who will be personally responsible for the registration.

26. Registers.—Registers of correspondence will include—

- (a) a register of miscellaneous correspondence (Form A appended) ;
- (b) a register of files (Form B appended) ,
- (c) a register of periodical returns (Form C appended) ;
- (d) register of judicial cases ,
- (e) register of passport applications ;
- (f) register of applications for employment ; and
- (g) any other register the maintenance of which may be ordered by the Secretary.

Separate registers (a), (b) and (c) may be kept for each branch of the office.

27 Except under orders in cases mentioned in paragraph 31, no correspondence should be registered in two different registers

28 Numbering—Except in the case of periodicals, the first column in every register of correspondence should be a serial number which in the case of (b) will be the file number, the serial numbers should run for one year only

All correspondence relating to a case should bear a number consisting of (a) the serial number in the register and (b) letters, to distinguish (i) the branch and (ii) the register in which it is registered and (c) the year, except in the case of periodical returns for which permanent files are kept. The letters to be used to distinguish the different branches and registers are shown in Appendix II. No separate letter (b) (ii) need be used to distinguish correspondence registered in the register of files. For example, in a small office where only one register of files is kept the number to be quoted may be "65/23", in an office divided into branches the number would be "65A/23", if A is the letter used to distinguish the branch. For instance, $\frac{65 \text{ P M}}{23}$ would indicate that the case related to the Political Branch and was registered in the Miscellaneous Register. The system of numbering of letters must not be altered without the approval of the Secretary.

Note—The letters for registers and branches authorised for use in the Rajputana Agency Office are shown among the abbreviations in Appendix II.

29 Abbreviations—In making entries in registers the abbreviations shown in Appendix II and any other abbreviations which the Secretary may from time to time authorise should be used.

30 When Registration is unnecessary—Generally, every paper received in the office will be registered in some register but in certain cases registration may be dispensed with under the general orders of the Secretary, e.g., acknowledgments of papers received summonses returned and petitions which are not properly stamped and are returned accordingly.

31 Register of Miscellaneous Correspondence—A register of miscellaneous correspondence in Form A appended may be kept for unimportant communications such as petitions which are either filed or disposed of by a communication to which no reply is expected, and which do not relate to an existing file. All such papers may be kept in one or more 'Miscellaneous' files. Care must however, be taken to see that papers which relate to an existing file or regarding which the opening of a new file is or may become desirable are not treated as miscellaneous. Papers entered in this register will not be indexed in the absence of a special order to that effect.

If a reference is ordered to which a reply is expected, an ordinary file should be opened at once and the correspondence should then be registered in the register of files.

32 Register of Files—All ordinary correspondence not entered in any other register should be registered in a register of files in Form B appended. The register should be paged and one page should ordinarily be allowed for each file. If this is exceeded registration should be continued on the next blank page, forward and backward references being given.

33 Periodical Returns—A list of periodical returns due to or from the office should be prepared and each return should be given a number. A copy of the list should be hung up in every branch concerned. A permanent file should be opened for each return and it will bear the number of the return and the letter R to show that it is a permanent file relating to a periodical return, the branch letter may also be added e.g., "27 P R". In all correspondence relating to the return, this number should be used as the file number—no year need be added. Periodical returns, on receipt or despatch, should be registered in a register in Form C appended. This register may be kept in the branch concerned.

34. Opening new registers—New registers will be opened for each branch at the beginning of each calendar year, but papers should ordinarily continue to be registered in the register of the year in which the file was opened even after the expiry of that year. It may however, be desirable sometimes to carry forward registration to the register of the current year e.g., when a file becomes bulky and a fresh volume is started. In such case a note should be made in the old register showing where the registration is continued, e.g., 'continued at page 95 of 1924 register'.

35 Central Record Room—If a single record room is kept for all or several branches, a separate register of files should be kept for each branch.

✓ VIII—INDEXING

36 Object—An index of records and pending files must be kept. The main object of the index is to enable all the previous papers in the record or elsewhere in the office relating to a specified subject or person to be traced and submitted as quickly as possible. The index should be so devised that no previous papers relating to the matter in hand are accidentally overlooked.

37. Card or loose-leaf Index—The index should be in the form of card index or loose leaf index. It will be assumed in the following description that card index is used but the procedure is similar if a loose leaf index is used, as may be found convenient in certain cases.

38 Major and Minor Heads—Great importance attaches to the selection of proper subject headings. The index showing major and minor heads for use in the Rajputana Agency as approved by the Secretary is printed separately and no addition may be made to the list without the Secretary's sanction.

list. It is
General sh

respect of matters on which there may be correspondence with the Agent to the Governor General, this will facilitate the indexing of papers in the office of the receipt.

39 Cross References—Every file must be ordinarily indexed under approved major and minor heads as soon as it is opened, but cross reference under other heads may be made at the discretion of the record keeper or head clerk. Sub heads must be used to keep the subject of each file as definite as possible (see paragraph 49).

40 Cards.—Major heads must be written or type-written in block or capital letters at the top of a card the minor head on the next line and the sub head on the following line. The file number (including the branch letter and year) should be entered on the right of or below the sub head. When the card is intended to make a cross reference to another head the head from which the cross reference is to be made will be entered in block or capital letters at the top of the card and the major minor and sub heads to which the reference is to be made will be entered on the next three lines with the word "see" before the major head, in this case no file number need be entered. In an index of names the name should be written or type written in block or capital letters on the first line further particulars, such as designation father's name or address, should be given in the second and following lines to enable the person named to be distinguished from others bearing the same name.

41 Branch Index.—In a small office or branch where the record keeper's index is readily available for the use of the clerks, a single index will generally suffice in a large office with a central record room not readily accessible to all the clerks a duplicate index should be kept in each branch of the files relating to it. In the e rule it is assumed that a duplicate index is kept, where this is not done references to the duplicate index should be disregarded.

42 Pending Cases.—In both indexes the cards relating to pending cases should be kept in a separate portion of the index from those relating to closed files. The branch clerk will be able to determine from the card index whether it is necessary to open a fresh file or not on the receipt of a communication in which there is no quotation of a pending file number. When a file is disposed of, the index card or cards relating to it should be transferred to the portion of the index which relates to closed files.

43 Subsidiary Indices.—The index may be divided into portions by means of guide cards. Besides the general index (1) of closed files and (2) of pending files it may be found convenient to have an index of names. Separate indices for each State are not necessary. If a name-index is kept names may be excluded from the main or subject index, if a subject index is kept for a State, additional cards will have to be made for the State subject index as the subject matter also be included in the main subject index.

44 Where Indexing is unnecessary.—In certain cases it will not be necessary to index records at all e.g., the records of judicial cases which should be kept in the record room in serial order of numbers of cases for each class of case. It may not be worth while to make an index of applications for permits under the Arms Act, the applications and connected papers should be kept in the record room in serial order of numbers in the register of such applications, if an index is necessary it should be an index of names. It is not necessary to index periodicals, a file for each periodical should be kept and these files should be arranged in the record room in a special place according to the numbers of the periodicals.

The Secretary will issue general orders as to the classes of cases which need not be indexed.

IX — ARRANGEMENT OF A CASE

45 Single Case — A case for orders will be submitted in a file board. The papers should be arranged in the following order beginning from the bottom —

- 1 Files put up for reference in order of number and year—the oldest files being in the bottom
- 2 The file of the case
- 3 Draft if any
- 4 Paper under consideration
- 5 Notes
- 6 Outside the flaps under the tape any books plans etc put up for reference

The papers in the file should be arranged in chronological order so as to read forward like a book. The P U C should be separate from the file but as soon as the draft has issued or other action been taken the paper under consideration and the draft if any should be placed in the file in chronological order.

46 Binding of Files — Care should be exercised in binding papers into a file that the whole of the paper is easily legible without undoing the file lace. In the case of half margin drafts or office copies which are duplicates of

47 Linking Cases — Cases can be linked when it is desired to refer in connection with one case for orders to files which are already included in another case which has not been disposed of so as to avoid breaking up either case or when two cognate cases are put up together for orders.

X — FIRST ACTION IN THE BRANCH

48 New Subjects — If a receipt does not obviously relate to a pending case the branch clerk should ascertain by reference to the index of pending cases

attached
either the
the head
clerk to

enter the receipt in the Register of Miscellaneous Correspondence (see paragraph 31)

49 Great care should be taken to keep the subject of each file as definite as possible, any tendency to place on one file a large number of receipts dealing with different aspects of a general subject should be checked as this will make the file unwieldy and impede business. It is better to have a larger number of files each with a more limited subject, new files should be freely opened when desirable. Appendix III contains some general instructions regarding the opening of files.

M2AGGR

50 If the head clerk decides that a new file should be opened or orders to that effect are received in the Register of Miscellaneous Correspondence he will enter at the head of the receipt in red ink the major minor and sub heads under which it is to be indexed and underline, in the minor and sub heads any heads under which, in his opinion it should be cross-referenced in the index, he must not use any unauthorised major or minor head without the permission of the Secretary which should be obtained, if necessary through the record keeper and Registrar

51 The branch clerk will prepare index cards and a file cover (in Government of India Standard Form S 97) accordingly and attach the receipt to the file cover by a tag through the left hand top corner and after consulting the index will note on the file cover any previous files which he requires he will take or send them all in a file board to the record keeper. The latter will register the receipt and enter the file number on the file cover on the index cards (other than those for cross references) and at the top of the receipt, he will place his index cards in their places in his index put up the previous files asked for or a slip showing in what pending cases they are to be found or stating that they

the branch clerk and initialled by them respectively whenever the pending file is transferred from one to the other

52 Pending Subject —(a) When a receipt relates to a pending case and pending cases are kept in the branch the head clerk will before giving the receipt to the branch clerk obtain the case and register the receipt by entering

January 1921' an abstract of the receipt should be entered in the notes

(b) If the register of files is kept in the branch and not by the record keeper the Head Clerk will register the new receipt in it or have it registered under his direct personal supervision before giving the receipt to the branch clerk,

(c) in other cases the new receipt should be sent or taken at once by the branch clerk to the record keeper, who will register it in the register of files, the record keeper will return the receipt to the branch clerk, he will at the same time give the branch clerk the pending case if he has it, taking the latter's initials in the slip for the case

The branch clerk will add the receipt to the file and number its pages in continuation of the last page number in the file. A cross reference should be added at any time before a file is closed if a receipt renders it desirable

If the receipt completes the replies due to a reference, the branch clerk will score out the file number in his reminder diary against the date on which the next reminder was to have been issued this date is given on the margin of the draft of the reference (cf, paragraph 115)

53 Letters from Princes—When a letter is received from a Ruling

amine the question raised and to send a further reply

54 Printed copies of letters received—When a printed copy is received of a communication previously received in manuscript the printed copy should be substituted for the manuscript copy which should be destroyed

XI—REFERENCING

55 "Referencing" defined—The object of referencing is to enable papers referred to or likely to be referred to by persons dealing with a case to be easily consulted, it includes the following processes—

- (a) Putting up any previous correspondence likely to assist in the disposal of the paper under consideration
- (b) Noting in the margin of the paper under consideration and of the notes where any previous paper referred to in them can be found,
- (c) "Flagging" a paper to which reference has been made in the paper under consideration or in the notes or is likely to be made in the notes

In a marginal note made under clause (b) above the file number and page of the file as well as the flag attached to it, should be quoted (e.g. '65A/23 page 12 fl L') unless the information is already given in the letter press

56 New File.—When a new file is opened, the branch clerk will as already described obtain from the record keeper any previous files in the record room which appear likely to assist in the disposal of the paper under consideration if any file likely to be wanted is put up with another pending case the branch clerk should obtain that case and link it temporarily with the case under consideration

The branch clerk will examine the files put up and remove any which are not wanted, obtaining if necessary, the verbal orders of the head clerk before doing so, any files so removed should be returned at once to the record keeper. The branch clerk, as he goes through the files should flag any paper referred to in the P U C, or to which he thinks it likely that a reference will be made

57 Additions to File—Every paper added to the file at any stage must be at once 'referenced' by the branch clerk, such papers include notes, drafts and further correspondence

XII—NOTES

58 Object—The object of the notes written by the clerks is to make it easier for the officers dealing with the case to arrive at correct decisions on the questions raised in the correspondence. No note should be submitted which does not have this effect or which is not likely to be read by the officer concerned

59 **Contents.**—The notes will vary in different cases; in a simple case no note at all may be required. In a complicated case the note may contain all or any of the following —

- (a) a précis of previous history or correspondence;
- (b) a verification of facts alleged in the P. U. C.;
- (c) the points for decision;
- (d) a verification of the correctness of references to Acts, rules, previous correspondence, etc., in the P. U. C. and of their interpretation as bearing on the points for decision;
- (e) further references to Acts, rules, previous correspondence, etc., bearing on the points for decision; and
- (f) precedents.

The occasions *however* on which a full note of this description will be required will be rare. Notes, however, should be as brief as possible.

60 **Incorrect Notes.**—The following should never occur in notes:—

- (a) *For reference to the previous correspondence or from*

- (d) Notes by (1) the branch clerk, (2) the head clerk, (3) the Secretary and (4) the Agent to the Governor General,
- (e) Branch and head clerk's note, "Draft submitted",
- (f) To whom a reference is made and date (red ink) *e.g.*, "To—P O, M W R, 18th December 1923",
- (g) From whom a reply is received and date (red ink) *e.g.*, "From—P O, M W R, No 75/23, 18th December 1923",
- (h) Further notes by the branch clerk, etc

64 **Marginal Notes**—Marginal notes should not be written by any one below the rank of Registrar except in brief reply to a question, in other case the passage should be marked in the margin by a line and letter and the clerk should deal in a further note below the last notes with "the point at (letter)" the Secretary's note dated "

65 **Notes to be initialed**—Every note must be initialed and dated by the writer. The date must include the year. Initials below the previous note without comment indicate acceptance of the correctness of the previous note and approval of any suggestions for action made in them.

66 **Branch Clerk's notes**—The branch clerk should note only in accordance with the general or special instructions (to be obtained verbally) of the head clerk. He should not attempt to suggest the disposal of a case, unless it is of a routine nature and in accordance with well established precedent.

67 **Head Clerk's note**—The head clerk should delete from the branch clerk's note anything that is wrong or unnecessary and should correct it and have it copied as corrected, if necessary. He may of course supplement it by a further note but should be careful not to repeat what is in the branch clerk's note. The head clerk should not discuss questions of policy or administration or suggest how cases of complexity or importance should be disposed of, in ordinary

which he considers unnecessary

68 **Further duties of Branch Clerk**—The branch clerk must at every stage see that every note is 'referenced' and that the notes are paragraphed, this must be done for example, before the head clerk's note is submitted to the Secretary and before the Secretary's note is submitted to the Agent to the Governor General. Before a case is submitted to the Agent to the Governor General or Secretary, the branch clerk will see that there are one or two blank sheets of paper at the end of the notes for the use of the office concerned.

69 If in doubt, head clerk or branch clerks, with the permission of the head clerk, should consult the Secretary before noting

78 The letter form should be used in addressing Darbars and when the subject matter cannot conveniently be expressed in memorandum form, e.g., when an argument has to be developed at some length

79 The memorandum form is designed to secure economy and expedition and communications in this form should be in brief telegraphic form. It is suitable for straightforward statements of facts or orders

80 Endorsements—The endorsement form should be used when transmitting copies for information or with brief instructions but it should not ordinarily be employed in addressing unofficial persons other than petitioners. Sample endorsements and memoranda on petitions will be found in Appendix IV. Routine memoranda and endorsements to Political Officers, etc., may issue under the attestation of Registrar after they have been passed by Secretary

81 Telegrams—Telegrams should be used as sparingly as possible. It should always be considered what saving in time will be effected by sending a telegram instead of a letter. A telegram received after office hours has generally

The wording of tele-
the meaning is clear
inary' telegram will

suffice

82 When the purport of a general communication has to be sent to Political Officers for the information of Darbars, it should where possible be drafted in the form of a note and despatched with a covering letter. Sufficient copies of the note should be sent to each Political Officer for the information of each Darbar

83 Demi-official Letters—Demi official correspondence may be used, where necessary, to supplement or explain official correspondence or to take the place of official correspondence in cases of extreme secrecy or great urgency

Demi official letters should be included in the file in their places in chronological order. A demi official letter received should, if necessary, be pasted on to a foolscap sheet. Demi official correspondence should not be referred to in official correspondence

84 Unofficial References—An unofficial reference consists in marking

1

in respect to which the reference is made should be stated as precisely as possible. An unofficial reference must not be made without the permission of an officer not below the rank of Registrar

85 Preparation of the Draft—(a) In simple cases or where there is established precedent or there is little doubt as to the message to be sent, a draft and fair copy for signature should be submitted with the case in the first instance, this practice should be adopted as much as possible, what is in the draft need not be repeated in the note which may generally be very brief, e.g., "Draft submitted, for precedent see No. 21/22, page 6", this practice saves

trouble and delay in disposal as the officer to whom the case is submitted can at once deal with the draft and if he approves of it with little or no alteration sign the fair copy and will be saved the re-examination of the case, which is necessary when a draft is submitted after orders have been passed on the notes.

(b) In other instances the case may be returned by the Agent to the Governor General or Secretary with a draft (or with a fair copy already signed and a copy initialed for the file) in such cases the head clerk must examine the draft or communication and if he considers that any portion of it is based on a misapprehension or that any point for decision has been overlooked or that a draft passed by the Secretary should under the general instructions of the Agent to the Governor General have been submitted to the latter it is the head clerk's duty to bring the matter to the notice of the Secretary at once either personally or by a separate note placed at the top of the case.

(c) When the orders passed on the notes necessitate the preparation of a draft in the office the head clerk will decide whether he will prepare or dictate the whole draft himself or entrust the preparation of the whole or a portion of it to the branch clerk. His decision should depend on the complexity of the case and on the abilities and experience of the branch clerk but he should train the branch clerk in preparing drafts.

If there are any doubtful points in the orders on the notes, the Secretary's instructions should be obtained at once verbally if possible.

86 Form of Draft—A draft of an official reference should ordinarily be written half margin on a separate foolscap sheet. When a fair copy is submitted with a draft under paragraph 85 (a) and the draft is made by carbon paper at the same time as the fair copy, the draft need not be half margin.

87 At the head of every draft of an official letter memorandum or endorsement should be put the number to be placed at the head of the fair copy, e.g. 60-1/23. In the case of a telegram this number should be placed at the beginning of the message.

88 At the beginning of every letter or memorandum to an official should be placed first the subject as entered in the file register and then the number and date of any communication to which it is a reply, e.g.,

' Sir,

Allowance—Grain Compensation Menials

Your letter 773 C/23 dated the 20th September 1923'

This form facilitates the distribution of the Dak in the office of receipt

89 Enclosures—When enclosures have to accompany a communication the drafter should enter a long list of enclosures including the margin of the draft in red ink or red pencil against the entry showing the list of enclosures as a guide to the copying and comparing clerks and to the despatcher.

90 Copies of Enclosures—When it is intended that a Political Office should communicate a copy or copies of a communication or of its enclosure

to Darbars or other authorities, a sufficient number of copies of the communication or enclosures should be sent with the communication to save the trouble of having fresh copies made in the Political Officer's office. In such cases the branch clerk must note the number of copies of the enclosure to be sent at the end of the draft for the instruction of the copying and comparing clerks and of the despatcher.

91 Addressees—A list of persons to whom the communication is to be sent must be placed at the head of the draft. Authorised abbreviations should be used (Appendix II).

92 First Reminder—At the end of the draft in the margin above the list of addressees, if a reply is necessary to the communication the drafter should enter the date on which he thinks that the first reminder should issue, if a reply is not received by that date, e.g., "1st Rem 2nd February 1924".

93 Reference in Reply—When a reply is sent to a communication regarding which one or more reminders have issued the reply should quote the date of the original communication and not that of the reminder, e.g., "with reference to your letter No 71 P/23, dated the 17th March 1923 and subsequent reminders".

94 Place in Case—A draft when ready should be put in its proper place in the case and the whole case should be submitted to the Secretary for orders.
 ' ,

the Governor General or not

XV—FAIR COPYING

95 Definition—A "fair copy" means a copy of a communication to be sent out of the office.

96 When Fair Copy to be made in Branch—When no draft is necessary, or when the fair copy is submitted with the draft the fair copy will be prepared by the branch clerk and submitted by the head clerk direct for signature.

97 When draft sent to Copying Branch—In other cases, after a draft has been passed, the branch clerk will—

(a) if a reply is expected, note the file number in his reminder diary against the date on which the first reminder is to be issued, and

(b) remove the draft from the case and send it to the copying branch.

If he desires to have a clean copy for the file in place of the draft as passed he must give instructions accordingly. He should ordinarily obtain a clean copy if the draft is in manuscript.

If copies of papers in the case are to be enclosed, the case or file concerned should be sent to the copying branch with necessary instructions.

98 Date—The copying clerk will enter the date above the draft and below the File No., this will be the date of the communication and will be entered as such in the fair copy.

MAGGR

99 Number—The number at the head of the draft should be entered at the head of the communication, *e g*, "No 65 A/23"

100 Comparing—After the copies have been made, they should be compared by the head clerk or a branch clerk. The clerks copying or comparing fair copies should initial all the fair copies (including any to be kept in the file) in token of their having done so. The fair copies of all important letters should be checked by the Secretary.

101 Signature—The comparing clerk will send the fair copies for signature in a portfolio. Any fair drafts should be submitted for initials at the same time as the fair copy. Letters to be sent to the Government of India, a local Government or the General Officer Commanding a Division should ordinarily be submitted to the Agent to the Governor General for signature. The Secretary will issue such instructions as may be necessary as to the

a fair draft has been initialled by the officer signing the fair copy, the original rough draft may be destroyed at once by the comparing clerk.

XVI—DESPATCH

102 Papers to be sent to Despatcher—All communications to be despatched from the office should be sent to the despatcher. When a fair copy is signed by the Registrar or any subordinate officer, it will be sent direct, after signature by him to the despatcher with any papers put up with the fair copy. A fair copy signed by a higher officer will be returned to the clerk who submitted it, this clerk will at once send to the despatcher the fair copy with the draft if there is one or with a register or note in which the despatch can be noted. In the case of papers registered in the Register of Miscellaneous Correspondence the register should be sent with a note in the last column as indicated in paragraph 76.

103 Return of Office Copies, etc—The despatcher, after verifying the correctness of the papers (including any enclosures to be sent) will mark at the end of the draft or in the register or note by means of a rubber stamp, the word "Despatched" and the date. He will then return to the branch concerned all the papers except the fair copies to be despatched. Papers relating to files of which the record keeper keeps the registers in Form B should be sent through him so that he may enter the same in his register.

104 Telegrams, etc—The despatcher will be responsible for the prompt despatch of telegrams and of papers marked "urgent" for local delivery.

105 Hours of closing Desk to be fixed—The Registrar will fix hours at which messengers should ordinarily be sent out for local delivery, every messenger must have a book in which the despatcher will enter the name of every addressee and the number of packets sent to each, on the messenger's return the despatcher must at once examine the book to see that all covers sent out have been correctly delivered and duly acknowledged.

106 The Registrar will fix hours at which the despatcher will close packets for delivery by post in different directions. Different hours may be fixed for registered and other letters.

107 The hours fixed under the preceding paragraphs should be included in a notice to be hung up in every branch

108. **Accumulation of Papers**—Papers received during the day for despatch will be placed by the despatcher in appropriate pigeon holes pending the closing of the dāk, unless they are addressed to petitioners, when they may be put in covers, addressed and stamped at once

109 **Closing Dāk**—At the time fixed for closing the dāk the despatcher will place all the papers for the same person in one cover unless they are too bulky

110 **Economy Labels**—Economy labels should be used for ordinary non confidential communications to officials

111 **Registered Post**—If papers are to be sent by registered post the clerk will personally take them to the despatcher and give him necessary instructions, the despatcher will enter in manuscript after the entry "Despatched" the word "Registered" (and "Ack Due" if an acknowledgment is to be obtained) and initial this entry. The despatcher will keep the postal receipts for registered letters in chronological order for six months and then destroy them

112. **Stamps.**—When all the covers are ready for the post, the despatcher will mark the date of despatch on each cover with his rubber stamp just below
 sary entries in his Stamp Register (Form
 and the Register to the Superintendent,
 correct, he should check any wastage of
 stamps or the employment of unnecessarily large covers. The Superintendent should also see that the value of stamps issued to the despatcher is entered in the latter's Stamp Register

113 The stock of service stamps should be kept under lock and key under the charge of the Accounts Clerk, who should maintain an account showing all receipts of stamps from the Treasury and disbursements made to the despatch clerk. The despatch clerk should also keep the stamps under lock and key and issue them in small quantities to the Daftri, who should

!

 clerk should be made primarily responsible for this check. The Accounts Clerk should exercise supervision and check the issues and balance on hand at least once a month at irregular intervals

114 **Further Action by Branch Clerk**—On receipt of the draft, etc., from the despatcher, the branch clerk will—

- (a) add the draft to the file;
- (b) page it in continuation;
- (c) note its despatch in the notes in red ink, see paragraph 63 (f);
- (d) if the register of files is maintained in the branch in Form B appended, register the issue or have it registered;

- (c) take or send the case to the record keeper, if he keeps pending cases, or otherwise put it in its place in the serial order of file numbers among the pending cases of the branch

XVII — REMINDERS

115 Diary.—Every branch clerk will keep a diary, this will consist of three fool cap sheets stitched lengthwise in the middle so as to form 12 pages, each page being half a fool cap sheet, one page will be allotted for each month of the year. The pages should be ruled so as to give one line for every day of the month. The object of this diary is to ensure and facilitate the issue of reminders.

As directed in paragraph 97 (a) the branch clerk will note in the reminder-diary the file number against the date on which the first reminder is to be issued. If all replies are received before that date the branch clerk may score out the entry of the file number in the reminder diary against the date relating to the reminder.

116 Issue of Reminders.—Every morning the branch clerk will ascertain from his reminder diary on what files he should issue reminders on that date. He will at once obtain the cases mentioned. If the date on which a reminder should be issued according to diary is a Sunday or public holiday, the reminder may be issued on the next working day. Except where the reply is due from the Government of India the branch clerk will prepare necessary reminders for signature (no office copy should be kept) and at the same time note in the margin of the draft under the entry relating to the reminder issuing the date on which the next reminder should, in his opinion, issue e.g. '2nd Rem 2nd March 1924'. The case should then be submitted through the head clerk to the Registrar with the reminder or reminders on top and immediately below them the page of the draft on which the dates of the reminders are entered.

117 Signature and Despatch.—The Registrar, after signing the reminders and initialling against the entry relating to the reminders issued in token of his signature and of his approval of the date on which the next reminder is to be sent will send the case to the despatcher who will take out the reminders for despatch and mark the date of despatch with his rubber stamp against the entry relating to the reminder on that draft. The despatcher will then return the case to the branch concerned.

118 Subsequent Reminders.—The branch clerk will note in his reminder-diary the file number against the date on which the next reminder is to be issued, he will then return the case to be kept among the pending cases.

The same procedure should be followed with reference to subsequent reminders.

119 Reminders to Government of India.—It is unusual to send official reminders to the Government of India in the absence of special reason for so doing which should be explained in each case. If a reply is not received from the Government of India by the date on which it was expected that first reminders should be issued the case should be submitted to Secretary for orders.

120 Further Uses of Diary.—The reminder diary may be used for noting other files on which action is to be taken on a particular date, *e.g.*, if special report has to be submitted on a particular date the connected file should be noted against a date sufficiently early to enable the report to be prepared in time

121 Periodical Returns—The diary may also be used for noting dates on which action should be taken in connection with periodical returns, *e.g.*, by the issue of advance reminders or reminders or by the compilation of information in the office, only the number of the periodical need be entered against the date on which action is to be taken, *e.g.*, '27 P A'.

XVIII —CLOSING FILES

122 When File should be closed—A file should be closed as soon as the question raised when the file was opened is decided. This will generally occur when a reply has been received to a reference which has been issued or a final reply is issued from the office

123 Orders to close—If the branch clerk thinks that the file should be closed, he will enter the word 'Record' on the last page of the file below the last communication or draft and will submit the file to the head clerk. If the head clerk agrees that the file should be closed, he will initial the word "Record"

124 Procedure in closing—The branch clerk will then take the following measures to close the file —

- (a) He will remove and place together all rough drafts, drafts not approved, duplicate copies, routine slips reminders, replies to reminders etc, which should immediately be destroyed. He should also remove for future use any blank sheets attached to the notes which have not been used
- (b) If he keeps a card index of files, he will transfer the cards relating to the file (including cross references) to the portion of the index relating to closed files.
- (c) He will, if necessary, divide the file into parts if one portion of the file should under the rules relating to destruction of records, be kept longer than the other portion (see Appendix V)
- (d) He will note on the file-cover (Government of India Standard Form S 97) the dates on which the file or each part of it should be destroyed (Appendix V)
- (e) He will arrange each part, the notes first and the correspondence second, the notes should either be bound in with the correspondence or, if bulky, bound in a separate file cover marked with the file number, the word "notes" being added. They should not be kept loose
- (f) He will then obtain the approval of the head clerk who will signify it by initialling on the cover below the entry regarding destruction

- (g) The head clerk will then consider whether the file should be indexed under any other heads than that on the file cover and if so direct the branch clerk to prepare the necessary cards.
- (h) The branch clerk will take or send the whole case to the record keeper, with any new cards for the record-keeper's index.
- (i) He should obtain from the record-keeper the slips for the case and for previous files now returned by him.

125. The record keeper will proceed as directed in paragraph 131.

XIX — RECORD-KEEPER.

126. **Responsibilities.**—The record keeper is responsible for the maintenance of the register of files, for his index of files, for the safe custody, preservation and dusting of records and for their destruction in accordance with the rules on the subject. He must also comply with requisitions for records with promptitude. No unauthorised person should be allowed access to the records.

127. **Pending Cases.**—Unless a central system of records is in force, the record keeper may also take charge of pending cases, he should keep these unbroken apart from the ordinary records.

128. **Closed Files.**—Records of closed files must be kept in serial order in bundles of convenient size each bundle being between boards of wood or stiff card boards bound by two tapes, on the top board of each bundle should be noted the numbers of the files which are to be included within it, e.g.,—

" Branch B

1928.

Nos. 601 to 700".

The bundles must be arranged on shelves and should be piled so that the space between two shelves is nearly filled. On the outside of each shelf below each pile of bundle should be entered the year and the numbers of the files on that portion of shelf, e.g. "601 1000 B-1928."

129. **Files taken out.**—When any file is taken out for reference from the records, a slip showing the file in connection with which it is required and initialled by the branch clerk, must be put in its place. This slip should be returned to the branch clerk and may be destroyed by him when the file is put back in its place. The record keeper will also note on the cover of the file the files which are put up for reference in connection with it.

130. **Opening a file.**—When a file is to be opened, the record keeper will receive the paper in a file board from the branch clerk with the index cards; the record keeper will register the file in the register of files and enter the file number on the paper under consideration and on the cards on which the new file is indexed. He will then put in the case any previous files in the records for which the branch clerk has asked or, if these are not in the records, put in a slip showing in what pending cases they are to be found if they have not been destroyed. He will then return the case to the branch clerk with the latter's index card and the necessary slips for files issued to be initialled and returned by the latter, these slips should include a slip relating to the pending file and headed "pending file"; this slip should be exchanged between the record keeper and

the branch clerk and initialled by them respectively whenever a pending file changes hands

131 Closing a File—When a file is closed the record keeper on receipt of the case will verify that the file is in order and will return to the branch clerk the slips for the file and for any previous files put up in the case the previous files should be at once returned to their proper places in the records the record keeper will have the newly closed file stitched and paged and will note the number of pages in each part on the file cover he will note in the register of files the date of closing the form of disposal and the date of destruction of each part He will also enter the date of destruction in his destruction register (paragraph 133) He will at the same time put a circle round the number in the first column of the register e.g. (65) to indicate that the file is closed

The index cards relating to the file should be transferred from the portion of the index relating to pending cases to that relating to closed files the file should be put in its place in the records

132 Reopening closed File prohibited—A file once closed must not be re opened without the permission of the Secretary but such papers as acknowledgment of receipt of communications which do not involve further correspondence may be added to file after it is closed

133 Destruction of Records—The record keeper is responsible for the destruction of records when it is due For this purpose he will maintain a destruction register showing under the appropriate month and year the files

Secretary with the list for orders as to their destruction. If in any case the Secretary considers that a file should be retained for a longer period the destruction register the register of files and the entries on the files must be altered accordingly When a file has been destroyed the destroying officer will enter the date of destruction in column of the register and will also put a cross over the file number in the first column e.g. (60) The record keeper will note in his card index the fact that file has been destroyed by noting by the rubber stamp *Dest (date)* on the card if all the files on the card have been destroyed it may be the card index relating to the card index records is kept in records to the branch clerk so that similar action may be taken there

The record keeper will also carry out similarly the destruction of periodicals returns judicial case records and other papers which are ripe for destruction but are not kept in ordinary files

The destruction of records for the year should ordinarily be completed in the first quarter of the year A report must be submitted by the record keeper through the Registrar to the Secretary at the beginning of April.

XX—DUTIES OF CONFIDENTIAL CLERK.

134 The Confidential Clerk will be the only person in the office under the Secretary to deal with confidential and secret papers He will maintain registers of such papers in forms A and B and a reminder-diary and will keep in

own custody all confidential and secret records and an index of them. When however, such a course appears desirable and unobjectionable he may send to the record keeper an index card relating to a confidential paper so that, if the matter is referred to in ordinary correspondence, the confidential file may not be overlooked. The Confidential Clerk will also be responsible for the destruction of confidential and secret records under the orders of the Secretary. He may also obtain the Secretary's orders to transfer confidential papers to the general records when there is no longer any adequate reason for keeping them confidential, such records should be acknowledged by the record keeper and should be kept in serial order in special place in the record room and should be indexed.

135 The Confidential Clerk will copy and despatch all confidential and secret papers to be sent out of the office, such papers should be put into double sealed covers the inner cover only should be marked 'confidential' or 'secret' and superscribed with only the name of the addressee, the outer cover should bear his usual official address.

XXI — MISCELLANEOUS

136 Check on disposal of business.—All branch clerks will submit once a week a list of cases which have been pending with them for more than six days. This list will be submitted through the head clerk and the Registrar to the Secretary.

137 that it is for disposal of for this purpose, the record keeper may clip together, by means of a paper fastener in one corner, the pages of the register on which there are no files pending.

138 The Registrar should examine all the registers in the office for the current year at the beginning of each month and submit them to the Secretary with his remarks when the latter is at Headquarters. These registers should be returned as soon as possible to the record keeper concerned, any note from the Secretary asking for any pending files being attended to separately.

139 Interviews.—If there are any papers regarding a person to whom the Agent to the Governor General grants an interview, they should be submitted to him on the day before the interview is to take place.

140 Corrections.—When slips are pasted in files books or registers to cover untidy or cancelled writing they should be initialed by the clerk doing it and by the head clerk.

141 Signature and initials to be dated.—All signatures and initials should be dated, date includes the year.

142 Stationery.—Stationery will be supplied at the beginning of each month to each branch on a separate list. Stationery or the Secretary responsible Registrar will refuse to purchase or issue requisitions from head clerks.

143. **Furniture.**—The furniture of the Agency Office is in the charge of the Registrar. Proposals for the purchase of new furniture or for renewals of existing stock will be submitted for the orders of the Secretary. A stock book will be maintained by the Registrar in which all new purchases must be entered; no stock should be written off without the Secretary's permission. The Registrar will check the stock in April every year and report the result to the Secretary.

144. **Tents.**—The tents and camp equipage will be in the charge of the Secretary. A stock register will be maintained by the Mir Munshi. The Secretary will inspect the stock once a year.

or damage.

XXII.—TOUR ARRANGEMENTS.

145. **Modifications in procedure.**—The Secretary will decide what modifications in procedure are necessary owing to his absence or to the absence of any clerk, etc., on tour.

146. **Camp Register.**—When the Agent to the Governor-General is absent from Headquarters on tour, a camp register will be opened in which all correspondence received in or issued from camp will be entered. The serial numbers in this register will be independent of numbers in the registers at Headquarters. When a new file is opened in the Camp, a number from the Camp Register will be assigned to it, which, with the subject of the file, will be communicated to the Headquarters office. The Headquarters office will then register the file after the last serial number in the File Register, and communicate it to the Camp Office who will replace it by the number intimated. The Camp Register will be destroyed on the return of Camp, after comparing the entries with the Headquarters Register. For facility of reference, the number intimated by the Headquarters' office will be noted against the Camp numbers which have been replaced by them.

147. **New File.**—A memorandum will be sent daily to Headquarters of all receipts in and issues from Camp Office. The details will be entered in the headquarters register and the memorandum will be returned to the Camp Office with the word "entered" and the signature of the person who has made the entry. The letters 'E. H. Q.' ('entered in headquarter register') will be placed against the entries in the Camp register.

148. **List of Files.**—Whenever cases are sent to Headquarters or *vice versa*, a list showing the file numbers of the cases should be sent at the same time, an office copy made by carbon paper being kept in the office of despatch. The list should be signed on receipt after check by the clerk responsible for opening the Dāk and should be returned to the office of despatch. The office copy may then be destroyed.

APPENDIX I

FORM A

REGISTER OF MISCELLANEOUS CORRESPONDENCES
(Paragraphs 26 (a) 31, 48 and 134 and Appendices II and VI)

1 Serial	2 From whom	3 Outside number and date	4 Purport.	5 Office Note	6 Orders	7 Action taken.	8 Remarks
1	2	3	4	5	6	7	8

ALTERNATIVE FORM B.

(1) REGISTER OF FILES,
(Paragraph 26 (b), 32, 33, 130, 131, 131 and Appendices II and VI.)

File No. 1	Date of opening. 2	Subject (Major, minor or sub-head) 3	Date of closing 4	R or D (year) or N. 5	Record keeper's initials. 6	Year of destruction 7	Initials of destroyer. 8

[See Government of India Standard form S 97]

FORM D

FILE COVER (FRONT PAGE)

[Paragraphs 51, 124 (d) and Appendix VI].

Office.

Branch ()

Year

File No.

Subject.

Retain	{	Part I.
		Part II.

Destroy	{	Part I in
		Part II.

FILES PUT UP FOR REFERENCE

Previous references.

Later references.

APPENDIX II

ABBREVIATIONS

[Paragraphs 28, 29, 52, 63, 91 and Forms A and B]

Agent to the Governor General	A G G
Chief Commissioner	C C
Resident, Mewar	Rest, M W I.
Resident, Western Rajputana States	Rest J P P
Resident, Jaipur	Rest, J P R
Political Agent, Eastern Rajputana States	P A, E R S
Political Agent Haraoth and Tonk	P A, H & T
Political Agent Southern Rajputana States	P A, S R S
Political Superintendent, Hilly Tracts, Mewar	P S Kherwara
Assistant Political Superintendent, Hilly Tracts, Mewar	A P S Kotra
Commandant Mina Corps	Comdt M C
Commandant, Mewar Bhil Corps	Comdt, M B C
Assistant Commandant Mina Corps	Asstt Comdt, M C
Assistant Commandant, Mewar Bhil Corps	Asstt Comdt, M B C
Principal, Mayo College, Ajmer	Pr, Mayo
Chief Medical Officer in Rajputana	O M O
Secretary, Public Works Department	Sec, P W D
Inspector General of Police	I G P
Railway Magistrate Ajmer	Ry Magt
Commissioner Ajmer Merwara	Comr
Assistant Commissioner, Ajmer Merwara	Asstt Comr
District Magistrate, Abu	D M, Abu
Foreign and Political Minister Bikaner	F & P Minister
Chief Minister, Sirohi	C M S R I
Administrator, Jhalawar	Adm, J H R.
Postmaster General, Central Circle, Nagpur	P M G, Nagpur
Accountant General, Central Revenues Delhi	A. G, C R, Delhi
Government of India, Foreign and Political Department	F and P
Government of India, Home Department	Home
Government of India Department of Industries and Labour	Industries
Government of India, Department of Commerce	Commerce
Government of India, Department of Finance	Finance.
Government of India, Department of Army	Army
Government of India, Department of Education, Health and Lands	Education
Excise Commissioner for Central India	Exc Comr, Indore
Secretary to the Agent to the Governor General, Central India	Sec., Indore
Agent, B. B & C. I. Railway	Agent B. B & C. I.,

Traffic Superintendent, B., B. & O. I. Ry.	T. S., Ajmer.
Superintendent, Government Printing, India..	S. G. P., Calcutta.
Controller of Printing, Stationery and Stamps	C. P. S., Calcutta.
Political Agent, Palanpur	P. A., Palanpur.
Superintendent of Education, Delhi and Ajmer-Merwara	Supdt., Edn., Delhi.
Chief Secretary to the Government of United Provinces	U. P.
Chief Secretary to the Government of Punjab	Punjab.
Chief Secretary to the Government of Bengal	Bengal.
Chief Secretary to the Government of Central Provinces	C. P.
Chief Secretary to the Government of Madras	Madras.
Chief Secretary to the Government of Bihar and Orissa	B. and O.
Chief Secretary to the Government of Burma	Burma.
Chief Secretary to the Government of Assam	Assam.
Agent to the Governor-General and Chief Commissioner, Baluchistan	A. G. G., Baluchistan.
Agent to the Governor-General, North-West Frontier Province..	A. G. G., N. W. F. P.
Director-General, Indian Medical Service	D. G., I. M. S.
Director-General of Observatories	D. G., Observatories.
Director-General, Commercial Intelligence	D. G., Com. Int.
Director-General of Posts and Telegraphs	D. G., P. & T.
Controller, Military Accounts	C. M. A.
Bharatpur	B. R. P.
Alwar	A. W. R.
Karauli	K. R. I.
Dholpur	D. O. P.
Kotah	K. T. H.
Jaipur	J. P. R.
Kishanganj	K. S. G.
Jodhpur	J. D. P.
Jaisalmer	J. S. M.
Tonk	T. N. K.
Bundi	B. N. I.
Mewar	M. W. R.
Dungarpur	D. N. P.
Partabgarh	P. B. G.
Banswara	B. N. W.
Bikaner	B. K. R.
Sirohi	S. R. I.
Jhalawar	J. H. R.
Khushalgarh	K. L. G.
Shahpura	S. H. P.

Registers—

Miscellaneous	M.
Periodical Returns	R.
Judicial cases	J.
Passports	PP.
Applications for employment		E.

Branches—

Confidential	Conf.
Political	P.
Vernacular	V.
Accounts	A.
Chief Commissioner's	C. C.

APPENDIX III

DIRECTIONS REGARDING INDEXING AND OPENING FILES

INDEXING [File paragraph 49]

The object of indexing is to ensure that papers on any subject are readily available, and that so far as possible all the papers on one subject are indexed in one place.

The index has been drawn up with this object. It will be observed that some heads are placed in square brackets with a cross reference, e.g.,

[Travelling Allowance see Allowances]

No papers may be indexed under such heads, the object of this prohibition is to limit the discretion of the indexer to index some papers under Allowance and some under Traveling.

2 A file will be indexed under a major head, a minor head if one is given in the index, and a sub head or individual head. New major and minor heads may only be opened with the permission of the Secretary and when this is done all holders of the index must be notified. The choice of sub heads is left to the indexer and new sub-heads must be opened whenever necessary. They should, however, be as brief as possible and a new sub head should not be given when a suitable one already exists. It is possible for more than one file to be indexed under the same sub head.

3 Major heads are as general as possible and when there is an alternative, the most general head should be used in indexing. It will be observed that the following major heads appear in the index —

Ruling Princes, the names of individual States

Adoption, Jagirdars, Arms and Ammunition, Treaties.

In connection with these heads the following principles should be observed —

(a) Under Ruling Princes only matters of common interest affecting Ruling Princes generally should be indexed under the various minor heads provided.

(b) Under the names of individual States should be indexed such matters as relate specifically to the State concerned and have no direct connection with other States or States as a whole for instance —

Individual

in the

only,

individual States or under Istimrardars, etc. But where an important general principle emerges in a particular case, it should be indicated under the head Adoption etc.

body

h action

sub head

adopt

useful

(c) Where, however, a matter is of more common interest to all States and forms in itself a well defined subject, it should be indexed under the more general head, e.g., Arms and Ammunition—Armed Police Treaties—Revision Jagirdars—Disputes. A cross reference may in such cases be given under the names of the States if thought desirable.

(d) Some cases may appear to fall equally under one or more of such major heads,

e.g.,

mix

hant

so f

be

be

4. The principle that major heads should be as general as possible is subject to the limitation that if a head is too general it is useless. Thus major heads 'Ajmer Merwara', 'Rajputana' are too general to be useful. It is for this reason that an apparent inconsistency appears under the head Budgets, for while Ajmer Merwara and Rajputana Agency Budgets appear under the head Budgets, other Budgets appear under the name of the body or department, e.g. Mina Corps, Forests. Municipal Budgets should similarly be indexed under the major head Municipalities and the minor head of the Municipality concerned, e.g. Municipalities—Ajmer—Budget—1923-24.

5 Under certain important Major heads a minor head General appears, e.g., Military, Education, Judicial. The object of this minor head is to cover all matters not covered by the remaining minor heads. It is possible that in some cases experience will show that further sub division is necessary.

6 A major head for the title of any Act may be opened without special permission.

7 Separate indices will be maintained for Chief Commissioner's Branch and for the rest of the Agency Office. Under each of these divisions, again, there will be a name index as well as a general index. The name index will often merely refer to files indexed in the

Major X—Motor advance, application for .File-45 A/23

Posted as Political Agent, Southern Rajputana States File 30 P /23

Applications for leave, however, will not be indexed in the general index, but only in the name index.

8 It is not necessary to index cases for which special registers are maintained under the provisions of Rule 26 (a), (c), (d), (e), (f), (g) of the Manual.

OPENING FILES

1 With a card index system properly maintained there is little fear of relevant papers being overlooked. There is therefore no object in having a large number of papers only loosely related to one another on one file and new files should be freely opened.

2 As far as possible a file should form a consecutive narrative and the narrative should not be interrupted by the intrusion of extraneous matter

[illegible]

4 Separate files should be maintained for individual officers, individual States, etc. For instance, deputation of B to State A as Superintendent of Police should be dealt with on a separate file from that dealing with the deputation of A, his predecessor. The connecting link will be the index. Both will appear under Foreign Service—X State, and the deputation will also be referred to under the names of the officers in the name index.

5 Subject to the provisions of the Manual, old files can, where suitable, be re opened, e.g., D, Istimrardar of W estate, dies in 1917 and the succession of E is sanctioned. In 1924, E dies and Commissioner recommends the succession of F. The 1917 file can conveniently be re opened.

6 Great care must, however, be taken in all cases to prevent files becoming bulky and unwieldy. If in a case similar to that mentioned in Rules 3 or 5 there is any likelihood of the correspondence being protracted, a separate file should be opened at once.

7 Similarly, when any correspondence contains a large number of references and exhibits, it is desirable that exhibits should be kept separate in a file bearing the same number.

but marked Part II. It may also be desirable to keep routine papers such as reminders and other unimportant communications, which would ordinarily be destroyed before the file is closed, in a separate temporary file.

8 On important subjects a separate file should be maintained for general orders, *et cetera*. Care must be taken not to lose sight of the principle that the file should be necessary to maintain a

file of general orders, it will be sufficient to indicate such principles by cross references in the index. Suppose, for instance, a file is indexed —

Posting Political Officers—Colonel Spence as Resident, Mewar

In this connection, a question arises regarding the allowances which may be granted for holding a dual charge. This can be indicated by the following cross reference —

Allowances—Charge—Dual charge of Mewar and S R S *vide* Postings, etc

9 In some cases it may be desirable to maintain several temporary files which can be amalgamated into one before closing, *et cetera*, in the case of tours or ceremonial visits. It is desirable that the papers regarding each visit should be kept separate until the visit is over. At the end of the season routine papers should be destroyed and the files weeded. Important papers can then either be combined into one file, *et cetera*, tour arrangements, winter 1923-24 or placed on existing files regarding each place visited, if such are maintained, *et cetera*, Ceremonial—Visits—Jaipur

10 Before a file is closed, it should be carefully weeded, unimportant papers such as reminders, interim replies, duplicate copies, etc., being destroyed. A note should be made in the notes against each paper that has been destroyed

INDEX FOR RAJPUTANA AGENCY OFFICE

A

*Major Head**Minor Head*

[Abdication see Ruling Princes and individual States.]

ABU (see also Courts Municipality, Education and Postings)

General

Lease

Leased Area

[Abu High School see Education.]

[Accession see Ruling Princes and under individual States]

ACCIDENTS (see also Railways.)

ACCOUNTS - - - -

Adjustment

Objections.

Refund.

ACCOMMODATION

[Acquisition of land (see also Europeans Ruling Princes and Land Acquisition Act)]

[Acts see under Title of Act]

[Address see Correspondence Government Servants and Speeches]

[Adjustment see Accounts.]

ADMINISTERED AREAS

Abu

Deoli

Erinpura

General

Kierwara

Kotra

Railways

ADMINISTRATION REPORTS - - -

Ajmer Merwara

States

ADOPTION (see Ruling Princes Istimardars and under individual States)

General

ADVANCES .

House-building

Motor Car

Passage

Provident Fund

Permanent

Transfer

ADVERTISEMENTS

[Advocate see Pleaders]

ADVOCATE GENERAL

ADVISORY COUNCIL

MAGGP

*Major Head.**Minor Head*

[Aeroplanes see Aviation]

[Affiliation see Education]

AFFRAYS

AFGHANS

AGENCIES, POLITICAL

[Agents, Political see Postings]

AGENT TO THE GOVERNOR GENERAL (see also Postings)

[Agreements see Extradition Treaties]

AGRICULTURE

[Aircraft see Aviation]

AITCHISON'S TREATIES

AJMER (to be used only for matters concerning Ajmer
not falling under any other head)

ALIENATION

[Aliens see Foreigners]

ALEXANDRIA

ALLOWANCES

Abu.

Bonus.

Charge

Compassionate.

Compensation

Conveyance

Deputation

Duty.

Exchange Compensation.

Grain compensation

Halting

Honorarium.

Horse.

Local.

Motor Car

Officiating

Overseas.

Personal.

Shorthand.

Sumptuary.

Travelling—Daily and Mileage.

ALWAR.

Ammunition see Arms and Ammunition]

AMBULANCE, St Johns

Andamans see Convicts]

*Major Head.**Minor Head*

ANIMALS, CRUELTY TO

ANIMALS, WILD

[Annual Reports see Returns]

[Anti Malaria see Medical]

ANTI RABIES TREATMENT

[Appeals (see Judicial, Memorial, etc)]

[Appeals for funds see Funds]

APPRENTICE (see also Railways).

APPOINTMENTS General orders (see also Estab-
lishment and Postings)

[Arbitration see Boundary Disputes and Judicial]

ARBORICULTURE

ARCHAEOLOGY

ARCHITECTURE.

ARMS ACT Ajmer Merwara.
StatesARMS AND AMMUNITION . . . Armed Police
General orders.
Licenses
Saluting Guns
State Forces
Smuggling
Troops and other than State
Forces

[Armed Police see Arms and Ammunition and Police]

ARMISTICE

ARMORIAL BEARINGS

[Army see Military]

ARREARS

[Arrest see Extradition and Judicial.]

ART

ARTIZANS

ARYA SAMAJ

[Assault see Affrays]

[Assembly see Legislative Assembly]

[Assessment see Income tax, Land Revenue, Mun-
icipalities]

ASSIGNMENT

[Assistant Surgeons see Medical]

ASSISTANT COMMISSIONER (see also Postings)

ASYLUM (see also Lunatics)

ATTACHE.

ATTACHMENT

BONDS

[Bonus see Allowances]

BOOKS AND PUBLICATIONS

BORDER COURTS

BORING

BOTANICAL

BOUNDARY	Disputes.
					General orders.
					Settlement Officer

BOY SCOUTS

BREEDING	Cattle
				Horse

[Bridges see Civil Works]

BRITISH SUBJECTS (see also Births, Deaths and Marriages, Extradition)

BUDGETS (see also Education, Municipalities, etc)	..	Ajmer Merwara.
		General
		Rajputana Agency.

[Buildings see Civil Works]

[Building sites see Municipalities, Delhi, Abu Leased Area]

[Bungalows see Residences, Rest Houses]

BUNDL

[Bunds see Civil Works]

BURIAL PLACES (see also Ecclesiastical)

C

[Cadets see Military]

[Camp see Tours]

CAMP EQUIPMENT

CANALS

CANTONMENTS	(Deoh see Administered Area.)
					(Erinpura do)
					(Kherwara do)
					(Kotra do)

CANTONMENT ACT	[Deputy Magistrate see Courts, Judicial]
					Executive Officer.
					General.

CARS (see also Advances)

CASH CERTIFICATE

CASUALTIES

[Casual leave see Leave]

CATALOGUES

[Cattle see Breeding, Grazing]

CATTLE SLAUGHTER

[Cavalry see Military]

[Cemetries see Ecclesiastical]

CENSOR

CENSUS	Ajmer Merwara Rajputana
--------	----	----	----	----	----	----------------------------

CEREMONIAL (see also Tours)	General Visits
-----------------------------	----	----	---	-------------------

CERTIFICATES (see also Medical).

[Cess see Land Revenue]

[Chakri see Jagirdars]

CHAMBER OF PRINCES

[Chaprasis see Establishment]

[Chaplains see Ecclesiastical]

[Charas see Hemp Drugs]

[Charge Allowance see Allowances]

CHARITY.

CHEMICALS

CHEMICAL EXAMINATION

[Chiefs see Ruling Princes]

[Chiefs Colleges see Education and Mayo College]

CHIEFS AND LEADING FAMILIES IN RAJPUTANA

CHILD WELFARE

[Cholera see Medical]

[Christians see Births, Deaths and Marriages]

[Church see Ecclesiastical]

CINEMA.

[Circulars see under subjects dealt with]

CIVIL ACCOUNT CODE.

CIVIL PROCEDURE CODE.

CIVIL SUITS	General
-------------	----	----	----	----	----	---------

[Civil Service Regulations see Fundamental Rules]

*Major Head**Minor Head.*

COPIES					
COPYRIGHT					
CORONATION					
[Corps see Bhil Corps and Mina Corps]					
CORRESPONDENCE	[Direct see Returns]	
				Methoda.	
				Police	
				Sepoys	
COTTON					
COUNCIL OF STATE					
[Council see Advisory Council]					
[Council of Regency see Ruling Princes and under individual States.]					
COURTS (Constitution and Powers) (see also Judicial)				Abu	
				Bench	
				Civil	
				Criminal	
				District	
				High	
				Magisterial.	
				Sessions	
				Small Cause	
COURT FEES ACT					
COURT OF WAKILS			.	Appeals	
				Appointment, Leave, etc., of	
				Wakils	
				Convicts	
				General	
				Jurisdiction	
				Rules	
COURT OF WARDS	(For individual Estate see under individual Istumrardars)
[Criminal Intelligence Department see Police]					
CRIMINAL PROCEDURE CODE					
CRIMINAL SUITS	General.	
CRIMINAL SETTLEMENT					
CRIMINAL TRIBES					
[Crops see Returns]					
CRUELTY TO ANIMALS.					
CURRENCY	Coinage
					Mints
					Notes.
CUSTOMS (see also Jagirdars)					

D

DACCOTIES

[Dais see Medical.]

DAK

DALY COLLEGE.

[Dami see Istimsardars.]

[Darbars see under individual States.]

DARBAHS - - .. Agent, Governor General's.

DARGAH.

[Deaths see Ruling Princes, Births, Deaths and
Marriages, Registration Act, Europeans.]

DECREEES.

DECORATIONS

DEESA

[Defence Force see Military.]

DEFALCATIONS

DEFAMATION

DEFENCE (see also Military.)

[Degrees see Education.]

DELEGATION (see also Finance.)

DELHI SITES

DEMARCATION (see Boundary.)

[Demobilisation see Military.]

[Deoli see Administered Areas.]

DEPORTATION

DEPOSITS

DEPUTATION (see Foreign Service.)

[Deputation Allowance see Allowances.]

[Deserters see Sepoys.]

[Destruction of Records see Records.]

[Destruction of Wild Animals see Animals.]

DHOLPUR.

DIARIES.

[Diet Charges see Judicial.]

[Diploma see Mayo College.]

[Direct relation see States.]

[Discipline see Office.]

[Diseases see Medical.]

DISMISSAL - - - - . General orders
(Individual cases will be
decreed under the name of
person concerned.)

MAGGR

*Major Head**Minor Head*

[Dispensary see Medical]

DISPOSAL OF ENEMY PROPERTY

[Disputes see Grazing Boundary Jagirdars etc]

[District Boards see Boards]

District Magistrate see Courts, Judicial and Post
ings]

DISTRICT SOLDIERS BOARD.

DISTURBANCES AND RIOTS

DOCUMENTS

DOMINIONS

[Donations see Charity Funds]

DRAINAGE (see also Municipalities)

DRAMATIC PERFORMANCES

DRESS REGULATIONS

DRUGS (see also Hemp Drugs and Excise)

e

Cocaine

Lucences

Morphia

Smuggling.

DUFFELIN FUND

DUNGARPUR

DUPLICATORS

[Duty Allowance see Allowances]

F

EARTHQUAKES

EASEMENTS

ECCLESIASTICAL

Bishop

Budget

Cemeteries

Chaplains

ECONOMIC CONDITIONS

EDUCATION (Ajmer Merwara and Rajputana)

Abu

Abu High School

Affiliation

Budget

College's Colleges

Degrees

Educational Commissioner

General.

Grants in aid

Indian Educational Service

Inspection

Lawrence School

[Mayo College see Mayo College.]

Post Graduate

Primary

Scholarship

Secondary

Technical

University

*Major Head,**Minor Head,*

EDUCATION DEPARTMENT

[Elections see Legislative Assembly, Municipality and Cantonments]

ELECTRIC

ELEPHANTS

EMIGRATION

EMPIRE

ENCROACHMENT (see also Municipality)

ENDOWMENTS

[Enemy Missions see Missions]

[Enemy property see Disposal]

ENEMY TRADING

ENGINEERING (see also Education)

ENGRAVINGS

ENTERTAINMENT

[Epidemic Diseases see Medical]

[Equipage see Camp]

[Erinpura see Administered Area]

ESCORT (see also Ceremonial)

ESTABLISHMENT, MINISTERIAL ..

Ajmer Merwara
Rajputana Agency

ESTABLISHMENT, MENIAL

Ajmer Merwara
Rajputana Agency

[Establishment, Revenue see Revenue]

[Establishment, Judicial see Judicial]

[Estates see Jagirdars Istimrardars]

[Estimates see Budgets, P W D]

ETIQUETTE

EUROPEANS	Acquisition of Land and property by Europeans Deaths of Europeans [Employment in States and Districts] [Qualification of States and Districts]
-----------	---	---	----	----	---

EUROPEAN VAGRANTS

EVIDENCE (see also Extradition)

EXAMINATIONS.

FINANCE.

[Exchange Compensation see Allowances]

FIXED (see also Hemp Drugs)

Ajmer Merwara
States

EXECUTIVE FACILITIES

*Major Head**Minor Head***EXHIBITIONS.**

[Expenditure see Finances]

EXPLORATION.**EXPORT**

EXPLOSIVES (see also Arms and Arms Act) . . . Licences.

EXTERNAL RELATIONS

EXTRA ASSISTANT COMMISSIONER (see also Revenue)

EXTRADITION	Act
	Agreement.
	Arrest
	General
	Transit Charges.
	Treaty

F**FACTORIES.****FAIRS**

[Family Pensions Fund see Pensions]

FAMINE

[Feasts see Festivals]

FERRIES**FESTIVALS**

FEUDATORIES (see also Jagirdars)

[Films see Cinema]

FINANCE	Book of Financial powers
	General (only general questions
	of finance to be indexed
	under this head, e g , Ajmer
	Merwara Finance, other
	questions, e g , reappropria-
	tion will be dealt with under
	departmental heads)
	Powers of Local Administration
	Standing Finance Committee

FINANCE DEPARTMENT

[Fines see Judicial, Government Servants]

FIRE

[Finger Impressions see Police]

FLAGS**FLOODS****FODDER.****FOOD-STUFFS.****FORECAST****FOREIGN AND POLITICAL DEPARTMENT****FOREIGNERS**

FOREIGN SERVICE	General charges (Furniture and books as per the estimate of the Government of India)
FORESTS (Ajmer Merwara)	Forest Forest (Forest) Indian Forest Service
FORESTS (see also Jagirdars)	
[Forms see Stationery.]	
FORTS.	
[Fortnightly Reports see Reports]	
FRAUD.	
FRONTIER	
FUNDS (see also Provident Fund)	Appropriation (Furniture and (Furniture) (Miscellaneous for each fund)
FUNDAMENTAL RULES.	
[Furlough see Leave.]	
FURNITURE (see also Easements)	
()	
GAMBLING :	
GAME	
GAMES.	
[Ganja see Hemp Drugs.]	
GARDENS (see also Plantations)	
GAZETTE OF INDIA ,	
GAZETTES.	
GAZETTEERS,	
[Geological and Survey]	
GEOLGY.	
GERMANS.	
GRATS.	
GIFTS (See also Government Grants)	
GIRASSIAS.	
GIRL GUIDES.	
[Governments see Companions]	
GOVERNMENT OF INDIA ACT.	

*Major Head**Minor Head*

EXHIBITIONS.

[Expenditure see Finances]

EXPLORATION

EXPORT

EXPLOSIVES (see also Arms and Arms Act) . . . Licences.

EXTERNAL RELATIONS

EXTRA ASSISTANT COMMISSIONER (see also Revenue)

EXTRADITION	Act
				Agreement
				Arrest
				General
				Transit Charges.
				Treaty

F

FACTORIES.

FAIRS

[Family Pensions Fund see Pensions]

FAMINE

[Feasts see Festivals]

FERRIES

FESTIVALS

FEUDATORIES (see also Jagirdars)

[Films see Cinema]

FINANCE	Book of Financial powers
				General (only general questions
				of finance to be indexed
				under this head, e.g., Ajmer
				Merwara Finance, other
				questions e.g., reappropria-
				tion will be dealt with under
				departmental heads)

Powers of Local Administration
Standing Finance Committee

FINANCE DEPARTMENT

[Fines see Judicial, Government Servants]

FIRE

[Finger Impressions see Police]

FLAGS

FLOODS

FODDER.

FOOD-STUFFS

FORECAST

FOREIGN AND POLITICAL DEPARTMENT

FOREIGNERS

[Horse allowance see Allowances]

[Horse Breeding see Breeding]

HOSE SHOW

[Hospitals see Medical]

[Hot weather charges see Budgets]

[House sites see Municipal, Delhi, Abu Leased Area.]

[House tax see Municipal.]

I

. IDENTITY CERTIFICATES

IMMIGRATION

IMPERIAL BANK

IMPERIAL CONFERENCE

[Imperial Indian Relief Fund see Funds]

IMPERIAL INSTITUTE

IMPERIAL SERVICE TROOPS (see also State Forces.)

IMPORTS.

IMPOSTORS.

[Improvement see Land Improvement Loans Act]

INCOME TAX

Assessment

Refund

Super tax

INDEMNITY BONDS (see also Stamp Act)

INDENTS (see also Stationery)

[Index see Office Procedure]

INDIAN CIVIL SERVICE

[Indian Army see Military]

[Indian Army Reserve Officers see Military]

[Indian Defence Force see Military]

[Indian Educational Service see Education.]

[Indian Forest Service see Forests.]

[Indian Medical Service see Medical.]

INDIAN PENAL CODE.

INDIAN SOLDIERS BOARD

[Indian State Forces see State Forces]

INDIAN STATES

INDUSTRIES.

INFANTICIDE.

[Informal Conference see Conferences.]

[Influenza see Medical.]

[Inoculation see Medical.]

[Insanity see Lunatics.]

INSCRIPTIONS.

INSECTS.

INSOLVENCY.

[Inspector General of Police see Police.]

[Installation see under individual States and Ruling Princes]

INSTITUTES.

INSURANCE.

INTELLIGENCE BUREAU.

[Internal Security see Military.]

[Interrogatories see Judicial]

INTERVIEWS.

INTRODUCTORY LETTERS.

INVENTIONS.

[Investiture see Honours]

INVITATIONS.

[Irrigation see Civil Works]

ISTIMRARDARS (also includes Ajmer-Merwara Jagir-dars)	Adoption.
	Dam
	Disputes.
	Estates.
	Loans.
	Succession.
	Tazim.

J

JAGIRDARS (see also Istimrardars)	Chakri.
	Customs.
	Disputes.
	Estates
	General.
	Succession,
	Tazim.

JAILS.

JAINS.

JAIPUR.

JAISALMER.

[Jangi Inams see Military.]

JHALAWAR.

JODHPUR.

JOURNALS.

JUBILEE.

[Judge see Judicial.]

[Judgment see Judicial]

Major Head.

Minor Head.

JUDICIAL	Appeals
						Appointments (Ajmer Mer- wara)
						Appointments (Administered Areas)
						Arrests
						Bench
						Commissions
						[Courts see Courts]
						Diet Charges
						Establishment (Ajmer Mer- wara)
						Establishment (Administered Areas)
						Fines
						General
						Interrogatories
						Judges
						Judgments
						Judicial Commissioner
						Justice of the Peace
						Magistrates
						Powers
						Privy Council
						Procedure
						Summonses
						Warrants
JURISDICTION (see also Administered Areas, Court of Valids, Extradition)	Disputes General
[Justice of the Peace see Judicial]						
K						
KABAULI						
KHARITAS						
[Kherwara see Administered Areas]						
KHETRI						
KHILLATS						
KING, HIS MAJESTY THE						
[King Edward Medical School see Medical]						
KISHENGARH						
KOTAH						
KUSHALGARH						
L						
LABORATORIES						
LABOUR						
[Ladies see Companions]						
LAND	Disputes General (see also Municipalities)
LAND ACQUISITION ACT	Injection
LAND IMPROVEMENT LOANS ACT						
[Landed Property see Europeans and Ruling Princes]						
LAND REVENUE (Ajmer-Merwara)	Assessment, General Excess Exemption [Establishment see Princes Revenue] [Exemption see Exemption] Settlement (see also Rents)

Major Head

Minor Head

INQUAGL (see also Examinations.)

LAW REPORTS

LAWA

[Lawrence School see Education]

LEAGUE OF NATIONS

LEASE (see also Abu and Mining)

LEAVE ..

..

General orders

(For individual cases see under names of officers concerned)

LECTURES.

LEGAL PRACTITIONERS

LEGISLATIVE ASSEMBLY

..

Elections

General

Questions

Voices

LEGISLATION (see also Administered Areas Abu, Municipality)

LEGISLATIVE DEPARTMENT

LEISERS

LEISERS

LIBRARY

[Licences see Arms Drugs Excise Explosives etc]

IMITATION ACT

[Liquor see Excise]

LITERATURE

[Liversies see Establishment]

LOANS (see also Advances, Istimrarjars and under individual States) (Minor heads for each kind of loan)

[Local Allowances see Allowances]

LOCAL FUND

LOCAL GOVERNMENT

LOCALISTS

LOTTERIES

LUNATICS

[Lymph see Medical]

M

MACHINERY

MAGAZINES

[Magistrates see Courts, Judicial]

Major Heal

[Malaria see Medical]

[Manœuvres see Military]

[Manual see Office]

MARS

MARKETS

[Marriages see Ruling Princes, Births Deaths and Marriages]

[Marwar see Jodhpur]

MATERNITY (see Child Welfare)

MAYO COLLEGE

MEDALS

MEDICAL

Minister

Admissions

Budget

Constitution

Diploma

Education

Establishment

Finances

General

Lost Diploma

Imports

Anti Malaria

Appointments in Rajputana (see also Foreign Service)

Assistant Surgeons

Boards

Chief Medical Officer

Civil Surgeons

Dais

Dispensaries

Epidemic diseases

Establishment

General

Hospitals

Indian Medical Service

Inoculation

King Edward Medical School

Lymph

Midwives

Minto Nursing Association

Post Graduate Studies

Quarantine

Sub Assistant Surgeons

Supplies

Training

Vaccination

MEMORANDA ON INDIAN STATES.

MEMORIALS

.. .. . General orders (for individual cases are under subject of cases)

[Memorial Funds see Funds]

MERCHANTS.

*Major Head**Minor Head*

MERCINARIES

METEOROLOGY

MEWAR (see Udaipur)

[Midwives see Medical]

[Mileage allowance see Allowances]

MILITARY [see also Arms and Ammunition, Sepoys,
State Forces]

Accounts

Appointments

Auxiliary Forces

Cavalry

Defence Force

Demobilization

General

Indian Army

Indian Army, Reserve Officers

Internal Security

Jangi Inams

Labour Corps

Manœuvres

Mobilization

Mutiny

Ordnance

[Pensions see Pensions]

Prince of Wales Military College

Reconnaissance

Recruiting

Sandhurst

Territorial Forces

Volunteer

MILLS

MINA CORPS

Budget

Finances

General

Officers

Pensions

MINAS

MINERALS

MINES

Lease

Royalty

[Minority Administration see Ruling Princes and under
individual States]

MIRORS

[Minto Nursing Association see Medical]

[Mints see Currency]

[Mir Munshi see Establishment Ministerial]

MISCONDUCT (See also Government Servants)

MISSIONS

[Mobilization see Military]

*Major Head**Minor Head*

MOGHULS.

MONOPOLIES

[Monsoon see Meteorology]

MONUMENTS (see also Archaeology)

[Morphia see Drugs]

MOSQUES

MOTOR CARS (see Cars)

[Motor Cars Advances see Advances]

[Motor Cars Allowance see Allowances]

MUHAMMEDAN LAW

MUHAMMEDANS

MULTIGRAPH

MUNICIPALITIES

Abul

Ajmer

Beawar

General orders.

Kekri

[Munitions see Arms and Ammunition]

MUSEUM

[Mutines see Military, State Forces.]

N

[Narb Tahsildars see Revenue]

NATIONAL ANTHEM

NATIONALITY

NATURALIZATION (see also British Status and Naturalization of Aliens Act)

NAYARS

NEGOTIABLE INSTRUMENT ACT

NEWSPAPERS

NOBLES (see also Jagirdars) Disputes
General.

NON CO OPERATION

[Notes Currency see Currency, Treasury]

NOTES, PROMISSORY.

NOTICES

NOTIFICATIONS.

[Nursing see Minto Nursing Association.]

O

OATHS.

[Objection Statements see Accounts.]

OBSERVATORY.

*Major Head**Minor Head*

OCTROI (see also Municipality)

OFFICE

Discipline

[Establishment see Establishment]

General

Manual

Orders

Procedure

[Officers see Government Servants Political Officers]

OFFICIAL RECEIVER

OFFICIAL TRUSTEE

OPIUM (see also Drugs)

ORDINANCES

[Ordnance see Arms and Ammunition and Military]

[Oriental Languages see Examinations]

OUTLAWS

[Out posts see Police]

OUTRAGES

[Overseas Allowance see Allowances]

P

[Panchayat Courts see Court of Vakils]

PANCHAYATS

[Paper Currency see Currency]

PARLIAMENT

PARTITION (see Istimrardars Jagirdars)

PARWANAS

[Passage Advance see Advances]

PASSAGES

[Passes see Arms Act]

PASSPORTS (General orders only Individual applications registered in Passport Register)

PAUPERS

[Pay see Salary]

PEACE

PENAL SETTLEMENT

PENSIONS

Civil

Commutation

Compassionate

Gratuity

Military

Proportionate

Retiring

Superannuation

Wound

[Periodicals see Returns]

[Permanent Advance see Advances]

PERSONAL ASSISTANT,

LISTS

*Major Head**Minor Head.*

PETITIONS (see also Memorials: general orders only
Individual cases under subject head)

PETROL.

PETROLEUM

PHOTOGRAPHY

POSTSTICKING

PILGRIMS

[Plague see Medical]

PLEADERS

POISONS

POLICE (Ajmer Merwara and Railway)

Appeals
Armed Police
Budget
Criminal Intelligence
Finance
Finger Impression.
General
Inspector General of Police
Outpost
Superintendent

[Political Agency see Agency]

POLITICAL AGENTS

POLITICAL AGITATION

[Political Department see Foreign and Political Department]

POLITICAL MEETINGS

POLITICAL OFFICERS (General Individuals under names)

POLITICAL OFFICERS MANUAL

POLITICAL PRISONERS

POLITICAL SUSPECTS

POLO

POSTS AND TELEGRAPHS

Ajmer Merwara
States

Post Diploma see Mayo College]

Post Graduates see Education and Medical]

Post office Cash Certificates see Cash Certificates]

POST OFFICE (see also Posts and Telegraphs)

POSTINGS Political Officers

POWER OF ATTORNEY

PRATAPJAHN

[Pratap Memorial Fund see Funds]

PRECEDENCE

[Presents see Gifts]

PRESS

*Major Head**Minor Head.*

PRICES

PRIESTS

PRINTING

PRISONERS

[Privilege leave see Leave]

[Privileges see Easements and Ruling Princes]

[Privy Council see Judicial]

PRIZE MONEY

PRIZES

PROBATE

[Proceeding see under subject head]

[Process see Judicial]

PROCESSIONS

PROCLAMATIONS

[Programmes see Tours]

[Promissory see Notes, Promissory]

[Promotions see Establishment Judicial, Revenue]

PROPAGANDA

[Property see Europeans and Ruling Princes]

PROPERTY, TRANSFER OF—

PROPERTY, UNCLAIMED.

[Proportionate Pension see Pensions]

PROPOSITION STATEMENTS (General orders only)

[Prospecting see Mines]

PROSTITUTES.

[Provident Fund see Funds, Pensions]

PUBLIC PROSECUTOR

PUBLIC SERVANTS (see also Government Servants)

PUBLIC SERVICES

PUBLIC WORKS DEPARTMENT

Budget

[Civil Works see Civil Works.

Establishment

Estimates

Finance

General

[Publications see Books and Publications]

PUBLICITY.

Q

[Quarantine see Medical]

QUARRIES.

*Major Head.**Minor Head.*

- [Quarterly returns see Returns]
 [Quarters see Civil Works, Establishments]
 [Questions see Legislative Assembly]
 [Quinine see Medical]

R

- [Rabies see Anti Rabies]

RAILWAYS (see also Administered Areas) . . .

- .. Accounts
 Apprentice.
 Bombay, Baroda and Central
 India Railway.
 General.
 Refreshment Rooms.
 States
 Workshops.

- [Railway Magistrate see Courts and Judicial]

- [Railway Police see Police]

- [Rainfall see Returns]

RAINGAUGE

- [Re appropriation Statement see Budgets, Education,
 etc]

RECEPTIONS.

RECESS

- [Reconnaissance see Military]

- [Record of Rights see Land Revenue]

RECORDS (see also Land Revenue) Destruction.
 General.
 System
 Weeding.

- [Recruiting see Military]

- [Red Cross see Ambulance, St John.]

REFORMATORIES

REFORMS

- [Refreshment Rooms see Railways]

- [Refunds see Accounts, Income tax, etc]

- [Regency see Ruling Princes and under individual States]

- [Regiment see Military]

REGISTERS.

REGISTRAR.

REGISTRATION ACT.

- [Registration see Births, Deaths and Marriages]

REGISTRATION DEPARTMENT.

- [Regulations see under subject head.]

- [Relief, Imperial Indian Fund see Funds.]

- [Religious Institutions see Ecclesiastical, Mission.]

- [Religious Endowments see Endowments.]

RAGGR

*Major Head**Minor Head*

[Remissions see Land Revenue]

[Remittance see Treasury.]

REMOUNTS.

REPATRIATION.

[Reports see Administration Reports and Returns under subject-head]

RESIDENCIES.

[Residencies see Civil Works]

RESIDENTS (see also Agents, Political)

REST (see also Residencies, Civil Works)

[Rest Houses see Civil Works]

RETIREMENT (General orders).

RETRENCHMENT (General see also under separate Departments).

RETURNS

REVENUE (see also Land Revenue)

Appeals.

Appointment.

Establishment (Ajmer Merwara)

General.

REWARDS

[Riots see Disturbances]

[Roads see Civil Works]

ROBBERY.

ROUTE BOOKS

ROYAL FAMILY (see also King)

[Royalty see Mines, Royal Family]

RULING PRINCES

Abdication.

Accession

Acquisition of landed property.

Adoption.

Council of Regency.

Family events.

General.

Installation.

Minorities.

Precedence.

Regency.

Rights & Privileges.

Salutes.

Visits.

SARFA.

SARFARAZ.

*Major Head**Minor Head*

SALARY (see also Establishment, etc.)

SALT (see also Treaties).

SALOON.

SALUTES (see also Ruling Princes).

[Saluting Guns see Arms & Ammunition.]

SAMBHAR.

[Sanads see Honours]

SANITATION (see also Municipality).

[Sardars see Nobles.]

SAROPAO.

SATTAGRAHA.

SAVINGS BANK.

[Scarcity see Famine]

[Scholarship see Education.]

[School see Education.]

[Scouts see Boy Scouts]

[Sea Customs see Customs]

[Season Report see Returns]

SECRETARIAT.

SECRETARY TO A. G. G.

[Secretary, Indian Soldiers Board see Indian Soldiers Board]

SECRETARY, PUBLIC WORKS DEPARTMENT.

SECRETS, OFFICIAL

SECURITIES (see also Loans).

SEDITION.

SELF-GOVERNMENT.

SEPOYS	--	--	--	--	--	Arrest (see also Extradition).
						Deserters
						[Pensions see Pensions]
						[Petitions see Petitions.]
						Verification Rolls.

SERVICE BOOKS

[Sessions see Courts and Judicial.]

SETTLEMENT (see Land Revenue and Boundary).

SHANTRA.

SHOOTING.

SHORTHAND.

SIKAR.

[Silver Wedding Fund see Funds.]

*Major Head**Minor Head.*

TONE.

TOPOGRAPHY.

TOSHAKHANA.

TOUR	Arrangements. Charges. General. Programmes.
------	----	----	----	----	----	--

[Tracking Rules see Court of Vakils.]

[Trade see Industries.]

TRAFFIC.

TRAMWAYS (see also Railways, States).

[Transfer of property see Property.]

TRANSFER OF PROPERTY ACT.

TRANSFERS	..	-	-	..	General order. (See also Postings, Establishment, etc.).
-----------	----	---	---	----	---

[Transit charges see Extradition, Prisoners.]

TRANSLATION.

TRANSPORT.

TRANSPORTATION.

TRAVELLERS (see also Court of Vakils).

[Travelling Allowances see Allowances.]

TREASURE TROVE.

TREASURER.

TREASURY.

TREATIES	-	!	..	Execution. General General orders. Revision.
----------	---	---	----	---

[Trees see Arboriculture]

[Tribes see Criminal Tribes.]

TRIBUNALS.

TRIBUTE (see also under individual States)	General.
--	----	----	----------

[Trigonometrical Survey see Survey]

[Troops see Military, State Forces]

TROPHIES.

TRUSTEES.

TRUSTS.

TYPEWRITERS.

U

[Udaipur see Mewar]

[Unclaimed property see Property.]

UNDER SECRETARY (see also Postings, Political Officers).

*Major Head.**Minor Head.*

UNDESIRABLES.

{Uniform see Dress Regulations.}

{University see Education.}

V

{Vaccination see Medical.}

VAGRANTS.

VAKALAT.

{Vakils see Pleaders.}

{Verdacular see Language.}

VETERANS

VETERINARY.

VICE-REGAL VISITS (General orders for particular visits see under individual States, etc.)

VICEROY.

VILAYATIES.

{Visas see Passport.}

VISITORS.

VISITS (see also Tours.)

VITAL STATISTICS.

{Volunteers see Military.}

VOTES (see also Legislative Assembly, Municipal and Cantonments).

W

WAGES.

WALTERKRIT RAJPUTHA HITKARINI SABHA.

WANDERING GANGS (see also Criminal Tribes).

WAQF.

WAR.

{War Legislation (see Legislation)}

{War Loans see Loans.}

{Wards see Court of Wards.}

{Warm clothing see Establishment.}

{Warrant see Judicial.}

{Warrant of Precedence see Precedence.}

{Water Supply see Municipal.}

WEATHER REPORTS.

WEAVING.

WEDDINGS.

WEIGHTS AND MEASURES.

Major Head.

Minor Head.

WELLS.

[Weeding see Records.]

WHIPPING.

[Wild Animals see Animals.]

WILLS.

WITCHCRAFT.

[Workshops see P. W. D. and Railways.]

[Wound Pensions see Pensions.]

X

Y

Z

ZOOLOGY.

APPENDIX IV

No I

[See Government of India Standard Form S 8]

RAJPUTANA AGENCY

No. V OF 192 .

Dated Abu, the 192 .

Forwarded to the
for disposal

By order,

RAJPUTANA AGENCY OFFICE.

Dated Abu, the 192 . }

Secretary to the Hon'ble the Agent
to the Governor-General,
Rajputana.Petition, dated the 192 .
from

No II

[See Government of India Standard Form S. 8]

RAJPUTANA AGENCY.

No. V OF 192 .

Forwarded to the
for disposal2 The petitioner has been informed of the action taken by this office.
petitioners have

By order,

RAJPUTANA AGENCY OFFICE.

Dated Abu, the 192 . }

Secretary to the Hon'ble the Agent
to the Governor-General,
Rajputana.Petition dated the 192 .
from

MAGG R

No III

[See Government of India Standard Form S 8]

RAJPUTANA AGENCY

No V, dated Abu, the 192

Forwarded to the

for favour of inquiry and a brief report

By order,

Secretary to the Hon'ble the Agent
to the Governor General
Rajputana.

Vernacular petition, dated the 192, from
English

No IV

[See Government of India Standard Form S 8]

No V. of 192

Forwarded to

with the request that, if he sees no objection, the petitioner may be informed that at least
the memorial concerns a matter of internal administration and discloses no grounds for
action by the Hon'ble the Agent to the Governor General

By order

RAJPUTANA AGENCY OFFICE

Dated, Abu, the

192 . }

Secretary to the Hon'ble the Agent
to the Governor General
Rajputana.

English petition dated the 192
Vernacular

from

addressed to the

No V.

[See Government of India Standard Form S S]

No. V. of 192 .

Forwarded to the

for disposal, in continuation of this office endorsement No V, dated the
 192 , with the request that the petitioner may be informed
 that no further petition on the same subject will be attended to unless submitted through
 him

By order,

RAJPUTANA AGENCY OFFICE .	}	Secretary to the Agent to the Governor General Rajputana.
Dated Abu, the 192 .		

English
 Vernacular petition dated the 192 ,

from

No. VI.

RAJPUTANA AGENCY.

MEMORANDUM

No V of 192 .

Dated Abu, the 192 .

With reference to his application, dated the 192,

is informed that there is no post vacant to which he can be appointed.

Secretary to the Agent to the
 Governor General in Rajputana,

No VII

RAJPUTANA AGENCY.

MEMORANDUM.

No V OF 192

Dated Abu, the 192

With reference to his application, dated the 192 ,

is informed that appointments in the Indian States of Rajputana are in the gift of the Darbars whom the applicant may address direct, should he be so advised

*Secretary to the Agent to the
Governor-General in Rajputana*

To

No VIII

[See Government of India Standard Form S-8]

DRAFT ENDORSEMENT.

FOR APPROVAL.

No of 192 .

By order,

RAJPUTANA AGENCY,
MOUNT ABU,

The 192 . }

*Secretary to the Agent to the
Governor-General, Rajputana*

LIST OF PAPERS FORWARDED.

No. 1N

[See Government of India Standard Form S (Agency) 37.]

RAJPUTANA AGENCY.

No V, dated Abu, the 192

With reference to his petition, dated the 192 ,

is informed that it has been forwarded to

for disposal, and that no notice will be taken of future petitions relating to the same subject unless forwarded through that officer.

Secretary to the Hon'ble the
Agent to the Governor-General,
Rajputana

No. X.

No. V. of 192 .

Dated Abu, the 192 .

To

With reference to his petition, dated 192 ,
her
their

is
are informed that the petition has been forwarded to the
for disposal.

Secretary to the Hon'ble the
Agent to the Governor-General,
Rajputana

No XI

No V of 192 .

From

Secretary to the Hon^{ble} the Agent
to the Governor General, Rajputana

To

*Dated Mount Abu the**192 .*

In reply to your letter dated the *192* , I am desured to inform you that an appointment in a State in Rajputana is in the gift of the Darbar concerned and that under the rules the Hon^{ble} the Agent to the Governor General is unable to make recommendations unless the Darbar invite such recommendations

*Secretary to the Hon^{ble} the Agent
to the Governor General, Rajputana*

No XII

RAJPUTANA AGENCY

 MEMORANDUM

No V of 192 ,

Dated Mount Abu, the 192 .

With reference to his application, dated the *192* ,

is informed that no appointment suitable for him is at present vacant but that his name has been noted

*Secretary to the Agent to the
Governor General, Rajputana*

To

APPENDIX V

RULES FOR THE DESTRUCTION OF RECORDS AND PERIODS FOR WHICH THEY ARE TO BE KEPT

(Paragraph 133)

1 The broad principle is that no papers which are likely to be of any value, at any

destroyed. These rules however do not exhaustively state the classes of papers which have to be dealt with. They are merely intended to indicate in a general way the plan on which destruction should be carried out. In all doubtful cases reference to be made to Registrar and if necessary to the Secretary.

2 *Classes of papers to be preserved*—Documents belonging to any of the descriptions noted below, shall be exempt from destruction, that is to say—

- (a) Documents relating to the rights of Government, or to matters of political or administrative importance
- (b) Documents prepared under the provisions of any enactment which appear to contemplate their being preserved
- (c) Documents the destruction of which would prejudice the rights or affect the liabilities of individuals

- (e) Standing orders of every description
- (f) Originals of legal documents or documents of legal value
- (g) Discussions relating to important public services
- (h) Papers which are important or likely to become important in the future, from a historical, biographical or other important point of view
- (i) Treaties, agreements and such like documents
- (j) Original letters on important matters from the Indian Princes and Chiefs
- (k) Papers containing discussions of important principles or policy
- (l) Maps sketches and similar papers relating to important or disputed questions
- (m) Gazettes of India.

3 *Classes of papers to be destroyed*—Papers not falling under any of the heads mentioned in the foregoing paragraph

- (a)

- (l) Periodical Returns and papers of the kind described in the lists attached.

- (c) *Office Registers*—Registers of letters received and issued should never be destroyed, but bound up for each year and Agency separately, the registers of issues and receipts being bound separately. Registers, e.g., Station Dak Book, Stamp Register, Register of unofficial cases may be destroyed after three years.

- (d) *General correspondence*—Unless the subject is of a very trivial nature and the Head of the Office is certain that it will never be referred to again and the space for keeping records is limited, no letters should be destroyed but

re-paged and indexed,
following papers which

Manuscripts of all printed papers except such as are mentioned in paragraph 2 above

(e) *Printed papers*—More than six spare copies of printed papers

4 When the cases have been weeded according to the foregoing instructions, a list of papers intended for destruction will be prepared and submitted to Registrar. He may order the destruction of such cases which can under the rules be destroyed after one, two and three years. The cases of the remaining classes will be put up for the orders of the Secretary.

5 When orders have been passed for destruction, Registrar should see that the cases are destroyed and that note to that effect is made by the insertion against the entry of a capital "D" in red ink in the File Register. As soon as a case is ready for restoration to the record room, a slip indicating the time for which it is intended to be retained according to the classification given to it in the lists should be attached. All doubtful cases will be referred to the Secretary for orders.

6. Printed slips (size 3 inches by 2 inches) coloured red for files to be retained permanently and blue for files to be kept for a period as noted below will be affixed —

To be retained until

To be retained
permanently

months, years.

7 The Record keeper will be responsible that no weeded files are received by him in the Record Room unless slips are affixed to them, and he will also maintain a register showing files which are to be destroyed in a particular year.

8 When there is doubt whether a case should be destroyed or retained, it should be retained.

9 A list of papers suggested for destruction will be prepared and submitted to the Registrar or the Secretary as the case may be.

10 Papers which have to be destroyed shall be collected and burnt under proper supervision.

11 Important files ordered by the Secretary or the Agent to the Governor General to be printed, should be weeded first and then sent to one of the Government Presses for printing. Proofs should be compared in the branch concerned after which the printed files will be restored to the Record.

I

List of papers to be destroyed after 1 year

-
- 1 Applications for copies of papers
 - 2 Distribution of books and pamphlets
 - 3 Selections from vernacular papers
 - 4 Half yearly return of leave contemplated by Political Officers
 - 5 Petitions (unimportant)
 - 6 Service of Summons
 - 7 Miscellaneous notices regarding unclaimed property and stolen cattle, etc
 - 8 Weather and Crop report
 - 9 Applications for re appropriation of Budget grant.
 - 10 Indents for Stationery
 - 11 Programmes of Viceroys's tours
 - 12 Meteorological returns
 - 13 Weekly plague and cholera returns
 14. Applications for interview
 - 15 Movements of troops (Routine papers)
 - 16 Dockets forwarding forest reports of other Administrations
 - 17 Distribution of annual reports and publications of other Administrations
 - 18 Railway accident reports.
 - 19 Grants of Casual leave
 - 20 Commissariate and other notices
 - 21 List of unanswered references.

II

List of papers to be destroyed after 2 years

-
- 1 Prices current, weekly, fortnightly and monthly
 - 2 Summonses, Interrogatories, Notices and Mukhtarnamas (if not connected with any judicial case within the Administration.)
 - 3 Dockets to letters or forwarding statements or returns.
 - 4 Camp Chalanas.
 5. Returns of commodities carried by railway
 6. Returns of railway accidents (other than those at No. 18 in List I above.)
 - 7 Monthly jail returns.
 8. Certificate of "No marriage" by Registrar

III

List of cases to be destroyed after 3 years

- 1 List of title holders
- 2 Petitions finally disposed of
- 3 Addenda and corrigenda to list of title-holders and leading officials, etc
- 4 Annual statement of English, Foreign, Anglo-Vernacular, and Vernacular news papers published in India and Burma as supplied by Criminal Intelligence Department
- 5 Returns of work performed by the subordinate courts.
- 6 Deportation of Vahaties.
- 7 Correspondence regarding maintenance of lunatics.
- 8 Death report of prisoners.
- 9 Absolute release of transmarine convicts.
- 10 Death reports of transmarine convicts.
- 11 Yearly returns of arms and ammunition.
- 12 Annual statement of Excise receipts and charges.
- 13 Annual Statement of receipts and charges under stamps
- 14 Applications for refunds of value of spoiled stamps.
- 15 References under the Income Tax Act.
- 16 Return of Excise duty on country made cloth.
- 17 Return of Excise in Cantonment's.
- 18 Forecasts of opium.
- 19 Applications for additional budget grants.
- 20 Arrears of pay and pension.
- 21 Office copies of salary, contingent and travelling allowance bills.
- 22 Cases of treasure trove
- 23 Civil Accounts.
- 24 Compensation for dearness of grain.
- 25 Daily allowance
- 26 Land revenue accounts
- 27 Purchase of articles charged to contingent grant.
- 28 Purchase of books.
- 29 Purchase of furniture
- 30 Question of leave allowances.
- 31 Question of pay and allowances to individual officers.
- 32 Cantonment and Local Fund budgets.
- 33 Petitions against local and cantonment rules.
- 34 Report on the working of mints.
- 35 Post Office
- 36 Wrongful use of service postage stamps
- 37 Telegrams erroneously classed as "State."

III—*concl'd**List of cases to be destroyed after 3 years—cont'd*

- 38 Agricultural statistics
- 39 Minerals and gem returns
- 40 Crop forecasts
- 41 All returns which appear in volumes of statistics compiled by the Director General of Statistics
- 42 Returns of large industries
- 43 Report on mines
- 44 Horse fair reports and appointment of judging Committee
- 45 Applications for appointments.
- 46 Returns of Europeans and Eurasians employed in Indian States.
- 47 Return of boundary pillars
- 48 Changes in Cantonment establishments
- 49 Petitions applying for patronage for books or translations.
- 50 Liveries for peons
- 51 Papers relating to privilege leave of gazetted officers.
- 52 Gazettes of other Local Governments.
- 53 Charge Allowances
- 54 Extension of joining time
- 55 Deputation allowances
- 56 All monthly, quarterly, half yearly and yearly returns
- 57 Miscellaneous papers connected with Census returns.
- 58 Diaries
- 59 Budgets Local Funds Cantonments and Indian States (the orders on them being retained)
- 60 Returns of prisoners or any returns of casualties in jails.

IV

List of cases to be destroyed after 5 years.

-
- 1 Inspector General Indian State Forces—Annual Reports .
 - 2 Permission to Indian Chiefs and Princes to visit Hill Stations
 - 3 Appointment of Cadets to Imperial Cadet Corps (or three years after they have left the Corps)
 - 4 Memorials
 - 5 Extradition of prisoners
 - 6 Payment of fees to Government pleaders
 - 7 Appeals against capital sentences
 - 8 Conditional release of transmarine convicts
 - 9 Licenses granted under the Arms Act
 - 10 Licences to solemnize marriages
 - 11 List of holidays
 - 12 Orders prescribing the headquarters for purposes of travelling allowance of an officer on deputation
 - 13 Camp equipage and carriage
 - 14 Exchange compensation allowance cases
 - 15 Agent Governor General's tour programme
 - 16 Papers relating to the grant of leave other than leave on full average pay to gazetted officers
 - 17 Papers and notifications relating to appointment of officers
 - 18 Statement of memorials withheld
-

V

List of cases to be destroyed after 10 years

-
- 1 Armament returns
 - 2 Investment of Cantonment Magistrate with Powers
-

VI

List of cases to be destroyed after 20 years

-
- 1 Pension and gratuity cases excepting those in connection with which any general ruling has arisen.
-

APPENDIX VI

Instructions regarding forms (Vide Appendix I)

FORM A—A separate register should be kept for each branch, the office and branch should be entered on the cover. The date of receipt should be written in red ink across the register above the entries of that date.

FORM B—A separate register should be kept for each branch, the office and branch should be entered on the cover.

Column 1—Unless columns 4 to 7 are omitted, sufficient space must be left between each entry in column 1 to allow for probable entries in columns 4 to 7, if it is subsequently found that the space allowed is insufficient a fresh entry should be begun later in the book and a reference should be made to it across columns 9 to 12 of the original entry, e.g., "continued after No 453". The original number should be kept in the new entry, the entries already made in columns 4 to 7 need not be repeated against the new entry.

Column 3—The 'subject' should be the same as in the index.

Columns 4 & 6—Authorised abbreviations should be used.

Columns 4 to 7—These columns may be omitted with the sanction of the Secretary, provided that

- (a) every receipt after the first is registered in the notes of the file by, or under the direct personal supervision of the record keeper or head clerk, and
- (b) in the case of a file not kept in the office when it is closed the person to whom it is sent and the date of despatch shall be entered across the last two columns of the Register of Files.

Column 9—These entries are to facilitate destruction.

R = Retain.

D (Date) = Destroy in the year mentioned.

N = Nothing retained in office, i.e., the whole file was finally sent away on the date of closing and is not expected back.

If different parts of the file are to be destroyed in different years, this should be stated in column 9, e.g.,

"I. R, II D (1933)"

would mean that part I is to be retained indefinitely and Part II is to be destroyed in 1933.

Column 10—If the register is maintained by the record keeper column 10 may be omitted.

Alternative to Form B—Form B may be divided into two registers in Form B 1 and Form B 2. Form B 1 would be a register of Files in the same form as Form B omitting columns 4 to 7. The Register should be kept by the record keeper. No space need be left between entries in column 1. Form B 2 would be register of correspondence in Files in the Forms of Columns 1, 4 to 8; Column 10 should be added if the Register is kept by the head clerk and not by the record keeper. The remarks above about column 1 of form B will apply to Column 1 of form B 2.

The instructions relating to form B in paragraph 32 will apply to both registers.

A receipt beginning a new file would involve entries in both registers (paragraph 51), subsequent correspondence in the same file in the same manner. (Paragraphs 131 and 132).

FORM C—A separate register should be kept for each branch, the office and branch should be entered on the cover

Column 1—In this column should be entered the number of the permanent file in which the periodical is filed

Columns 4 and 6—Authorised abbreviations should be used, one line should be allowed for each individual *e g*, if a return is due from all Political officers, each should be entered on a separate line

Columns 1 to 4, 6, 7 should be filled up at the beginning of the year

FORM D—(Government of India Standard Form S 97)—The letter of the branch should be entered in the bracket after the word "Branch".

The "subject" should be the same as in the index.

Trade Returns and Commercial History.

CAMPHOR.

Camphor obtained from Borneo from the trunk of *Dryobalanops* *aromatica*, and by sublimation from *Med*, 222). Dr. accepts this opinion the Sanskrit writer modern Camphor

HISTORY.

the Sanskrit writers, and Camphor referred to may be which at the period India or imported from appear to have been sufficient the strongly camphoraceous in the first plant resorted to as a substitute or adulterant for the prized Camphor of Sumatra. As a matter of fact, this Camphor is much more nearly related to the Malayan than to the China Camphor, and even at the present day it is ten times the price of the Formosa Camphor, and is extensively consumed in China, partly as a medicine and partly in perfuming the finer qualities of Chinese ink. Moodeen Sheriff mentions four kinds of Camphor as met with in the bazars of South India, viz., (a) *Kāfūre-gaisūrī*, (b) *Sārati kāfūr*, (c) *Chini-kāfūr*, and (d) *Batāi-kāfūr*.

TRADE RETURNS AND COMMERCIAL HISTORY.

TRADE.
263

Commerce.—While some of the less important camphors do, to a limited extent, reach Europe and India, the commercial or Chinese form is that which has been called "Common Camphor." This arrives at the English and Indian markets chiefly in a crude state, and is in both countries resublimed.

than the Formosa Camphor.

small
uses,

view
the

chiefly from China, is worth not more than R40 to R65 per cwt. This enormous difference is accounted for by the reputation (scarcely merited) which the Bhimsaini kind enjoys of peculiar excellence" (*Para. 16, pages 9 and 10*)

Of Borneo and Sumatra Camphor probably not more than 2 or 3 cwt. are annually imported into India.

CAMPHOR.

Trade Returns and Commercial History.

INDIAN
TRADE IN
CAMPHOR

The Import and Re-export trade in Camphor between India and foreign countries for the past seven years was as follows —

YEAR	VALUE OF CAMPHOR			
	IMPORTED INTO INDIA		RE EXPORTED FROM INDIA	
	Bhimsaini or Barus	Other kinds	Bhimsaini or Barus	Other kinds
	R	R	R	R
1879-80	20,909	5,34 001	2,316	23,174
1880-81	22,924	5 53 732	140	26,559
1881-82	38 574	5,52,335	1,640	21,138
1882-83	43 618	8 68 794	529	25 231
1883-84	38,579	6,27,278	790	28 730
1884-85	35 501	6 83 333	270	13 432
1885-86	25,944	6,53,545	Nil	16,779

YEAR.	VALUE	ANALYSIS OF EXPORTS FOR 1885-86	
		Country to which exported	Province from which exported
	R		
1879-80	7,574		
1880-81	7,142		
1881-82	6,510		
1882-83	9,475		
1883-84	6 682		
1884-85	6 135		
1885-86	6 055		
		Ceylon	Bombay
		Other Countries	Madras
		TOTAL	TOTAL

Indian
Refined
264

the process as practised in Bombay "The process of resublimation is a peculiar one, the object being to get as much interstitial water as possible into the camphor cake. The vessel used is a tinned cylindrical copper drum, one end of which is removable; into this is put 14 parts of crude camphor and 2½ parts of water, the cover is then luted with clay, and the drum, being placed upon a small furnace made of clay, is also luted to the top of the furnace. In Bombay four of these furnaces are

Purification of Camphor.

CAMPHOR.

Ind., 1st Ed., 549). This same practice seems to be followed at Delhi and at a few other cities in India, but the method is crude and unsatisfactory, when the purified article is compared with that imported into India from Europe. The European process of refining camphor was long kept a secret, and towards the end of the seventeenth century the entire camphor of Europe had to be sent to Holland to be sublimed. A monopoly was also held for some time in Venice, but at the present day camphor-refining is largely accomplished in England, Holland, Hamburg, Paris, New York, and Philadelphia.

European
Refined.
265

by means of a fire, where flame might ignite the gas given off during the process of sublimation, dishes of fusible metal, kept warm by a furnace below the room, are used. The heat is suddenly raised from 120° to 190° C, and kept at that point for half an hour, so as to expel the water from the camphor. The temperature is then raised to 204° C, and maintained at that point for 24 hours. When the crude camphor has melted, the sand

The *rationale* of the process consists in preserving the temperature uniformly at the point of volatilization; the quicklime retains resin or empyreumatic oil, the iron fixes on any sulphur that may be present,

Camphor
Plants.
266

India. In the report of the 12-83 it is mentioned that a re well. It seems likely that, rupees worth of China Camphor, since there is every reason were made, the tree could be

successfully introduced. The amount of Camphor consumed in

CAMPHOR.

Chemical Formula for Camphor.

India is not sufficiently great to tempt experiments being undertaken with *Dryobalanops Camphora*, but the extended cultivation and manufacture of *Blumea* and *China Camphors* would seem highly desirable.

CAMPHOR OIL.

OIL
267

Oil of Camphor—There are two very distinct substances known by that name in commerce. The first and most important is the oleo-resin or camphor oil of Borneo. This is obtained by tapping the trees. Some times this accumulates to such an extent that (as with the South American *copaiba* tree) the trunk, no more able to resist the pressure of the fluid, spontaneously bursts open or has its tissue broken into large internal chambers, producing while this occurs a loud noise, "as if the tree were rent in twain." The *Pharmacographia* states that Motley, in cutting

distinct and should not be
phor-oil of Formosa. This
is a brown liquid, holding in solution an abundance of common camphor,

CHEMICAL AND MEDICAL PROPERTIES OF CAMPHOR

268

Chemistry—It is not necessary to enter into this subject in great detail. For a full account of the chemistry of Camphor the reader is referred to works on chemistry, but more particularly to the *Pharmacographia* and the *United States Dispensatory*, as these are more likely to be accessible than the numerous and scattered papers in which this subject

camphor. When mixed with resins or concrete oils, camphor often partially or completely loses its odour. The formula given for this form of camphor is $C_{10}H_{16}O$, by treatment with various reagents it yields a number of interesting products. Prolonged boiling with nitric acid oxidises the camphor into *Camphoric acid*, $C_{10}H_{14}O_4$, and *Camphoronic acid*, $C_8H_8O_6$, water and carbonic acid being eliminated. When repeatedly distilled with chloride of zinc, it is converted into *Cymene* or *Cymol*, $C_{10}H_{16}$.

$C_{10}H_{16}O$. It is somewhat does not consequently it. It is also heavier, having the sp gr 1.009. It is easily pulverised without the aid of alcohol, it is, in fact, a more compact and brittle substance than ordinary

Medicinal Properties of Camphor.

CAMPHOR.

camphor It requires for fusion 198° C. In optical properties an alcoholic solution is found to be $12\frac{1}{2}^{\circ}$ dextrogyre. By the action of nitric acid, camphor is converted into camphoric acid, and by continued oxidation, it is converted into camphoric acid, and by continued oxidation, it is converted into camphoric acid, and by continued oxidation, it is converted into camphoric acid.

CHEMISTRY.

and differe
remains
phor is converted into ordinary camphor

MEDICINE.
269

secondary, that of a sedative, anodyne, and antispasmodic. In large doses it is an acro-narcotic poison. Camphor has been extensively used in the advanced stages of fevers and inflammation, insanity, asthma, angina pectoris, hooping-cough, and palpitations connected with hypertrophy of the heart, affections of the genito-urinary system, comprising dysmenorrhœa, nymphomania, spermatorrhœa, cancer, and irritable states of the uterus, chordee, incontinence of urine, hysteria, rheumatism, gangrene, and gout. It has also been employed as an antidote to strychnia, but with doubtful results. It is regarded as a medicine in impotence.

be discussed here at great detail. The reader is therefore referred to the *Pharmacopœia of India*, pp 190, 192, and other standard works on materia medica. As having a special bearing on India, however, the following extract may be republished from Waring's most useful little book, *Basar Medicines*—

"In chronic rheumatism, in addition to its use externally, it may be

Care, however, is necessary to prevent the patient inhaling the vapour, which is of comparatively little consequence when simple water is used.

"In asthma, camphor in 4-grain doses, with an equal quantity of asafoetida, in the form of pill, repeated every second or third hour during a paroxysm, affords in some instances great relief. Turpentine stupes to the chest should be used at the same time. Many cases of difficulty of breathing are relieved by the same means. These pills also sometimes
childhood, cam-
chest at nights,
the strength of
addition of some

bland oil

"In rheumatic and nervous headaches, a very useful application is one ounce of camphor dissolved in a pint of vinegar, and then diluted with one or two parts of water. Cloths saturated with it should be kept constantly to the part.

"In spermatorrhœa, and in all involuntary seminal discharges, no

CAMPHOR

Medical Properties of Camphor.

MEDICINE

medicine is more generally useful than camphor in doses of 4 grains

pill twice or three times a day, according to the severity of the symptoms, will sometimes afford great relief. In each of these cases it is important to keep the bowels freely open.

"In painful affections of the uterus, camphor in 6 or 8-grain doses often affords much relief. The liniment should at the same time be well

rubbed over the region of the heart. It should be discontinued if it causes headache or increased heat of the scalp. Its use requires much discrimination and caution.

"To prevent bed sores, it is advisable to make a strong solution of camphor in alcohol and have the patient bathe in it to bathe, likely to

process of

Assistant Surgeon Jaswant Rai, Multan). "It is an irritant and rubefacient, good for a cold in the head with coryza, summer diarrhoea" (Brigade Surgeon W. R. Rice, Jubbulpore). "Largely used as a liniment for muscular pains. Is a good expectorant" (Surgeon R. Gray, Lahore). "Used in 3 or 4-grain doses and mixed with about $\frac{1}{4}$ grain of extract of belladonna. I have found this to be of very great value in neuralgic pains."

Ilang-ilang.

CANANGA
odorata.

MEDICINE.

preserves clothing and other articles against insects and worms" (Surgeon Shib Chunder Bhattacharji, Chanda, Central Provinces) "Useful in cholera" (Surgeon H. D. Masani, Karachi). "In the form of spirit, camphor is very effective in cholera passages. In the case of cholera, its use, and think of it as a remedy." (Surgeon S. H. Browne). "that when given in 10-grain doses every fourth hour in cholera, good results are obtained."

Assistant Chuna Lal, Jubbulpore) "Is taken in large doses to procure abortion" (Surgeon-Major D. R. Thompson, Madras) "Camphor is daily used as a stimulant, antispasmodic, sedative to the genito-urinary system, and parasiticide. The spirit of camphor is a useful remedy in cholera, in 1 to 5-drop doses" (Assistant Surgeon Nundo Lal Ghose, Bankipur) "Camphor Used in 3 or 4-grain doses and mixed with about $\frac{1}{2}$ grain of extract of belladonna. I have found this to be of very great value in cholera." (Assistant Surgeon N. D. Ghose, Bankipur)

DOMESTIC
270

but it is a camphorated soap greatly to enable them to take root when placed in the soil

Camphora glandulifera, Nees, see *Cinnamomum glanduliferum*, Meisn.; LAURINEÆ

Canada Balsam, see *Abies balsamea*, Aiton.; CONIFERÆ.

CANANGA, Rumph.; Gen Pl, I, 24.

Cananga odorata, H. f. & T. T., Fl Br. Ind., I, 56; ANONACEÆ

271

THE ILANG-ILANG of European perfumers.

Syn.—UVARIA ODORATA, Lamb

Vern.—"Ilang-ilang" (Tagalog), "Ilang-ilang" (Hindi)

Refere

371

Encyclopædia, I, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100

CANARIUM
commune.

Java Almond.

ILANG-
ILANG

Habitat.—A large evergreen tree of Burma (Ava and Tenasserim), distributed to Java and the Philippines Cultivated in many parts of

OIL.
272

273

CANARIUM, Linn.; Gen Pl, I., 324
Canarium bengalense, Roxb; Fl Br Ind., I, 534; BURSERACEÆ

Vern —Gogul dhap, NEPAL, Narockpa, LEPCHA, Tekreng, GARO; Bis-jang, dhuna, ASS

References —P. 1, 5, 12, 13, C B C, Kurr, For Fl Burma, I, Hort Sub Cal, 149, Him Bot, 177, Cooke,

GUM.
274

Habitat.—A tall tree, with a straight cylindrical stem, it is met with in the eastern moist zone, eastern Himalaya, Bengal, and Burma

Gum.—Yields a brittle, amber-coloured resin, resembling copal, which is used as incense The natives set little value on it In Calcutta bazars

TIMBER
275MEDICINE.
276

swellings

FOOD

Food —"Fruit edible

277
TIMBER.

Structure of the Wood —"Strong and durable, used for common house building" (Trimen).

278

279

C. commune, Linn; Fl Br Ind, I, 531.

JAVA ALMOND TREE.

Vern —Jangali badam, HIND, Jangali bādāna, CUTCH, Kagli mara, kagga libiya, java badamiyanne, KAN, Canari, MALA, Rata kakana, SING.

References —Roxb, Fl Ind, Ed C B C, 504, Vogt, Hort Sub Cal.

GUM.
280

intr

long

Phc

as

C. 280

Bengal Incense: Elimi.

CANARIUM
commune.

Blanco, a botanist of Manilla, described in 1845 under the name *Isica Abilo*, but which is completely unknown to the botanists of Europe. Blanco's description is such that

in either of the old genera *Isica* or
and Hooker in that of *Bursera*,
in fact, even the order to which it belongs is somewhat doubtful.

"Manilla Elemi is a soft, resinous substance, of granular consistence
more
enders
impure
yellow

Manilla Elem.
281

tint. It has a strong and pleasant
yet withal somewhat terebinthine

higher temperature fuses into a clear

(15th Ed.), page 536, says "The
Manilla Elemi is conjecturally referred to *Canarium commune*." In their
Medicinal Plants Bentley and Trimen give a detailed description of
the plant. They say "It is also cultivated in India, and has been

by Blanco, should be even referred to the *BURSERACEÆ*.

The gum is used principally in the manufacture of varnishes, also in
felting and in medicine.

Oil.—The nut yields a semi-solid oil on expression, similar in appear-
ance to cocoanut oil. It is used for culinary purposes, and is regarded
palatable.

OIL
282

§
tine
to pc
C. f.

Bombay)

FOOD.
284

Celebes. If eaten fresh or too frequently, the nuts often produce diar-
rhœa (*Drury*).

CANARIUM
strictum

Black Dammar Tree

285

Canarium strictum, Roxb, Fl Br Ind, I, 534, Beddome, t 128

THE BLACK DAMMAR TREE

Vern

Reference

GUM
286

Habitat —A tall tree of South India. Common about Courtallum in the Tinnevely district and in Kanara.

Gum —It yields a brilliant resin called the Black Dammar of South

ten years between the months of April and November, and the resin is collected in January

'This substance occurs in stalactitic masses of a bright shining colour when viewed *en masse* but translucent and of a deep reddish brown colour when held between the eye and the light, homogeneous with a vitreous fracture, partially soluble in boiling alcohol, and completely so in oil of turpentine (*Pharm Ind*)

BLACK
DAMMAR
287

the manufacture of bottling wax varnishes, &c. Its colour when in solution is pale if compared with its dark tint when in mass. Thus, though insoluble in spirit, its solution in turpentine forms a tolerable varnish. When submitted to destructive distillation it yields about 78 per cent of oil, resembling that obtained from common colophony, but I fear, in the majority of its possible applications, it possesses few advantages over

as the

nearly

suit

common

with

t and

colourless as glass, in such amount that a single firm turns out 60 tons per week "

MEDICINE
Burgundy
Pitch
288

Medicine —The resin is used medicinally, according to Dr Bidie, as a substitute for Burgundy Pitch in making plasters

Special Opinions —§ 'Bathing in a tub painted inside with dammar is supposed to relieve the irritation of prickly heat' (*Surgeon Major A S G Jayakir, Muskat Arabia*) 'Employed as a liniment with gingelly oil, in rheumatic pains' (*Surgeon Major F F L Ratton, Salem*)

C. 288

The Sword bean.

CANAVALLIA
ensiformisCANAVALLIA, *Adins* (PDC); *Gen Pl*, I, 537Canavalia ensiformis, DC; *Fl Br. Ind.*, II., 195; *Wight*, Ic, t
753; LEGUMINOSÆ.

SWORD BEAN Sometimes called PATAGONIAN BEAN.

Syn — C. GLADIATA, DC, DOLICHOS GLADIATUS, Willd., as in Roxb., *Fl.*
Ind., Ed C B C., 559, D ENSIFORMIS LinnVern — *Makham shim*, *mekhun*, BENG., *Tihon*, SANTAL., *Sufed or Idi kud*
sumbal, HIND., *Sem*, PB & N-W P., *Garari*, MAR., *Gairara*, BOM.,References — *Thwaites*, *En. Ceylon Pl.*, 88 *Dals & Gibs*, *Bomb Fl.*,
C. B. C. 559, D

p 144 Fig 27

suspicious belief that it will protect their property from plunder (*Smith*)There are several forms of this plant met with in India, the seeds and
flowers being of different colours (*Drury*) These according to the *Flora*
of *British India*, are referred to three distinct varieties —Var 1st, *virosa*, W & A, *Prod.*, 253, *Dals & Gibs*, *Bomb Fl.*,
69, *Dolichos virosus*, *Roxb.*, *Fl Ind.*, Ed C B C., 559 Pods often 2-4
inches long, 4-6-seeded Speaking of this form, *Roxburgh* says 'I do not
find that any part of this species is in any shape useful to the natives or
others, indeed, the natives of *Coromandel*, where the plant is common,
reckon it poisonous, which is corroborated by *Van Rheede*' This is
known in *Bengal* as *Kath-shim*, or *Kala-shim* and *Garwara* (*Gowara*) in
Bombay

290

Var 2nd, *turgida*, *Grah in Wall Cat* C *Stocks*, *Dals & Gibs*,
Bomb Fl., 69 Pods large and turgid, 3 to 5 inches by 1½ to 2 inches

291

Var 3rd, *molis* *Wall* Found in Southern and Western India The
pods are smaller than in either of the above, when cultivated they are
tender and eaten like French beans.

292

Food — The young, tender, half grown pods, apparently of only var
3 are actually eaten, but these constitute the so-called French beans at
the tables of Europeans Natives also eat them in curry The form
with large white seeds is considered the most wholesome Some five
varieties are reported to be cultivated in *Lucknow*, of which the form
known as *hulwa*, a white narrow-podded variety, is considered the best Mr
Cameron informs the writer that the seeds of this pulse are highly
relished in *Mysor* *Atkinson* writes of the North-West Provinces that
the *sem* is "consumed by all classes"

FOOD

293

Professor *Church* gives the analysis of this pulse (p 144), and adds
that its nutrient ratio is 1 : 2 : 2 and the nutrient value 80

H

C. 293

CANES

White Cinnamon, Canes

294

Canavalia obtusifolia, DC, Fl Br Ind, II, 196References — *Thwaites En, Ceylon Pl, 88, Voigt, Hort Sub Cal, 235, Drury, Us Pl 125, Balfour, Cyclop, Kew Cat, 44*

Habitat — Met with on the coasts of the Western Peninsula, Ceylon, and the Malaya Peninsula

"Is a useful binder of loose sand" (*Balfour*)

295

CANELLA, Sw, Gen Pl, I, 121, 970**Canella alba, Murray, DC Prod, I, 563; CANELLACEÆ**

WHITE CINNAMON Eng, CANELLE BLANCH, Fr, WEISSER ZIMMET, Germ CANFLA BIANCA, It CANFLA ALBA, Sp CANFLA BLANCA Sp

References — *Voigt Hort Sub Cal 88, Pharm Ind, 25, Fluck & Hand, Pharmacog 73, U S Dispens, 15th Ed, 337, Year Book of Pharmacy 1873 p 43, Spens Encyclop 1419, Smith Dic, 84, Treasury of Botany Hanbury Sc Papers, 353, Kew Cat 14*

Habitat — A West Indian aromatic plant, the bark of which is imported into India, and is sold by druggists, the tree might be cultivated in India

OIL

296

Oil — "An essential oil, erroneously called 'white cinnamon,' is obtained by the aqueous distillation of the bark, it is a mixture of caryophyllie (engenic) acid, an oil resembling cajuput, and an oxygenised oil" (*Spens, Encyclop*) It is a rare article, not known to commerce

MEDICINE

Bark

297

Medicine — The bark is met with in rolls or quills two or three feet in length, having a bitterish acid peppery taste. The odour is something like a mixture of cloves and cinnamon. The bark is an aromatic stimulant used to a limited extent in combination with other articles in constitutional debility, dyspepsia, scurvy, &c (*Pharm Ind*). In the West Indies it is used as a condiment and has some reputation as an antiscorbutic

CANES.

CANES

298

Canes

CANNE, Fr, ROHR Germ, Bhate HIND, Nathur, GuzThe species of the genus *Calamus* — a genus of climbing palms — yields the canes of commerce. Few plants are more useful to the hill tribes of India and the Malay than are the various forms of cane yet very little of a definite nature is known as to the peculiar properties and uses of the individual species. They afford 'Dragon's blood' and the "Malacca" and "Rattan Canes" of commerce but it is probable that each of these articles is obtained from more than one species of *Calamus*. Reeds and small bamboos are sometimes, but incorrectly, spoken of as canesThe species of *Calamus* are delicate gr stunted or times, by

trees of the forest, they ascend as gigantic climbers, often attaining to as much as 600 feet in length. The stems, leaves, and tendrils are covered with spines and prickles. The fruit hangs in great clusters, the inner

Canes often 600 feet long

C. 298

Asiatic Uses of Canes.

CANES.

pieces. The roots and young sprouts are eaten as vegetables and somewhat resemble asparagus. Canes owe their value to their great strength, and more particularly to the strength of the outer layer of woody structure. As substitutes for ropes they are invaluable, and in some respects even superior to ordinary ropes. For walking sticks and canes, and for spear and lance shafts, they are in great demand and are justly popular, lightness, strength, and uniform structure and size, are properties of the greatest importance.

Substitutes
for Ropes
200
Shafts.
300

Cane-bridges
301

parallel canes forming the pathway, the canes being knit together with bamboo or bark, so as to constitute a band not more than 18 inches in breadth, through which the rushing water may be seen below. The railing affords additional support, it consists of two canes carried about three or four feet above the pathway, one on either side. These are here and there connected by perpendicular canes passing under the pathway, and the whole structure is bound together by a network of bark-ropes or smaller canes. With the weight of the traveller the bridge bends until it is often alarmingly near the water, and to prevent the railing closing on the person crossing the bridge, barriers are thrown across here and there, about 18 inches above the pathway, similar stays are also carried over head. These barriers constitute the chief difficulty in crossing a cane bridge, for on raising the foot, the swaying structure and the rushing

Bridges.

Ropes

and indeed throughout the Eastern Islands, vessels are furnished with cables formed of cane twisted or platted. This sort of cable was formerly extensively manufactured at Malacca (*Royle, Fibrous Plants*). Dampier says "Here we made two new cables of rattans, each of them four inches about. Our captain bought the rattans, and hired a Chinese to

them down, nor can we carry them out but by placing two or three boats at some distance asunder, to buoy up the cable while the long boat rows

CANES.

European Uses of Canes.

entire and cut Useful chairs, sofas, and couches are made all over India from cane, and cane *punkha* ropes are almost in universal use. In Bengal baskets (*dhama*) are made of entire canes by twisting the canes round

gether, by means of cane-strings, the canes being arranged so as to be flat and parallel.

THE EUROPEAN USES OF CANES

They are valued on account

They are extensively used as

as a substitute for whalebone

such ribs costing only from 1*d* to 2½*d* instead of 2*s* 6*d* to 3*s* for whalebone.

Cane is also extensively employed in saddlery and harness, and a wicker-work of rattan is now used in the construction of the German military helmet

of the central core In Europe this central portion is saved, a patented machine being used to split the rattans which cuts off the outer layer in bands of any required size or thickness, while leaving the central core in the form of a perfectly round and even rod This rod is utilised in the

to reveal, of the fact that the Nagas and other hill tribes of Assam dye human and goats hair a beautiful scarlet, as also tint with the same colour the outer silicious layer of the rattan cane Bands of stained rattan they use for decorating ear rings, bracelets, and leggings

Prepared strips of rattan are extensively used in Europe as in India for caning furniture, but a comparatively new and increasing trade in rattan is the construction of baskets, which are rapidly displacing willow baskets; these are used in cotton-mills, sugar refineries, and other factories, as well as employed extensively by Railway Companies and by gardeners, &c Rattan baskets are peculiarly adapted for carrying carboys containing acids, since the silica of the cane is not acted on by acids (*Spons, Encyclop*) The waste product, after stripping the cane, is, by certain manufactures, reduced to a fibre, and in this form is largely used for stuffing mattresses Cane mattresses are in great favour on the Continent, taking the place of the coir of India

TRADE RETURNS OF CANES

Very little can be learned regarding the internal trade in rattan canes; but, from the fact of the imports (which come chiefly from the Straits Settlements) into Calcutta, Madras, Burma, and Bombay, far exceeding the exports, it seems that with improved facilities of communication a trade might easily be opened up with Eastern Bengal, Assam, and Burma which would to a large extent check the importation, from foreign countries, of a product of which India has herself an unlimited amount The following

Trade Returns

CANES

summary of the foreign trade in Canes and Rattans will be found instructive —

TRADE

Foreign Trade in Canes and Rattans

YEAR.	IMPORTS		EXPORTS AND RE-EXPORTS	
	Quantity	Value	Quantity	Value
	Cwt	R	Cwt	R
1879-80	2067	193035	7483	73582
1880-81	2164	199557	16346	16363
1881-82	27559	292754	23801	206544
1882-83	24603	246476	14244	133061
1883-84	2883	251203	20836	34884
1884-85	33408	30675	1433	133734
1885-86	13	177536	6453	56844

Detail of Imports 1885-86

Province into which imported	Quantity	Value	Country whence imported	Quantity	Value
	Cwt	R		Cwt	R
Bengal	794	6698	Sam	43	358
Bombay and Sind	9871	79095	Straits Settlements	20350	172880
Madras	62	873	Other Countries	450	498
British Burma	2986	23530			
TOTAL	2123	77536	TOTAL	223	77536

Detail of Exports 1885-86

Province from which exported	Quantity	Value	Country to which exported	Quantity	Value
	Cwt	R		Cwt	R
Bengal	155	20770	United Kingdom	3827	35030
Bombay	623	2406	United States	427	8435
Madras	637	54	Italy	63	60
British Burma	3700	3354	Cape Colony	459	628
			Mauritius	87	680
			Other Countries	152	50
TOTAL	20836	34884	TOTAL	6485	56844

The reader is referred for further particulars to the format on given page.

Substitutes
for canes
317
When canes
318

CANES	European Uses of Canes.
	out the anchor" Ropes are regularly made in China by splitting the
Baskets 302	
Chairs 303	
Mats 304	
Cane work 305	chairs made in this way being light and cool A strong and durable floor mat for office purposes is constructed of small entire rattans, bound together, by means of cane-strings, the canes being arranged so as to be flat and parallel.
Walking Sticks 306	THE EUROPEAN USES OF CANES
Umbrella handles 307	They are valued on account They are extensively used as
Umbrella ribs 308	whalebone.
Saddlery 309	and a wicker- nan military
Harness 310	helmet, which is said to make it sword proof But the chief purpose to which cane is put in Europe is in furniture and basket making In India, canes are cut up by hand, the outer strips being separated at the expense of the central core In Europe this central portion is saved, a patented machine being used to split the
Furniture 311	bands of any
Central axis 312	the form of a T construction of
Window blinds 313	perty not possessed by the strong outer bands, namely that it takes with ease any desired colour European authorities do not appear to be aware, however, of the fact that the Nagas and other hill tribes of Assam dye human and goats hair a beautiful scarlet, as also tint with the same colour the outer silicious layer of the rattan cane Bands of stained
Dyed cane 314	
Fibre from cane 315	these are used in cotton-mills sugar refineries, and other factories, as
Cane- mattresses 316	panies and by gardeners, carrying carboys contain- acted on by acids (Spon- ing the cane, is by certain manufactures, reduced to a fibre, and in this form is largely used for stuffing mattresses Cane mattresses are in great favour on the Continent, taking the place of the coir of India
	TRADE RETURNS OF CANES
	Very little can be learned regarding the internal trade in rattan canes;

Trade Returns

CANES

summary of the foreign trade in Canes and Rattans will be found instructive —

TRADE

Foreign Trade in Canes and Rattans

YEAR.	IMPORTS		EXPORTS AND RE-EXPORTS	
	Quantity	Value	Quantity	Value
	Cwt	R	Cwt	R
1879-80	20 617	1 93 035	7 453	73 582
1880-81	21 164	1 99 557	16 346	1 6 363
1881-82	29 559	2 92 754	23 501	2 66 344
1882-83	24 603	2 46 476	14 244	1 33 061
1883-84	28 83	2 5 203	20 836	34 884
1884-85	33 408	3 0 675	14 33	1 33 734
1885-86	21 213	1 77 536	6 453	56 844

Detail of Imports 1885-86

Province to which imported	Quantity	Value	Country whence imported	Quantity	Value
	Cwt	R		Cwt	R
Bengal	7 94	66 98	Siam	4 3	3 58
Bombay and Sind	9 87	79 093	Straits Settlements	20 350	1 72 880
Madras	62	8 7 3	Other Countries	4 0	498
British Burma	2 986	23 530			
TOTAL	21 2 3	77 536	TOTAL	2 3	1 77 536

Detail of Exports 1885-86

Province from which exported	Quantity	Value	Country to which exported	Quantity	Value
	Cwt	R		Cwt	R
Bengal	1 525	20 770	United Kingdom	3 827	35 030
Bombay	6 3	406	United States	427	8 435
Madras	637	54	Italy	63	60
British Burma	3 700	32 354	Cape Colony	469	6 28
			Mauritius	87	880
			Other Countries	1 5 2	5 0
TOTAL	20 836	34 884	TOTAL	6 485	56 844

The reader is referred for further particulars to the information given under the species of Calamus. In concluding this account of Canes it is necessary to briefly mention a few of the more common articles sometimes sold though incorrectly under the name of cane. The most important is the manila —

comparing returns of of grasses

Substitutes
for canes
317
Whan-ee
canes
318

CANES.

European Uses of Canes.

out the anchor" Ropes are regularly made in China by splitting the rattan and twisting it into a rope of a certain thickness. This is used. The small entire and cut

Baskets.

302

Chairs.

303

Mats

304

Cane-work.

305

Walking

Sticks

306

Umbrella

handles

307

Umbrella ribs

308

Saddlery.

309

Harness.

310

Furniture.

311

Central axis

312

Window

blinds

313

Dyed cane

314

gether, by means of cane-strings, the canes being arranged so as to be flat and parallel.

THE EUROPEAN USES OF CANES are even more varied than the Asiatic. They are valued on account of their lightness, flexibility, and strength. They are extensively used as walking-sticks, umbrella handles, and even as a substitute for whalebone for umbrella and parasol ribs, each set of such ribs costing only from 1d to 2½d instead of 2s 6d. to 3s for whalebone. Cane is also extensively employed in saddlery and harness, and a wicker-work of rattan is now used in the construction of the German military helmet, which is said to make it sword proof. But the chief purpose to

construction of fancy
perty not possessed b
ease any desired colour

however, of the fact that the Nagas and other hill tribes of Assam dye human and goats' hair a beautiful scarlet, as also tint with the same colour the outer silicious layer of the rattan cane. Bands of stained

Fibre from

cane

315

Cane-

mattresses.

316

the place of the coir of India

TRADE RETURNS OF CANES

Very little can be learned regard the internal trade in cane;

Indian Hemp.

CANNABIS
sativa.CANNABIS, Linn.; *Gen. Pl.*, III, 357.

331

Cannabis sativa, Linn.; *DC. Prodr.*, XVI, I, 30; URTICACEÆ.

HEMP; INDIAN HEMP; CHANVRE, *Fr.*; HANF, *Germ.*; CANAPE,
It.; KONAFLI, *Rus.*; CANAMO, *Sp.*; HAMP, *Dan.*; KANAS,
Keltic; CANNABIS, *Latin and Greek.*

Syn.—C. INDICA, Lamk.

V-

References—*DC. Prodr.*, XVI, p. I, 30, published in 1869, *Roxb. Fl.*
Indica, p. 1, 30, published in 1832, *DC. Prodr.*, XVI, p. I, 30, published in 1869, *Roxb. Fl.*

Habitat—Cannabis indica has been reduced to C. sativa—the Indian plant being viewed as but an Asiatic condition of that species. This extends the region of the hemp-plant very considerably. It has been found

CANNA
indica.

Indian Shot

Palm walking
sticks.

319

Male bamboo

320

palm, and from the cocoa-nut palm, and are now-a-days largely used for umbrella handles. The "Malacca cane" is obtained from *Calamus Scirponum*, and the rattan from *C. Ratong* and one or two allied species, the former obtains its beautiful colour by being smoked

321

CANNA, Linn, Gen Pl, III, 654

Canna indica, Linn, Roxb, Fl Ind, Ed C B C, 1, SCITAMINEÆ

INDIAN SHOT

Vern.—*Sabba jaya*, HIND, *Kiwāra*, N-W P, *Sarba jaya* lat sarbo

Butsarana, SING

References—*Thwaites En Ceylon Pl* 320, *Dals & Gibs, Bom Fl Suppl*, 60, 634, *C Powell, Smith,*

Habitat—Several varieties are common all over India and Ceylon, chiefly in gardens, where they are grown as ornamental and flowering plants, they are in flower all the year

Dye—"The SEED is black, and round like a pea and yields a beautiful but evanescent purple dye" (*Dals & Gibs, Bomb Fl*)

Medicine—The ROOT is used as a diaphoretic and diuretic in fevers and dropsy (*Atkinson*), and also given as a demulcent (*Irvine*) It is considered acrid and stimulant (*Fleming*) When cattle have eaten

DYE

Seed.

322

MEDICINE

Root

323

Seed.

324

FOOD

Root

325

Starch

326

Allment or

arrow-root

327

DOMESTIC.

Leaves.

328

Seeds.

329

Necklaces

330

THE STARCH SUBSTITUTES

§ "In the West Indies arrow-root has been obtained from *C. glauca*, called '*Tous les mois*' (*O'Shaughnessy*)" (*Surgeon C F H Warden, Professor of Chem to C F H Warden,*

use

sew

necklaces and other ornaments of them. In the West Indies the leaves are used to thatch houses" (*Drury*) [See also under Beads, Vol I—Ed] "In Bangalore, the leaves are used by the natives in lieu of plates, to serve ragi pudding and other dishes" (*F Cameron, Esq*)

C. 330

CANNABIS
sativa

The History of the Indian Hemp

Hemp
Acclimatised
and
Cultivated
in
India.

doubtful of its being a native of Southern and Central Russia, but sus-

springing up spontaneously on the *churs* of the Subarnarekhá river and to be wild in the territory of the Mohurbhunge State on the frontier of Midnapur and also in Singbhum. It is cultivated more or less throughout India, either on account of the NARCOTIC derived from (a) the resin, *charas*, (b) the young tops and unfertilised female flowers—*ganja* (or *ganja*), (c) the older leaves and fruit-vessels—*bháng*, or on account of the fibre, HEMP, or the ripe seed from which an OIL is prepared. *Ganja* is de-

property is not developed until the fruits are mature, leaves at this stage, and sometimes the fruits also, afford *bháng*. With *Cannabis indica* differing in so marked a degree according to the climate, soil, and mode of cultivation, it was rightly concluded that its separation from the hemp plant of Europe could not be maintained. We have here, in fact one of the most notable illustrations of the effect of climate in changing the

The History of the Indian Hemp.

CANNABIS
sativa.

chemical processes which take place in the structure and physiological peculiarities of a plant. In most instances, a plant taken by man from one climatic condition to another, either dies quickly, or if it survives, it exists in a sickly condition. A few plants however, such as the potato,

FORMS OF
HEMP.

The plant for one or other of these purposes is now extensively cultivated throughout Persia, in India, from the level of the sea in Bengal to the inner Himālaya at an altitude of 10,000 feet, in China; in Arabia, and in Africa, from the extreme south to the north, and on the mountains as well as on the plains, in the north-eastern portions of America and on the table-land of Brazil. It is also to be met with in Northern Russia even as far as Archangel. In England it not unfrequently occurs as a weed, springing up most probably from rejected birdseed.

The modes of cultivation and the nature of the soil required, depend on the purpose for which the plant is cultivated. This subject will accordingly be discussed later on.

HISTORY OF HEMP.

THE NARCOTIC

Indian Literature.—“The earliest synonym appears to be *bhānga*, which occurs in the Atharva Veda—the last of the four scriptures of the Hindus. It is derived from a root which means ‘to break,’ and is supposed to imply the process of debarkation by which the fibres of the plant were separated from the stem. This would indicate that even at the remote period when the *Veda* in question was written, probably about 3,000 years ago, the use of hemp as a fibre-yielding plant was well known and the knowledge fully utilised. The *Veda*, however, reckons it, along with the *Soma*, as one of the five plants ‘which were liberators of sin,’ and this would imply that its narcotic property was also well known. The word is used in the masculine form with a short final vowel, and not, as in later literature, with a long one. Both the masculine and feminine

332

probability, the habit of speaking of the narcotic in the masculine form of the name, and of the fibre in the feminine. As a matter of fact, the nar-

CANNABIS
sativa

The History of the Indian Hemp

HISTORY

cotic yielding is the reverse to the popular belief the male or staminate
and female are

and Sanskrit writers were aware of the existence of male and female
flowers centuries before the sexes of plants were realised in Europe

Himalaya

The
Narcotic.
333

Classical Literature of Europe —The ancient SCYTHIANS seem to have
been acquainted with the narcotic properties of the plant as well as with
its fibre HERODOTUS tells us that they excited themselves by 'inhaling
its vapour' HOMER makes HEPHÆSTUS administer to TELEMACHUS, in the
house of MENELAUS, a potion prepared from nepenthes, which made him
forget his sorrows This plant had been given to her by a woman of
Egyptian Thebes,
much stress on this
among them since
ing a secret by which
secret is supposed
(*Johnston, Chemistry of Common Life, 337*)

Mythology
334

Mythological History of the Narcotic —"The notices of hemp in Arabic
and Persian works are much more numerous The oldest work in which
it is noticed is a treatise by Hassan, who states that in the year 638 A.H.,
Sheik Jafer Shirazi, a monk of the order of HAIDER, learned from his
master the history of the discovery of hemp Haider lived in rigid pri-

in wine or spirit seems to have been the favourite formula in which Sheik
Haider indulged himself (*Dymock, Mat Med, W Ind, 604*)

A curious story is told in the Hindu mythology about the origin of
this plant "It is said to have been produced in the shape of nectar

excited On the last day of the Durga Pooja, after the idols are thrown into
water, it is customary for the Hindus to see their friends and relatives and
embrace them After the ceremony is over it is incumbent on the owner
of the house to offer his visitors a cup of *bhāng* and sweetmeats for tiffin
(lunch) (*U C Dutt's Mat Med Hind, 236*)

C. 334

The History of the Hemp Fibre.

CANNABIS
sativa.

More Recent Historic Facts regarding the Narcotic.—The use of hemp (*dháng*) in India was particularly noticed by Garcia de Orta (1563), and the drug at time to in his *T. East In*. "It calling DeLacy drug in O Shaughnessy in 1829.

HISTORY.

established place in the Pharmacopœia' (*Fluck. & Hanb, Pharmacog, 547-48*).

HISTORY OF THE HEMP FIBRE.

The following extract may be here published as giving the most trustworthy facts which can be adduced regarding the history of the fibre "According to Herodotus (born 484 B C.), the Scythians used hemp, but in his time the Greeks were scarcely acquainted with it. Hiero II, King of Syracuse, bought the hemp used for the cordage of his vessels in Gaul, and Lucilius is the earliest Roman writer who speaks of the plant (100 B C.) Hebrew books do not mention hemp. It was not used in the fabrics which enveloped the mummies of ancient Egypt. Even at

The Fibre.

335

Canvas,

with *hasish* before performing certain ceremonies or perpetrating inhuman deeds. The word according to some would appear to have been originally

Assassin.

CANNABIS
sativa.

History of the Hemp Narcotic.

great havoc It seems probable that the English form of the word was adopted at the latter date, but that the more Arabic form was known in Europe for some time previous Hemp is alluded to in the "Arabian Nights" under its more ancient Arabic name, *beng*.

CULTIVATION

It has already been incidentally remarked that the cultivation of *Cannabis sativa* in India is naturally referable to two sections : (a) Cultivation with a view to preparing some of the forms of the narcotic, and (b) cultivation on account of the fibre. It has also been stated that the hemp plant has, to a large extent, changed its character under Indian or rather Asiatic cultivation It is very generally admitted, for example, that in the plains, while the narcotic principle is readily developed, the hemp fibre is but very imperfectly formed Let it, however, be distinctly understood that by hemp is here exclusively meant the fibre of *Cannabis sativa* This remark is all the more necessary when it is added that in the Government reports of the Trade and Navigation

" " " "

Expectations
regarding
Hemp Fibre.

a superior oil-seed, and the hemp plant a valued narcotic, but neither

enters into an account
object of proving that it
Panjáb, but he makes no
mention of the fact that the principal seats of hemp cultivation, as a commercial article, are in Eastern Bengal, the Central Provinces, and Bombay The *Encyclopædia Britannica* has also fallen into the same mistake, and, indeed, illustrations might be multiplied to show that undue prominence has been given to the fact that the plant is grown in Garhwal, the

* See a further page regarding Godavery District

The Cultivation of Hemp in India.

CANNABIS
sativa.CULTIVA-
TION

Panjab, and Kashmir, the more so since by most writers the true regions of Indian cultivation have been, to a large extent, overlooked.

his *Report on the Cultivation of and Trade in Gánjá in Bengal* (1877), has placed in the hands of the public a valuable treatise which deals both with the cultivation of the plant and the preparation of the narcotic. Dr Forbes Royle in 1855 issued his *Fibrous Plants of India*, a work

personal observations, supplemented by several less important publications, and Government reports, the following abstract regarding Indian hemp cultivation has been prepared.

(a) CULTIVATION FOR THE NARCOTIC.

For the
Narcotic.
337

Bengal Cultivation.—The method pursued in Eastern Bengal, according to Mr. Hem Chunder Kerner, is as follows:

over the field, and it is freely manured is ploughed into the soil, and the means of the cultivator will admit of The belief is that for hemp the land cannot be too often so treated. The thorough water, the rain into the Nuri ing of sandy May after need be used of Sept ready sary for TR on the up by

by fav and manured, the furrows ploughed, and all weeds removed. At this stage the plants begin to form their flowers, when the services of an expert, known

CANNABIS
sativa.

The Cultivation of Hemp in India.

CULTIVA-
TION.Fruits
injure
Ganja.

as the *gānjā*-doctor (*poddār* or *parakdār*) are called in This person
all the male or
lanis.
Kerr
njure
gānjā
action
scape
detection, the result being that a certain number of the female plants are
fecundated, fruits and seeds being produced. These are thrashed out as
far as possible in the manufacture of the drug, the quality of which may

For the Fibre.
338

(b) CULTIVATION FOR THE FIBRE HEMP.

Indian Methods.—Dr. Royle very appropriately remarks: "There is every reason for believing that the plant is of Eastern origin, while there is no sufficient reason for thinking that the climate of Europe is so peculiarly suited to the production of its fibre as to exclude those of its native climes, especially here attention is paid to those where the plant is grown on account of it where it is cultivated for latter requires exposure sowing, while the growth of the fibre is promoted by shade and moisture, which are procured by thick sowing" It has already been pointed out that the regions suited for *ganja* cultivation are perfectly distinct from those where it might be possible to develop an industry in the fibre. However much it may be regretted it seems impossible to combine the two industries, and it is an accepted fact that, unless utilisable as a paper stock, the immense amount of stems annually destroyed by the *gānjā* cultivators must continue to be so.

Godavery
Hemp.
339

At the same time Mr. Morris, in his account of the Godavery District, gives some interesting facts regarding the cultivation of hemp fibre. It is planted in November and cut by the end of March. It is grown in drills and never watered. Clay soils and those beyond the reach of inunda-

Rise a pull of land The bundles are buried in mud and left to rot for about a week when they are taken out and beaten in the water, and after all impurities are removed, the fibre is collected." The exports from the district are said to have been, in 1854-55, 4,269 cwt

Unless there be some mistake, *Sunn* hemp having been called "*Cannabis sativa*," for Mr. Morris gives that scientific name as well as the vernacular name *sannum* for the fibre he is describing, this information is of the greatest interest, as it would show, what the writer was not aware of until recently, that hemp fibre was actually produced on the plains of India

Cultivation of Hemp in India

CANNABIS
sativa

EARLY EXPERIMENTS IN HEMP CULTIVATION.—In 1802 the Government of India made various experiments on an extended scale to establish hemp fibre cultivation. European seed was imported, and farms and factories established but finally abandoned. Recourse was had to improving the cultivation of the Indian stock. The cultivation and manufacture was carried on at Rishra, Cussimpore, Maldah, Gorackpore, Mhow, Rohilkand, and Azimgarh, under the experienced supervision of European hemp dressers. The results were everywhere unsatisfactory and

CULTIVA-
TION
For the Fibre

THE POSSIBILITY OF MORE FAVOURABLE RESULTS.—In spite of disheartening results, it cannot be definitely stated that it is impossible that hemp fibre can be produced in India. The efforts alluded to were mainly

Possible
Prospects

printed as it expresses pretty clearly Dr. Royle's view.—This (hemp)

Dr. Royle alludes to successful experiments of hemp cultivation in the plains, especially at Chittagong. But in most cases—as was proved with the plant reared at Saharanpur, it is admitted that the plains crop is far

CANNABIS
sativa.

The Cultivation of Hemp in India.

CULTIVA-
TION.

as the *gánjá-doctor* (*poddár* or *parakdár*) are called in. This person

Fruits
injure
Ganja.

yielded by them is very inferior and scarcely saleable. The destruction of the *mádí* plants is, however, never so complete but that a few escape detection, the result being that a certain number of the female plants are fecundated, fruits and seeds being produced. These are thrashed out as far as possible in the manufacture of the drug, the quality of which may

For the Fibre.
338

(b) CULTIVATION FOR THE FIBRE HEMP.

Godavery
Hemp.
339

which are procured by thick sowing." It has already been pointed out that the regions suited for *ganja* cultivation are perfectly distinct from those where it might be possible to develop an industry in the fibre. However much it may be regretted it seems impossible to combine the two industries, and it is an accepted fact that, unless utilisable as a paper stock, the immense amount of stems annually destroyed by the *gánjá* cultivators must continue to be so.

At the same time Mr. Morris, in his account of the Godavery District, gives some interesting facts regarding the cultivation of hemp fibre. It is planted in November and cut by the end of March. It is grown in drills and never watered. Clay soils and those beyond the reach of inundation are those best suited. "About 2,200 bundles can be produced in one *putti* of land, each bundle yielding $1\frac{1}{2}$ viss of fibre, or a total of 3,300 viss or 412½ maunds, and is valued at one rupee a maund. The expenses of cultivation are estimated at Rs. 8, and those of the preparation of fibre at Rs. 100 a *putti* of land. The bundles are buried in mud and left to rot for about a week when they are taken out and beaten in the water, and after all impurities are removed, the fibre is collected." The exports from the district are said to have been, in 1854-55, 4,269 cwt.

Unless there be some mistake, *Sunn* hemp having been called "*Cannabis sativa*," for Mr. Morris gives that scientific name as well as the vernacular name *sannu* for the fibre he is describing, this information is of the greatest interest, as it would show, what the writer was not aware of until recently, that hemp fibre was actually produced on the plains of India.

Cultivation of Hemp in India.

CANNABIS
sativa

EARLY EXPERIMENTS IN HEMP CULTIVATION.—In 1802 the Govern-

CULTIVA-
TION
For the Fibre.

Mhow, Rohilkand, and Azimgarh, under the experienced supervision of European hemp-dressers. The results were every where unsatisfactory and the experiments abandoned.

er the rejected stems from
but the enquiry in this

Possible
Prospects

printed, as it expresses pretty clearly Dr Royle's view — 'This (hemp)

would also be softer and more pliable at the same time that it retained a great portion of its original strength, and probably in as large a quantity as is yielded by the *sunni* plant. Thus, an article might be produced which, judging from the Italian samples, might enter into competition with the Russian product, and at all events afford much more valuable cordage than the several *...*

Dr Royle alludes to successful experiments of hemp cultivation in the plains especially at Chittagong. But in most cases as was proved with the plant reared at Saharanpur, it is admitted that the plains crop is far

CANNABIS
sativa.

The Cultivation of Hemp in India.

CULTIVA-
TION
For the
Fibre

separate flowers and borne on separate plants The male plants (called

authors give accounts of the methods pursued in Europe in hemp culti-

Italian Hemp
340Male Fibre
341

sowing, each is uprooted singly, care being taken not to injure the stem.
"The fibre is separated either by retting or by breaking and scutching"
(*Spens' Encycl*)

*Properties and Uses of Cannabis sativa*ECONOMIC
PROPERTIES

From the STEMS, LEAVES or FLOWERS, and even the FRUITS a RESIN-
OUS EXTRACT, of a powerful narcotic character, may be prepared The
INNER BARK affords the valuable FIBRE HEMP The SEEDS are occa-

The Narcotic—Indian Hemp

CANNABIS
sativa.

sionally eaten, they are much valued for feeding birds. An OIL is expressed from them which is of some importance, but can scarcely be called commercial.

RESIN OR NARCOTIC.

There are primarily three forms of this substance, but under each there exist also local modifications, special preparations from these, and adulterations. The three forms are known as *C. indica*, *C. sativa*, and

BENGAL MANUFACTURE

(1st) *GANJA*—This is known in the trade as consisting mainly of two forms *Flat Ganja* and *Round Ganja*. Speaking of the manufacture of *ganja* in Bengal Mr Hem Chunder Kerr says—"In February and March, when *ganja* attains its maturity the cultivator proceeds to make arrangements for reaping the crop and preparing the drug. His first step is to present himself to the supervisor, show him the license under

GANJA
342

menacing operations.

Flat Ganja—The stems are cut with a sickle about 6 inches above

Flat
343

size. These are arranged on a mat in a circular form, with their points directed towards the centre and overlapping each other. The circle thus

firmly among the flowers in the desired form. Fresh twigs are then

mats are spread and the flowering twigs beaten two and two together so as to shake off the leaves or any fruits that may still remain and are re-arranged in a new circle, so that what was on the top before now forms the bottom

CANNABIS
sativa.

The Narcotic—Indian Hemp.

GANJA.

layer of the new circle. The treading is repeated stage by stage until the stack is again covered by the mats, and men take up their inexplicable seat on the top. After this each twig is trodden upon separately, being placed for

Round
344

thin sausage shape near the apex of the twig. This rolling is repeated

Char or rora
345

ganja Chi

Mr E. . . .
ganja of the
Garhwál is
exported
tracts. Tw

the *biluchar*. The *pattar* is imported chiefly from Holkar's territories and is of quality inferior to the Bengal *ganja*. It is purchased at from R5 to 6 a maund in Indur in the rough state, and "pays a duty of about 4 annas per maund on exportation to British territory." It is sold retail at from R3 to 4 a seer. The *biluchar* variety is imported from Lower Bengal, and is sold at R10 to 12 a seer.

BOMBAY AND THE CENTRAL PROVINCES.

IMITATIONS
OF GANJAExpressed
Juice
346
Decoction
347

Bengal is concerned, it may confidently be stated that adulteration can
C. 347

The Narcotic—Indian Hemp.

CANNABIS
sativa.

alone take place when the intoxicant reaches the hands of the dealer. In the *golas* it is quite pure.

The mention of *chur*, and of the extracts referred to by Dr. Irving,

which
int (see
accord-
to state

CHARAS.
348

ground. The crop is reaped about November and the powder stored in small 24^{lb} bags. About May these are sold to the traders, who cut the bags open and spread out the now partially agglutinated powder on cloths under the sun. It softens and deepens in colour and is hard pressed into bags or bales 1½ maunds in weight (a half pony-load ready for exportation). The quality is judged of by the amount of oil seen through the . . . until it is of t and exposing surface of th broken, is se pure steel. exposed, it linsed oil and a powder of the hemp leaves

From the above description it would appear as if Yarkand *charas* was

MONEA.
349Momea of
Nepal.
350

CANNABIS
sativa.

The Narcotic—Indian Hemp.

MOMEA

given internally in cases of wounds and ulcers along with *ghî*, dose one *masha*." It is noteworthy, in connection with Dr Gimlette's discovery regarding human fat used in the manufacture of Nepál *momea*, that amongst the ignorant classes of Northern India a superstition prevails that they may be captured and carried off to some distant land to be made into *momea*. This fact has been alluded to by various officers in

Mumtal
351

several localities where bituminous products occur, as they are commonly sold as drugs in the bazars of that country. According to Captain Hutton (*Cal. Jour., Nat. Hist., Vol. VI, 601*), a mineral pitch called

salts of lime. There was no trace of bitumen or sulphur. In fact, this

Momyal
352

353

354

exude from a crack on the face of a high rock

There are thus numerous allusions to a substance or substances known in the bazars of India under the name *momea*, but in none of the published accounts of this drug is there the slightest reference to its being a product of Indian hemp, although, in the early literature of that narcotic, it is repeatedly stated that a pure waxy form of *charas* obtained from Nepál is sold under the name of *momea*.

Charas from
Sind
355
Central India
356

Charas is collected in Sind and in Central India by causing men to run through the hemp fields. They are said to be generally clad in leathern aprons to which the resin adheres, but in some cases are reported to have their bodies first oiled and then to run naked through the fields

C. 356

The Narcotic—Indian Hemp

CANNABIS
sativa.

CHARAS.

Trans Hima-
laya

357

Garda or
Panjab
Charas

358

Surkhal,
Shangra, and
Khaki

359

stirring about the *bhāng* and making hay of it. Soon a fine dust flies out and, filling the room, settles down on the surface of the cloth spread over the heaps. When all the dust has been shaken out and settled on the cloth the cloth is rolled up and the dust is collected and pressed into balls.

(3rd) *BHĀNG* OR *SIDDHĪ*, *SABZI*, AND *SABZI*—Apparently the wild plant is the chief source of this form of the drug, which consists of the mature leaves and in some parts of India of the fruits also. The resin is

BHANG.

360

article is taken into consideration

INDIAN PREPARATIONS FROM HEMP

FORMS OF INDIAN HEMP—As already explained there are three forms of this poisonous drug (a) *ganja*, the agglutinated female flowering tops and resinous exudation on these, (b) *charas*, a resinous substance found on the leaves, young twigs, and bark, and (c) *bhāng* or *siddhī*, the mature leaves,

Smoking
mixtures.

361

Hashish.

362

Majun

363

PRICES.

... and a part growing within a certain radius of his hut, and it would be impossible to prohibit him gathering, from such a plant, the daily quantity used by himself and family. This is precisely the state

CANNABIS
sativa.

The Hemp Fibre of India.

of affairs which prevails over a great part of India, and, indeed, on the

Bedding for
Cattle.

plant, and the consumption can therefore be regulated by law. The Excise Act provides that licensed persons may cultivate the plant, prepare the narcotics, and retail these to the consumer. The right to vend is sold by public auction, a person purchasing thereby the sole right, for one year, to all or so many of the shops in a district. Any person, other than a licensed dealer, having in his possession more than a very small quantity at one time is liable to prosecution and fine. This system of farming the wholesale and retail shops exists all over India,—Madras

Excise
Arrange-
ments.

COLLS.

THE FIBRE-HEMP.

FIBRE
364

The reader is referred to the account given of the cultivation of the hemp plant in a preceding page. It will there be found that a con-

When
Mature.

Lignification.

nd modes of culture,
the plant in India,
rattle character com-
n due to the fact of
e fibre at an earlier
hus, for example, it
plains of India, at

Saharanpur, grew vigorously, attained a height of 12 feet, and gave every promise of proving successful. When reaped, Dr. Falconer, however, reported that "the hemp-fibre did not retain the strength or

Experiments
to be per-
formed in
India.

microscopically and chemically examined once a fortnight, right through

The Hemp Fibre of India.

CANNABIS
sativa.

their subsequent growth, or until in each locality the period when lignification was reached by the plants had been determined. It would also be

FIBRE.

failed to discover such regions or were imperfectly conducted, for, with the exception of certain limited tracts of the Himalayas, no part of the plains of India can be said to have been discovered in which there is the least hope of hemp or flax cultivation becoming of much importance. (See remarks as to hemp in Godavery District No. 339).

In portions of the North-West Himalaya the hemp plant has been cultivated for its fibre for a very long time. Mr. Atkinson gives a brief but practical account of this industry in his *Himalayan Districts*

abolition of the

Separation
of Fibre.

lessens the value of the fibre very much, since it increases the labour in cleaning it, each hank requiring to be opened out by the hand.

man
must
num
true

CANNABIS
sativa.

The Hemp Fibre of India.

FIBRE.

EUROPEAN
HEMP.
366

ropes and twine. Where this competition proved comparatively hopeless, substitutes were brought forward, and at the present day the most extensively used fibres in the rope trade may be said to be hemp, coir (or the fibre from the outer layer of the cocoanut), Manilla hemp, cotton, and sunn-hemp. Italy produces the finest hemp, France is perhaps next in importance, then Great Britain, Servia, Germany, and of Asiatic countries China is reputed to produce good hemp.

INDIAN FOREIGN TRADE IN "HEMP"

Raw Hemp
367Manufactures.
368Cordage
369

		Foreign Hemp imported	Foreign Hemp exported	Indian Hemp exported.
		R	R	R
Raw Hemp	Raw Hemp . . .	1881-82 . . . 1,10,875	.	5,59,112
		1882-83 . . . 1,82,993	.	4,30,325
		1883-84 . . . 1,76,765	.	6,85,316
		1884-85 . . . 2,14,118	.	5,82,679
		1885-86 . . . 1,96,052	.	9,88,825
Manufactures	Manufactured Hemp (excluding cordage)	1881-82 . . . 10,179	4,182	1,409
		1882-83 . . . 27,090	8,857	3,176
		1883-84 . . . 32,570	4,548	6,510
		1884-85 . . . 41,356	150	3,129
		1885-86 . . . 42,810	323	3,205
Cordage	Cordage and rope excluding jute, but otherwise the bulk probably Manilla Hemp and true Hemp	1881-82 . . . 3,22,485	24,886	3,25,178
		1882-83 . . . 4,31,693	15,586	2,84,166
		1883-84 . . . 3,90,584	11,198	4,92,068
		1884-85 . . . 3,52,413	13,076	3,53,389
		1885-86 . . . 3,24,519	7,437	3,28,320

Foreign Trade in Manufactured and Unmanufactured Hemp, excluding
Cordage

Year.	Imports	Exports and re-exports
	Value	Value
	R	R
1881-82	1,21,054	5,64,703
1882-83	2,10,083	4,42,358
1883-84	2,09,335	6,96,374
1884-85	2,55,474	5,85,053
1885-86	2,38,862	9,02,353

The Indian Hemp.

CANNABIS
sativa.

Detail of Imports, 1885-86

FIBRE
Imports.
370

Province into which imported	Value	Country whence imported	Value.
	R		R
Bengal	1,33,235	United Kingdom	83,431
Bombay	1,01,600	China	1,23,474
Madras	1,183	Philippines	2,609
Sind	2,844	Straits Settlements	17,827
		Other Countries,	11,521
TOTAL	2,38,862	TOTAL	2,38,862

Detail of Exports, 1885-86

Exports.
371

Province from which exported	Value	Country to which exported	Value
	R		R
Bengal	3,11,551	United Kingdom	6,78,607
Bombay	6,31,444	Belgium	2,56,566
Madras	40,358	Persia	11,438
		Arabia	15,698
		Other Countries	30,044
TOTAL	9,92,353	TOTAL	9,92,353

It has been found impossible to give the quantities, since the raw fibre is expressed in weight, cloth in pieces, and rope in balls of various lengths and weights.

OIL.

HEMP SEED
OIL.
372

gravity of 0.9252 at 15°C, it thickens at -15°C, and solidifies at -25° to -27.7°C. It dissolves in boiling hot water and in 30 parts of cold alcohol.

MEDICINE.

MEDICINE.
373

CANNABIS
sativa.

The Indian Hemp as a Drug.

MEDICINE

Chur or Round
Ganja best
suited for
Pharmacy
Flat Ganja
and Charas
should be
avoided

to be a small dose of oil of late years as to a feeling of agree-
ment
to purchase
en allowing
a dealer or
er a permit
raised as to
stered as of
From what
to use for

o
d
a
E
C

ducing uterine contractions.
It is admitted by most Indian physicians to be of special merit in the treatment of tetanus and cholera and has not the injurious after effects which but too frequently result from
ever, very similar to that of opium,
habitual opium eater may take large
quantities of hemp without injurious consequences
Sir William O'Shaughnessy was the first European writer to draw prominent attention to the peculiar properties and actions of the hemp-narcotics. He experimented with these in Calcutta and published his results. The reader is referred to his *Bengal Dispensatory* and to a "Memoir on the preparations of Indian Hemp" in the *Transactions of Medical and Physical Society of Calcutta* for 1839, and to two papers in the *Journal of the Asiatic Society*, Vol. VIII, of the same year. Shortly after the appearance of these most exhaustive accounts, the drug began to be experimented with in Europe.

The Indian Hemp as a Drug.

CANNABIS
Sativa.Ainslie, in his *Materia Indica*, 2nd Vol., gives an interesting account of MEDICINE.

ferent ingredients, of which datura and opium are frequent. In some parts of India a beer is brewed with *blāng*, and this, together with *blāng* itself, *mājām* and other preparations, are often employed in Native pharmacy.

and convenience, Indian Hemp is the next anodyne hypnotic and antispasmodic to opium and its derivatives, and often equal to it." Dr.

Makhsan, "the leaves make a good snuff for detarging the brain; their juice applied to the head ear it allays
rhœa and go

Mat. Med. West India).

The medicinal properties of hemp, in various forms, are the subject of some interesting notes by Mirza Abdul Russac. "It produces a ravenous

Leaves,
376

C. 376

CANNABIS
sativa.

The Indian Hemp as a Drug.

MEDICINE

tice has greatly decreased of late years owing to a feeling of insecurity as to the quality of the article. It is commonly recorded that no reliance can be placed on the strength of the material. The value of a measure of the

Chur or Round
Ganja best
suited for
Pharmacy
Flat Ganja
and Charas
should be
avoided

in cholera, menorrhagia and uterine hæmorrhage, rheumatism, hay fever, asthma, cardiac functional derangement, and skin diseases attended with much pain, and pruritus. In lingering and protracted labours depending upon atony of the uterus, it has been employed with the view of inducing uterine contractions.

It is admitted by most Indian physicians to be of special merit in the treatment of tetanus and cholera and has not the injurious after effects frequently result from nalar to that of opium, m eater may take large

The Indian Hemp as a Drug.

CANNABIS
sativa.

Ainslie, in his *Materia Indica*, 2nd Vol., gives an interesting account of

MEDICINE.

itself, *majun* and other preparations, are often employed in Native pharmacy.

... remarks, derived from some years' experience, from Calcutta, that it is a good remedy for pain, obtain sleep, and put an end to spasm in circumstances under which morphia either did not suit or was objected to by the patient, and after wide experience with it I am quite satisfied that it is an excellent substitute for it, if given in sufficient doses. The difficulty is, to be always sure of the quality of the extract, or rather of the *ganja* from which the extract is obtained.

Uniformity in
quality.

and convenience, Indian Hemp is the next anodyne hypnotic and antispasmodic to opium and its derivatives, and often equal to it. Dr.

applied externally in
nutation, a poultice
erysipelas, neuralgia
istered internally."

Mat Med West India)

The medicinal properties of hemp, in various forms, are the subject of some interesting notes by Mirza Asad Khan, "the subject of the
appetite and constipation, arrests

smokers of *ganja* generally die of diseases of the lungs, dropsy, and anasarca, so do the eaters of *majun* and smokers of *siddhi*, but at a later period. The inexperienced, on first taking it, are often senseless for a day, some go mad, others have been known to die.

Dr U O Dutt says that, according to the Sanskrit writers, "the leaves of *Cannabis sativa* are said to be purified by being boiled in milk

Leaves.
376

CANNABIS
sativa.

The Indian Hemp as a Drug

MEDICINE.

Dysentery.

Affections of
the eye
Piles.NASHA
377Oil used in
Rheumatism

Acute Mania

Hysteria.

Orchitis

Asthma
Chronic Colic

Charas of the trade, but it is terribly adulterated. The plant is called The OIL extracted from the remedy, applied by rubbing (Aitchison, Simla) "Used in useful in atonic dyspepsia and di

in dysuria, and in relieving pain in dysmenorrhœa" (Dr E G Russell, Superintendent, Asylums, at Presidency General Hospital, Calcutta). "Commonly used as a narcotic, a few grains of the leaves called *siddhi* rubbed in with cardamom and other spices to allay pain, taken as a drink

mixed with other drugs and spices, forms an useful compound in diarrhœa and indigestion of children" (Assistant Surgeon Shri Chunder Bhattacharya, Chanda, Central Provinces) "The leaves, which are known as

The Indian Hemp as a Drug

CANNABIS
sativa.

(Dr G Price Civil Surgeon, Shalabad) — It is also used in the form of tincture for the form of etc

MEDICINE
Ague Fits
Impotence

from a medical point of view, are the *Resin* and *Volatile Oil*

"The former was first obtained in a state of comparative purity by T and H Smith in 1846. It is a brown amorphous solid, burning with a bright white flame and leaving no ash. It has a very potent action

CHEMICAL
COMPOSITION

small crystals. With due precautions it may be separated into two bodies the one of which named by Personne *Cannabene* is liquid and colourless, with the formula $C_{15}H_{20}$, the other which is called *Hydride of Cannabene*, is a solid separating from alcohol in platy crystals to which Personne assigns formula $C_{17}H_{22}$. He asserts that *Cinnabene* has

Cannabene
378

from the oil which he obtained from the fresh herb, just after flowering, to the extent of 0.3 per cent.

"It remains to be proved whether an *alkaloid* is present in hemp, as suggested by Preobraschensky.

The other constituents of hemp are those commonly occurring in other plants. The leaves yield nearly 20 per cent of ash.

As to the resin of Indian hemp Bolas and Francis, in treating with

from purified resin of charas, but without success" (Fluck and Hanb, *Pharmacog* page 549)

Dr Dymock (in his 2nd Ed of the *Materia Medica of Western India*) goes into considerable detail on the chemistry of this drug. Preobraschensky discovered in China *haschisch*, a volatile alkaloid which he believed to be identical with nicotine. Dragendroff and Marquiss

published his conviction that hemp contained several alkaloids the principal one being a substance he named *Tetano-cannabine*. More recently to all these published results of the chemical investigation of the narcotic resin

CANOES.

The Indian Hemp Canoes.

oil contained phenol, ammonia, and several other of the usual products of destructive distillation.

"The nicotine like principle contained in this oil appeared to be an alkaloid. It formed salts which evolved a strong nicotine-like odour when acted on by alkalis. But physiologically it was found to be inert, and therefore was evidently not identical with nicotine" (*Ind. Med. Gas*, Dec 1884)

FOOD.

FOOD.
379

Food — Messrs Duthie and Fuller, writing about the Himálayan tracts within the North-Western Provinces, say that the seed is not uncommonly roasted and eaten by the hill-men, and that after the oil is expressed the oil-cake is given to their cattle. Dr Stewart writes that on the Sutlej the seeds are roasted and eaten in small quantities with wheat.

DOMESTIC AND INDUSTRIAL USES.

DOMESTIC.
380

Cannabic Composition — "This material for architectural decoration is described by Mr B. Albans to have a basis of *hemp* amalgamated with sheets of large pness of detail than half the elastic to be adapted to wall surfaces, bearing blows of the hammer and resisting all

of varnish, the material is so hard as to allow gold to be furnished after gilding the ornaments made of it" (*Ure, I*, 611).

CANOES

See Boats, Vol II, B 548

381

TIMBERS USED FOR CANOES, DUG-OUTS, TROUGHS, WATER PIPES, DRINKING CUPS, &c.

- 1 *Acer caesium*, *Will* (drinking cups made in Tibet)
- 2 *A. oblongum*, *Wall* (drinking cups)
- 3 *A. pictum*, *Thunb* (drinking cups made of knotty excrescences).
- 4 cups).
- 5
- 6
- 7
- 8 *Artocarpus Chaplasha*, *Roxb* (much used for canoes).
- 9 *A. Lakoocha*, *Roxb* (canoes)
- 10 *A. nobilis*, *Thw* (Ceylon canoes)
- 11 *Bœhmertia rugulosa*, *Wedd* (Lepchas make cups, bowls, and tobacco-boxes)

C. 381

Woods used for Canoes, Dug-outs, &c.

CANSCORA
decussata.

- 12 .
13 .
14 .
15 .
16 .
17 .
18 .
19 .
20 .
21 .
22 .
23 . boats
and canoes)
24 *Dnabanga sonneratioides*, Buch. (canoes, cattle-troughs cut out of green wood)
25 *Dysoxylum Hamiltonii*, Hiern (canoes)
26 *D. procerum* Hiern (Assam canoes)
27 *Givotia rottleriformis*, Griff (catamarans).
28 *Gmelina arborea*, Roxb (clogs, canoes, &c.).
29 *Gyrocarpus Jacquini*, Roxb. (preferred above all other woods for catamarans)
30 *Hopea odorata* Roxb (Burma canoes)
31 *Juniperus excelsa*, M. Bieb (drinking cups).
32 *Lagerstræmia Flos-Reginæ*, Retz (boats and canoes).
33 .
34 .
35 .
36 .
37 .
38 .
39 .
40 .
41 .
42 .
43 .
44 *Populus ciliata*, Wall (water troughs)
45 *Sarcosperma arborea*, Hook (Sikkim canoes).
46 *Schuma Wallichii*, Choisy (Assam canoes)
47 *Shorea obtusa*, Wall (canoes).
48 *S. robusta*, Gartin (Hills of Northern Bengal, canoes)
49 .
50 .
51 .
52 .

CANSCORA, Lam, Gen Pl., II, 811.

Canscora decussata, R. & Seb., Fl. Br. Ind., IV., 104; Bot. Mag., t. 3066, GENTIANACEÆ

Syn *PLADERA DECUSSATA*, Roxb., Fl. Ind., Ed. C. B. C., 135

Vern — *Sankhdhuli*, HIND; *Dankuni*, BENG; *Shun khapushopp*, CUTCH; *Sankhopushpi dandolpala*, SANS

References — *Thwaites En Ceylon Pl.*, 204, Voigt, *Hort. Sub. Cal.*, 510; U. C. Dutt, *Mat. Med. Hind.*, 201, 296, 316, Dymock, *Mat. Med. W. Ind.*, 451; also 2nd Ed., 542

**CANTHIUM
didymum.**

Cantharides; Canthium.

MEDICINE
383

Habitat—Common throughout India from the Himālaya to Burma, ascending to 4,000 feet, is abundant in the plains of Bengal and not uncommon in Ceylon

Med Hind, 201).
Special Opinions.—§ "This deserves a trial" (*Surgeon-Major C F. McKenna*). "Laxative, tonic, expectorant" (*Dr. W Barren, Bhuj, Cutch*).
and
itta,
ade
be
'at.

384

Canscora diffusa, *Br*, *Fl Br Ind*, IV, 103; *Wight*, *lc*, t 1327 (*not* [of *Clarke*])

Syn.—PLADERA VIRGATA, *Roxb*, *Fl Ind*, Ed C B C, 134

Vern—*Kyauk pan*, BURM

References—*Thwaites*, *En Ceylon Pl*, 204, *Dals and Gids*, *Bomb Fl*, 158, *Voigt*, *Hort Sub Cal*, 520

Habitat—Common throughout India, ascending to 4,000 feet, from Kumāon and Bhutan to Ceylon and Penasserim

Medicine.—Used as a substitute for *C. decussata*

MEDICINE.

385

386

C. sessiliflora, *Roem and Sch*, *Fl Br Ind*, IV., 104

387

CANTHARIS, Latreille

Cantharis vesicatoria, *Latreille*, COLEOPTERA.

CANTHARIDES, BLISTERING BEETLE, SPANISH FLIES, *Eng*,
MOUCHES D'ESPAGNE, *Fr*, SPANISCHE FLIEGEN, *Germ*,
CANTERELLE, *It*, HISCHPANSKIE MUCHI, *Rus*, CANTHARIDES, *Sp*

References—*Pharm Ind*, 274; *U S Dispens*, 15th Ed, 342, *Spons*,
Encyclop, 796, *Balfour*, *Cyclop*, *Ure & Dic of Arts and Manufactures*

Habitat.—A dried insect imported into India and sold by chemists
For indigenous insects used as substitutes, see *Mylabris cichorii*, *Fabr.*

Blistering
insect
388

389

CANTHIUM, Lam, *Fl Br Ind*, III, 131.

The *Genera Plantarum* reduces the above genus to *PICTETRONIA*, *Linn*; but *CANTHIUM* has been retained in the *Flora of British India*, which puts *PICTETRONIA* (in part) under *CANTHIUM*

390

Canthium didymum, *Roxb.*, *Fl. Br Ind*, III, 132; RUBIACEÆ

Vern—*Garbha gojha*, *SANTAL*, *Yerkoli*, *TAM*, *Yellal*, *porawa mara*,
Gal koranda, *SING*, *KAN*

References—*Roxb*, *Fl Ind*, Ed C B C, 180; *Kurz*, *Fl Burm*, II,
359, *Thwaites*, *En Ceyl Pl*, 152, *Bom Gas*, XV, 65

Habitat—A shrub or small tree found in the Sikkim Himālaya at an altitude of 1,500 feet and distributed east to the Khasia and Jyntea mountains. It also is met with in Chutia Nagpur and in the Western Peninsula from the Concan southwards to the Malayan Peninsula and Ceylon

C. 390

Canthium • Canvas

CANVAS.

Medicine — Bark used by the Santals in fever (*Rev. A. Campbell*)

Structure of the Wood — Hard heavy, and close-grained, yellowish, with central masses of black (*Pomb Gaz*) This is very much like the description of the wood, as given by Brandis and by Lisboa for *C. umbellatum*.

MEDICINE.

391
TIMBER.
392

Canthium parviflorum, *Lamk, Fl Br Ind, III, 136*

393

Syn — *WEDERA TETRANDBA*, Hill 1, KANLEN KARA in *Rhe de, Hort Mal*, 1, t 36

Vern — *Arni*, BOMB, *Karai-choddi*, TAM, *Tijéron kard*, MAL, *Dalusu, chettu, lalsu*, TEL (*AINSLIE*), *Kára*, SING

References — *P. L. IND. FL. COR. — C. L. H. T. I.*

Habitat. — A shrubby plant met with at altitudes of 4,000 feet, in the

MEDICINE.

394

FOOD

395

TIMBER.

396

397

C. umbellatum, *Wight, Ic, t 1034; Fl Br Ind, III, 132.*

Syn — *PLECTONIA DIDYMA*, *Benth & Hook*; *Brandis, For Fl*

Vern — *Arsál*, BOMB, *Neckanre, nalla, balsá*, TAM & TEL; *Abalu, KAN, Tolan, URIVA*

References — *Brandis, For Fl, 276* *Bedd, Flor Sylv, 221; Dals & Gibs, Bomb Fl, 113. Gamble, Man Timb, 230 (under Plectonia didyma, Benth & Hook); Lisboa, U Pl, Bomb, 67.*

Habitat — An evergreen tree met with in the Western Peninsula (on the Ghats at altitudes of 4,000 to 8,000 feet) and distributed south to Tenasserim and Ava

Structure of the Wood — Hard close-grained

white or chocolate-centre (*Brandis*)

small, numerous

and numerous

black wood (*Coy*)

TIMBER.

398

Timber is used for agricultural purposes

CANVAS.

Canvas.

399

SAILCLOTH, *Eng*, KANEVAS and SEGELTUCH, *Germ.*, CANEVAS and TOILE-A-VOILE, *Fr*, ZEHDOK, *Dut*; LONA, *It*, *Port*, *Sp*, CANEVÁZZA, *It*, *Port*; PARUSSINA, PARUSSNOE POLOTNO, *Rus*, KITTAN, *Tam.*, *Tel*

A coarse strong cloth

is employed by artists for painting on

CANTHIUM
didymum.

Cantharides; Canthium.

MEDICINE
383

Habitat —Common throughout India from the Himālaya to Burma, ascending to 4,000 feet, is abundant in the plains of Bengal and not uncommon in Ceylon

Med Hind 201

Special Opinions —§ This deserves a trial" (*Surgeon Major C F. McKenna*). "Laxative tonic, expectorant" (*Dr. W Barren, Bhuj, Cutch*).

384 *Canscora diffusa*, *Br, Fl Br Ind, IV, 103, Wight, Ic, 1 1327 (not of Clarke)*

Syn —*PLADERA VIRGATA*, *Roxb, Fl Ind, Ed C B C, 134*

Vern —*Ayauk pan*, BURN

References —*Thwaites, En Ceylon Pl, 204, Dals and Gibs, Bomb Fl, 158, Voigt Hort Sub Cal, 520*

Habitat —Common throughout India, ascending to 4,000 feet, from Kumāon and Bhutan to Ceylon and Tenasserim

Medicine —Used as a substitute for *C decussata*

MEDICINE
385
386

C. sessiliflora, *Roem and Sch, Fl Br Ind, IV., 104*

387

CANTHARIS, Latreille

Cantharis vesicatoria, Latreille, COLEOPTERA

CANTHARIDES BLISTERING BEETLE SPANISH FLIES, *Eng*,
MOUCHES DESPAGNE *Fr*, SPANISCHE FLIEGEN, *Germ*,
CANTERELLE, *It*; HISCHPANSKIE MUCHI, *Rus*, CAN-
THARIDES, *Sp*

Blistering
Insect.
388

References —*Pharm Ind, 274 U S Dispens, 15th Ed, 342, Spens, Encyclop 796 Balfour, Cyclop, Ure's Dic of Arts and Manufactures*

Habitat —A dried insect imported into India and sold by chemists
For indigenous insects used as substitutes see *Mylabris cichoru, Fabr.*

389

CANTHIUM, Lam, Fl Br Ind, III, 131.

The *Genera Plantarum* reduces the above genus to *PLECTRONIA Linn*, but CANTHIUM has been retained in the *Flora of British India*, which puts *PLECTRONIA* (in part) under CANTHIUM

390

Canthium didymum, Roxb; Fl. Br Ind, III, 132, RUBIACEÆ

Vern —*Garbha goyha, SANTAL, Yerkolı TAM, Yellal, porawa mard, Gal karanda, SING, KAN*

References —*Roxb, Fl Ind, Ed C B C, 180, Kurz Fl Burm, II, 359 Thwaites, En Ceyl Pl, 152, Bom Gaz, XV, 65*

Habitat —A shrub or small tree found in the Sikkim Himālaya at an altitude of 1500 feet and distributed east to the Khasia and Jyntea mountains. It also is met with in Chutia Nagpur and in the Western Peninsula from the Concan southwards to the Malayan Peninsula and Ceylon

C. 390

Canthium: Canvas.

CANVAS.

Medicine — Bark used by the Santals in fever (Rev A. Campbell)

Structure of the Wood — Hard heavy, and close-grained, yellowish, with central masses of black (Jomb Gas) This is very much like the description of the wood, as given by Brandis and by Lisboa for *C. umbellatum*

MEDICINE.

391
TIMBER.
392*Canthium parviflorum*, Lamk., *Fl Br Ind.*, III, 136

Syn — *WEPERA TETRANDRA*, Hilld., *KANDEN KARA* in Rhe. de, Hort Mal., t 36

Vern — *AIRNI*, BOMB., *Kavar-cheddi*, TAM., *Tajeron kard*, MAL., *Balusu, chettu, balsu*, TEL. (KINSLIF), *Kara*, SING.

References — *Forb.*, *Fl Ind.*, *E I C B C.*, 179, *Gamble*, *Man Timb.*, 230, *Amst. & Mat. Med.*, II, 63, *Dymock*, *Mat. Med.*, *W. Ind.*, 713, and 2nd Ed., 409, *Lisboa*, *U Pl.*, BOMB., 162, *Thwaites*, *En Cey Pl.*, 152, *Trinens Cat. Ceyl Pl.*, 44

Habitat. — A shrubby plant met with at altitudes of 4,000 feet, in the

393

MEDICINE.

394

FOOD

395

TIMBER.

396

397

C. umbellatum, Wight, *It.*, t 1034, *Fl Br Ind.*, III, 132

Syn — *PLECTONIA DIDYMA*, Benth & Hook., *Brandis*, *For Fl*

Vern — *ARUL*, BOMB.; *Neckanie, nalla, balsu*, TAM. & TEL., *Abalu, kan*, *Tolan URIYA*

References — *Brandis*, *For Fl*, 276, *Bedd.*, *Flor Sylv.*, 221, *Dals & Gibs*, *Bomb Fl.*, 113, *Gamble*, *Man Timb.*, 230 (under *Plectonia didyma*, *Benth & Hook.*), *Lisboa*, *U Pl.*, BOMB., 87

Habitat. — An evergreen tree met with in the Western Peninsula (on the Ghats at altitudes of 4,000 to 8,000 feet) and distributed south to Tenasserim and Ava

Structure of the Wood — Hard, close-grained, and heavy, yellowish

TIMBER.

398

It is numerous. Gamble makes no mention of the irregular masses of black wood (Compare with *C. didymum*) Weight 57lb a cubic foot.

Timber is used for agricultural purposes

CANVAS.

Canvas.

399

SAILCLOTH *Jng*, *KANEVAS* and *SEGELTUCH*, *Germ.*, *CANEVAS* and *TOILE A VOILE*, *Fr.*, *ZFIDOCK*, *Dut.*; *LOVA*, *It.*, *Port.*, *Sp.*, *CANEVAZZA*, *It.*, *Port.*; *PARUSSINA*, *PARUSSNOE POLOITNO*, *Rus.*, *KITTAN*, *Tam.*, *Tel.*

CAPPARIS
sepiaria

The Wild Caper berries

References — D J E E F K E K D M I A C m

277 Balfour Cyclop

ig in most parts of India

MEDICINE
Leaves

- 417
- Bark
- 418
- Fruit
- 419

FOOD

- 420

FODDER

- 421

TIMBER

- 422
- 423

on the Economic Products of Chutia Nagpur)

Special Opinion — § A decoction of the leaves is used in syphilis ”
(Surgeon Major D R Thompson 1st District Madras)

Food — In the Southern Panjab and Snd the fruit is made into pickle
(Stewart) The twigs shoots and leaves are greed ly eaten by goats and
elephants

Structure of the Wood — Yellowish white, moderately hard, weight
about 47lb per cubic foot Used as fuel

Capparis multiflora, Hook f & Th ; Fl Br Ind, I, 178

Vern — Suntri NEPAL

References — Kurs For Fl Burm I 61 Gamble Man Timb, 11

Habitat — A climbing thorny shrub of the Eastern Himalaya and
Upper Burma

Structure of the Wood — White moderately hard

TIMBER

- 424
- 425

C olacifolia, Hook f & Th , Fl Br Ind I, 178

Vern — Naski has NEPAL Jhenok LEPCHA

References — Gamble, Man Timb 15 11

Habitat — A thorny shrub of the Sub H malayan tract from Nepal to
Assam chiefly in the undergrowth of Sissu forests along river banks

Structure of the Wood — White, hard, weight about 44lb per cub ft

TIMBER

- 426
- 427

C sepiaria, Linn ; Fl Br Ind I 177

Vern — Hián garna hius Pb Kantá gur kámas káliakara BENG
Kants kapali URIYA Kanthár, GUJ Nella uppi TEL Ah nsra kakh
dan SANS

References — Roxb Fl Ind Fd C B C 425 Brandis For Fl 15
Kurs For Fl Burm I 66 Gamble Man Timb 11 Thwaites
Enum Ceylon Pl 16 Dalz & Gbs Bomb Pl 10 Atchison Cat
Pb Pl 10 ; Vogt Hort Sub Cal 75 Murray Drugs and Pl Sind
54 Royle Ill Him Bot I 72 Balfour Cyclop

MEDICINE
428

cau
ton

TIMBER

- 429

DOMESTIC

- 430

Sera nore)

Structure of the Wood — White hard, pores moderate sized

Domestic Uses — The branches make excellent hedges

C. 430

The True Caper-berry.

CAPPARIS
spinosa

431

Capparis spinosa, Linn, Fl Br Ind, I, 173

THE EDIBLE CAPER

Syn.—C MURRAYANA, Graham; Wight, Ic, t 379

Vern.—*Aabra ber*, HIND, *Aebra*, LADAK, TIBET, *Ulla kanta*, KUMAON, *Aaur*, *kiari bauri*, *ber*, *bandar*, *bassar*, *kakri*, *kander*, *taker*, *barar*, *keri*, *kabra kabarra*, *karari bauri*, P^W, *Aufari*, SIND, *habar*, BOMB *habarra kabarra* AFG *habar kabur* ARAB, *Acbir*, PERS (In Persia it is known as *habar*, *karak*) *habur*, SYRIAN, *habarish*, TURKISH

Habitat.—This is the plant which affords the Caper berry of Europe. It occurs in India in the central and northern parts of the Panjab and in Sind, is less frequent in Rajputana than *C. aphylla*.

Medicine.—Dr Stewart remarks that in Kangra the roots are said to be applied to sores. The author of the *Makhzan-ul-Adwiyā* considers the root bark "to be hot and dry and to act as a detergent and astringent,"

MEDICINE.
Roots432
Root-bark.433
Juice.

434

Buds
435

considered diuretic, and was formerly employed in obstructions of the liver and spleen, amenorrhœa, and chronic rheumatism."

Chemical Composition.—"The root-bark is said to contain a neutral bitter principle of sharp irritating taste, and resembling senegin. The flower-buds, distilled with water, yield a distillate having an alliaceous odour. After they have been washed with cold water, hot water extracts from them Capric acid ($C_{12}H_{24}O_2$), and a gelatinous substance of the Pectin group, Capric acid is sometimes found deposited on the calices of the buds in white specks having the appearance of wax (*Rochleder and Blas*)" (*Watts' Dict., Chemistry*)

CHEMISTRY.
436

Food.—In Europe this furnishes the Caper. Mr Edgeworth found the buds (prepared in the style of "Capers") to answer very well as a substitute for the European congener. In India the ripe fruit is either eaten raw or made into pickle. In Sind and in some parts of the Panjab, a compound of oil, mustard, fœnu greek, &c., is used in pickling capers. In Ladak the leaves are eaten as greens.

FOOD
437
Berries.
Pickle.
438
Leaves.
439
FODDER.
440

Fodder.—The leaves and ripe fruits constitute a favourite food of goats and sheep.

CAPSICUM
annuum

Capsicum or Red Pepper

441

Capparis zeylanica, Linn Fl Br Ind I 174Syn — *C ACUMINATA* Roxb *C BREVISPIA* DCVern — *Kalo-kera* BENG *Authoondy kai* TAMReferences — *Voigt Hort Sub Cal*, 74 *Dals & C Balfour Cyclop*

Habitat — Common in the Carnatic and Malabar, &c Western Dekkan and in the drier parts of Ceylon

Food — The green fruit is pickled

FOOD
Pickled
442CAPSELLA, *Manch*, Gen Pl, I 86

443

Capsella Bursa pastoris, *Manch*, Fl Br Ind I 159

SHEPHERD'S PURSE PICKPOCKET, Eng BOURSE DE T

Fr HIRTENASCHE Germ

Habitat — A weed in the vicinity of cultivation throughout the regions of India, particularly abundant on the N W H

Medicine — This very common weed is bitter and pungent volatile oil on distillation identical with oil of mustard and has b

MEDICINE

444

Oil

445

FOOD

446

447

CAPSICUM, Linn Gen Pl, II, 892

The greatest confusion exists in Indian literature as to the cultivated species of Capsicum. Popularly the larger fruits are usually designated Capsicum and the smaller Chilies. According to Firminger the powdered seeds of the latter constitutes Cayenne pepper. That author in his *Manual of Gardening for India* states that there are a great many varieties of Capsicum grown in

believed to be the same species as the

448

Capsicum annum, Linn DC Prodr XIII Pl 1 412 SOLANACE

RED PEPPER

Vern — *Mattisa wāngru lāl mirch march mirch gachn rech* HIND

UNM

References — *Roxb Fl Ind Ed C B C* 193 *Stewart Pb Pl* 156DC Orig of Cult Pl 289 *Voigt Hort Sub Cal* 510 *Plarm Ind*

909

428

can
ton

Ser

St

Dol

C. 430⁸

TIMBER

429

DOMESTIC

430

Capsicum or Red Pepper.

CAPSICUM
annuum.

Habitat.—A native of equatorial America, most probably of Brazil. Commonly cultivated for its fruit throughout the plains of India, and on the lower slopes of hills in Kashmir, and in the Chenab valley up to altitude 6,000 feet. When grown on the hills it is said to be very pungent. There are seven varieties, differing chiefly in the length, shape, and colour of the fruit, some being round, others oblong, of tusc, pointed or bifid, some smooth, and red, white, yellow, or variegated. It is probable that most Indian authors have confused this species with *C. minimum*, which see.

History.—This species has a number of different names in European languages, which all indicate a foreign origin, and the resemblance of the taste to that of pepper. In French it is often called *Pivre de Guinée* (Guinea pepper) but also *poivre du Brésil d'Inde* (Indian, Brazilian pepper) &c., denotation to which no importance can be attributed. Its cultivation was introduced into Europe in the sixteenth century. It was one of the peppers that Piso and Maxgraf saw grown in Brazil under the name *guia or guya*. They say nothing as to its origin. (*DC Orig of Cult Pl*). "Chilies are not mentioned by any Sanskrit writer, consequently their introduction into India must have taken place at a comparatively recent date. It is probable that the Portuguese brought the fruit from the West Indies. Up to the present time the cultivation of the plant is carried on more extensively at Goa than at any other place on the western coast and capsicums are well known in Bombay by the name of *Goza mirchi* (Goa pepper)" (*Dr Dymock, Mat Med W India*). Hove alludes to Capsicum as grown in Bombay in 1787 and expresses no astonishment at its existence in India.

CULTIVATION OF CAPSICUM.—A light well-manured soil is the best for all kinds in which the plants should be picked out at about four inches apart when they attain a growth of three inches, and afterwards put out into a bed of rich light earth when they attain six inches in height, giving them a good supply of water and keeping them clear from weeds. (*The Gardener*)

Medicine.—Dr Stewart says that the fruit is used externally in the form of plasters and taken internally in cholera, it is eaten from a conviction that it counteracts the effects of bad climates.

As a drug red pepper is considered by the natives as stomachic and stimulant, and is used externally as a rubefacient (*Dymock*). It has been employed with success as a topical application to elongated uvula and relaxation of the pendulous veil of the palate. Made into a lozenge with sugar and tragacanth, it is a favourite remedy for hoarseness with professional singers and public speakers. In putrid sore-throat whether symptomatic or

very usefully employed in febrile affections, bitters, tonics and other stimulants in weak states of the stomach, in cold leucophlegmatic habit dyspepsia and flatulence and as a gargle in relaxed states of the throat it is highly extolled and has also been used with success in the advanced stages of rheumatism. In native practice it is given in conjunction with asafoetida and sweet flag root, in cholera. By German physicians it is supposed to be particularly injurious in gonorrhoea" (*Murray's Pl and Drugs of India*).

Dr Sakhambari Arjun says that the fruit is used as a stimulant in snake bite.

Chemical Composition.—"Buchholz in 1816, and about the same time Braconnot, traced the acridity of capsicum to a substance called *capsicin*.

449

450

MEDICINE
Plaster.
451Lozenge.
452

CAPSICUM
annuum

Capsicum or Red Pepper

- 441 *Capparis zeylanica*, Linn , Fl Br Ind , I , 174
 Syn — *C ACUMINATA*, Roxb *C BREVISPIA* DC
 Vern — *Kalo kera* BENG , *Authoondy kai* TAM
 References — *Voigt Hort Sub Cal* , 74 , *Dals & Gibs* , Bomb Fl , 9 ,
Balfour , *Cyclop*
 Habitat — Common in the Carnatic and Malabar, occasional in the
 Western Dekkan and in the drier parts of Ceylon
 Food — The green fruit is pickled

FOOD
Pickled
442

CAPSELLA, Mærch , Gen Pl , I 86

- 443 *Capsella Bursa pastoris*, Mærch , Fl Br Ind , I , 159 , CRUCIFERÆ
 SHEPHERD'S PURSE , PICKPOCKET, Eng , BOURSE DE PASTURE,
 Fr , HIRTENASCHE, Germ
 Habitat — A weed in the vicinity of cultivation throughout the tem-
 perate regions of India , particularly abundant on the N W Himalaya

MEDICINE

444

OIL

445

FOOD

446

natives as a pot herb "

- 447 *CAPSICUM*, Linn , Gen Pl , II , 892

be given to all the species at ke

- 448 *Capsicum annuum*, Linn , DC Prodr , XIII Pl 1 412 , SOLANACEÆ
 RED PEPPER
 Vern — *Mattisa wánggrá lol mirch marcha mirch gáchmirch* HIND

DURR
 References — *Roxb Fl Ind Ed C B C* 193 *Stewart Pb Pl* 156

428

ca
ion
SeraSt
DoC. 4⁸TIMBER
429
DOMESTIC
430

Capsicum or Red Pepper.

CAPSICUM
annuum.

Habit.—A native of equatorial America, most probably of Brazil. Commonly cultivated for its fruit throughout the plains of India, and on the lower hills such as in Khasi, and in the Chenchu valley up to an altitude 6,500 feet. When grown on the hills it is said to be very pungent. There are seven varieties, differing chiefly in the length, shape, and colour of the fruit, some being round, others oblong, obtuse, pointed or bifid, smooth or rugose, and red, white, yellow, or variegated. It is probable that most Indian authors have confused this species with *C. minimum*, which see.

History.—"This species has a number of different names in European languages, which all indicate a foreign origin, and the resemblance of the taste to that of pepper. In French it is often called *poivre de Guinée* (Guinea pepper), but also *poivre du Brésil, d'Inde* (Indian, Brazilian pepper), &c., denominations to which no importance can be attributed. Its cultivation was introduced into Europe in the sixteenth century. It was one of the peppers that Piso and Maxgraw saw grown in Brazil under the name *guja* or *guaya*. They say nothing as to its origin" (*DC. Orig. of Cult. Pl.*) "Chillies are not mentioned by any Sanskrit writer, consequently their introduction into India must have taken place at a comparatively recent date. It is probable that the Portuguese brought the fruit from the West Indies. Up to the present time the cultivation of the plant is carried on more extensively at Goa than at any other place on the

449

apart when they attain a growth of three inches; and afterwards put out into a bed of rich light earth when they attain six inches in height, giving them a good supply of water and keeping them clear from weeds" (*The Gardener*).

450

Medicine.—Dr. Stewart says that the fruit is used externally in the form of plasters and taken internally in cholera; it is eaten from a conviction that it counteracts the effects of bad climates.

MEDICINE.
Plaster.

451

As a drug, red pepper is considered by the natives as stomachic and stimulant, and is used externally as a rubefacient (*Dymock*). "It has been employed with success as a topical application to elongated uvula

Lozenge.

452

white. Made into a lozenge, remedy for hoarseness with a putrid sore-throat whether fusion of red pepper are often

snake-bite

Chemical Composition.—"Bucholz in 1816, and about the same time Braconnot, traced the acidity of capsicum to a substance called *capsicin*.

CHEMISTRY.

453

C. 453

CAPSICUM
annuum.

Capsicum or Red Pepper.

CHEMISTRY.

It is obtained by treating the alcoholic extract of ether, and is a thick yellowish red liquid, but slightly soluble in water. When gently heated it becomes very fluid, and at a higher temperature is dissipated in fumes which are extremely irritating to respiration. It is evidently a mixed substance consisting of resinous and fatty matters.

"Felleter, in 1869, exhausted capsicum fruits with dilute sulphuric acid and distilled the decoction with potash. The distillate which was strongly alkaline and smelt like *conine*, was saturated with sulphuric acid, evaporated to dryness and exhausted with absolute alcohol. The solution, after evaporation of the alcohol, was treated with potash and

isolating it in sufficient quantity to allow of accurate examination.

"Dragendorff states (1871) that petroleum ether is the best solvent for the alkaloid of capsicum, he obtained crystals of its hydrochlorate, the aqueous solution of which was precipitated by most of the usual tests, but not by tannic acid.

"The colouring matter of capsicum fruits is sparingly soluble in alcohol, but readily in chloroform. After evaporation an intensely red soft mass is obtained, which is not much altered by potash, it turns first blue, then black, with concentrated sulphuric acid, like many other yellow colouring substances. By alcohol chiefly *palmitic* acid is extracted from the fruit, as shown by Thresh in 1877.

The crystals melted at 38°C . On keeping them for some days, at the

caustic lye removes *capsicin*, which is to be precipitated in minute crystals by passing carbonic acid through the alkaline solution. They may

Cayenne Pepper or Chillies.

CAPSICUM
frutescens.

CHEMISTRY.

be purified by recrystallizing them from either alcohol, ether, benzene, glacial acetic acid, or hot bisulphide of carbon; in petroleum *capsaicin* is but very sparingly soluble, yet dissolves abundantly on addition of fatty oil. The latter being present in the pericarp is the cause why *capsaicin* can be extracted by the above process.

"The crystals of *capsaicin* are colourless and answer to the formula $C_{15}H_{25}O_3$; they melt at $59^{\circ}C.$, and begin to volatilize at $115^{\circ}C.$; but decomposition can only be avoided by great care. The vapours of *capsaicin* are of the most dreadful acridity, and even the ordinary manipulation of that substance requires much precaution. *Capsaicin* is not a glu oxide; it is a powerful rubefacient, and taken internally produces very violent burning in the stomach" (*Pharm. ograp.* 11).

Special Opinions.—"Stimulant and rubefacient, useful in dyspepsia; recommended in infusion as an external application to the eye" (*Assistant Surgeon Nela Sirg, Shahranpur*). "Chiefly used as a condiment and considered to be stomachic" (*Assistant Surgeon Anund Chunder Mockersy, Neathally*). "Antimalarious to a certain extent" (*H. D. Masani, Surgeon, H. M.*).

native, cooling medicine. The seeds is used in cholera. In and sore-throat. It is an in Deccan, Guzerat, and Cutch" (*Bombay, Bhuj, Cutch*). "The known, are powerfully irritant by natives to dog-bites. An infusion made with 4 drams of chillies and a bottle of boiling water has been found useful in severe sore-throat" (*Assistant Surgeon Bhagwan Dist, Rawal Pindi*). "In delirium tremens in 20-grain doses" (*Surgeon-Major George Cumberland Ross, Delhi*). "Is used in liniments as a rubefacient; in cholera pills with camphor and

Food.—The fruit when green is used for pickling and when ripe is found for

FOOD.
454

or daily
curries,
ginger,
poor can

obtain to eat with their rice (*Balfour's Cyclop.*) Dr. Dymock gives the value of Ghâti chillies at Rs 3½ per maund, and Goway, Rs ½ to 4 per maund of 23lb in Bombay.

Capsicum, fastigiatum, Blume. See *C. minimum, Roxb.*

C. frutescens, Linn; Fl. Br. Ind., IV, 239.

455

SPUR PEPPER, CAYENNE PEPPER, GOAT PEPPER, AND CHILLIES.
THE SHRUBBY CAPSICUM

Ve-

CAPSICUM
frutescens

Cayenne Pepper or Chillies

ladamera china, MAL, *Menashina kays*, KAN, *marichu phalam*,
brahu or *bran maricha* ? SANS *Filfil-ahmar*, ARAB, *Fulfil i surkh*,

Supposed to have
in South America.
pecies of Capsicum,
now cultivated in India, have no Sanskrit names. Of the Indian culti-
cates the large t. being

the sun

Cayenne
Pepper.
456
Chillies
457

Opinions differ slightly as to the plants which afford Cayenne pepper. Speaking of this species, DeCandolle says "The great part of the so-called Cayenne pepper is made from it, but this name is given also to the product of other peppers. Roxburgh, the author who is most attentive to the origin of Indian plants, does not consider it to be wild in India" (*Orig Cult Pl*). Simmonds writes that "the Cayenne pepper of commerce is obtained chiefly from the pulverised chillies or fruit pods of one or two species of Capsicum (*C. annum*, Linn., and *C. fastigiatum*, Blume). So also in the *Kew Official Guide* (p 100) the dried and pulverised rind of the pods of *C. annum* and its allies is said

MEDICINE
458

(Atkinson)

Special Opinions — § 'When taken in curry in unusual quantities,

tent
In
ague

Seed,
459
Cholera
mixture
460

in gargles for sore-throat (*Brigade surgeon A. M. S. H. CORE SURGEON et al*
abad) "A powerful stimulant used as a gargle in sore throat, also in

Chilli
Vinegar
461
Chilli
Extract
462
Powder
463

C. 463

Bell Pepper Bird's eye Chili.

CAPSICUM
minimum

ana in 1867 in the collection forwarded to the U. S. Lib. bot. on (See
m. rds Trop. Agric. 45)

The pod is dried on a hot plate or in a slow oven and then pounded
in a mortar. The powder is then passed through a hand mill, but it is
brought to the fine powder state; thereafter it is well sifted and preserved
in corked glass bottles for use (Treatise of B. Lang)

Capsicum grossum, Willd., Fl. Br. Ind. II 239

BELL PEPPER

Vern.—*Acra mara* h. Beng. Hind.

References.—*Forsk., F. Ind. Ed. C.B.C. 191 F. & H. Hand. Pharm.*
ma. 4452 Dymock & al. 3rd W. Ind. 2nd Ed. 440 F. & H. Wood P. m. b.
Prod. 222 L.C. Orig. Cult. Pl. 270 Balfour Cylop. 1st Sm. th. D.
51; S. Monrods Trop. Agric. 59

Hab. tat.—Not much cultivated in India; native place uncertain.

Food.—Cultivated to a limited extent in gardens, but chiefly for Euro-
pean, who either cut this capsicum in stew or it is opened, stuffed with
certain spices, and pickled in vinegar. The thick fleshy skin is not so
hot as that of the other species.

C. minimum, Pers. Fl. Br. Ind., IV 239 Wright, Ic. t. 1617

BIRD'S EYE CHILLI

Syn.—*C. FASTIGIATUM Plume; C. BACCATUM Hall*

Vern.—*Gd. h. mara* h. Hind.; *Dhan-jung ka murich* *lanh morich* *Idi*
morich *BENG.*; *Lal mara* h. *marcha* *Cuj.*; *Merch* *lal m. rech* *DUK*
Us-mulaghai *TAM.*; *Sudm. rupa* *ka a* *TEL.*; *Chale* *Ida-ach* *na*
MAL.; *ka pal melaka* *MALABAR.* *Fili surah* *PERS.*; *Fili i-ahmar*
(red pepper) *ARAB.*; *Mris* *S. NG.*; *Nayd si gna yoke gna yoke no-*
pmyan *nayon* *BURM.*

References.—*Harb. Fl. Ind. Ed. C.B.C. 191 1047 Hort. Sub. Cal.*
510 Pharm. Ind., 180 Fl. & H. Hand. I. harma. og. 452 453; U. S.
D. pens. 15th Ed. 349 Bentl. & Tr. m. Med. Pl. t. 168 U. C. Dutt
Mat. Med. Ind. 221 Dymock Mat. Med. W. Ind. 1st Ed. 531;
Haring Basar. Med. 35, Diden Lovell Pb. Prod. 363; Spous
En yelo 1803; Balfour Cylop. Sm. th. D. c. 91; S. Monrods Trop.
Agric. 4790

Hab. tat.—Cultivated throughout India but not extensively. clo. ely

464

FOOD
465

466

energetically added to snapshots it greatly increases the reactivity
"Acts as an acid stimulant and externally as a rubefacient used in

CARALLIA
integerrima

Small Chillies, Carallia

MEDICINE

putrid sore throat and scarlatina, also in ordinary sore-throat, hoarseness, dyspepsia, and yellow fever, and in diarrhoea occasionally, also in piles' (*Buden Powell*)

Mixture
469

an excellent gargle in the sore throat which accompanies this disease as well as in ordinary relaxed sore-throat, hoarseness, &c" (*Waring, Bazar Medicines*)

FOOD
470

Food—This small "chilli" is rarely used by natives, but by Europeans is steeped in vinegar and mixed with salt, in this form it is employed as a seasoning in stews, chops, &c

CARAGANA, Lam, Gen Pl, I, 505

471

Caragana pygmaea, DC, Fl Br Ind., II, 116, Royle, III, t 34, fig 2, LEGUMINOSÆ

Vern—*Tama, dama trāma, LADAK, Shmalak SIND*

References—*Brandis, For Fl, 134, Stewart, Pb Pl, 61, Balfour, Cyclop.*

Habitat—A low shrub very much resembling furze. It inhabits the dry highlands of the Western Himalaya, altitude 8,000 to 17,000 feet

FOOD
Roots.
472
FODDER
473

Fodder—It is browsed by goats and is much valued for fuel in the treeless regions where it is met with. Balfour states that in China the roots of *Caragana flava* are eaten in times of scarcity

CARALLIA, Roxb, Gen Pl, I, 680

474

Carallia integerrima, DC, Fl Br Ind., II, 439, Wight, Ic, t 605, Beddome, Fl Sylv, t CXCVIII, RHIZOPHOREÆ

Syn—*C. LUCIDA, Roxb, Fl Ind Ed C B C, 396 Kurz 1, 451*

Vern.—*Kierpa BENG, Jur, KOL, Palamkat NEPAL, Kujitekra ASS, Punschi BOMB, Pansi phansi MAR, Karalli, TEL And punar, phansi KAN, Dawata davelite, SING, Bya ARRACAN, Manetoga, mani-am g BURM*

References—*Brandis For Fl 219, Gamble, Man Timb, 177, XY, Thwaites En Ceylon Pl 120, Dals & Gibbs Bomb Fl 96, Longt, Hort Sub Cal, 42, Royle, Ill Him Bot, I, 210, Lisboa, U Pl, Bomb 73, Balfour, Cyclop*

Habitat—An evergreen tree with thin, dark grey bark, found in the Eastern and Western moist zones, particularly in the Eastern Himalaya, Bengal, Burma, South India, the Andaman Islands and Ceylon

TIMBER
475

Structure of the Wood—Sapwood perishable, heartwood red very hard, durable, works and polishes well, weight from 42 to 51 lb per cub c

(Beddome)

C. 475

The Monkey's Horn; Carapa.

CARAPA
moluccensis.CARALLUMA, *R Br., Gen Pl., II., 782*

Fleshy, erect nearly leafless herbs, with very thick subterrene or angular stems. The generic Carallum is said to be derived from a South Indian vernacular name.

Caralluma adscendens, *Br.; Fl. Br. Ind., IV., 76; ASCLEPIADEAE* 476

Vern.—*Calamulayan*, TAM

References.—*Nervat, Fl. and Drugs, Sind, 1st, Balfour, Cyclop*

Habitat.—Met with in arid places in the Dekkan Peninsula

Food.—This fleshy plant is often eaten by the Natives in the form of pickles, or is made into chutney. 477

C. edulis, *Berth; Fl. Br. Ind., IV., 76* 478

Syn.—*HOLCEROSIA EDULIS*, *Edg*

Vern.—*Chang, chunga pipid, pipid, pipid, pipid, pipid, pipid, pipid, pipid, pipid, pipid*, Pa

References.—*Swartz, Ib. Fl., 141; Aitchison, Cat., Pb Fl., 90; Murvar, Fl. and Drugs, Sind, 1st; Baden Powell, Pb Pr., 24, Balfour, Cyclop*

Habitat.—Found in the arid tracts of the Punjab and Sind

FOOD
479

C. fimbriata, *Wall; Fl. Br. Ind., IV., 77* 480

MONKEY'S HORN

Vern.—*Matar-sing*, BOMB

References.—*Dalziel & Gibb, Bomb Fl., 155; Voigt, Hort Sub Cat., 535; Lisboa, U Pl., Bomb., 185*

Habitat.—Met with in arid rocky places of the Dekkan Peninsula, from the Konkan southwards, and also in the Ava district of Burma.

Food.—In the Bombay Presidency the plant is eaten as a vegetable

FOOD.
481

Carambola. See *Averrhoa Carambola*, *Linn*, GERANIACEAE

CARAPA, *Aubl., Gen Pl., 338*

Carapa moluccensis, *Lam., Fl. Br. Ind., I., 567, Bedd., Fl. Sylv., 136, MELIACEAE* 482

Syn.—*C. OBOVATA*, *Bl. (Kurz, 1, 226); XYLOCARPUS GRANATUM*, *Koen*

Vern.—*Poshur, pussur, BENG. Kandalanga, TAM; Pinlayo ing, pinl*

...

Habitat.—A moderate sized evergreen tree of the coasts of Bengal, Malacca, and Ceylon

semi-solid fat This
as a hair-oil, and also

GUM
483
OIL
484

for burning purposes

C. 484

CARBONATE OF LIME.

Indian Lime

Lime.

IV. LIME is an oxide before being slaked with water to its corrosive property. OF LIME deprived of its into CALCIUM HYDRATE (CaH_2O_2) which on being mixed with sand forms mortar or cement "As an earth, lime is properly disseminated in nature, as a rock, it enters largely into the composition of the earth's crust, it is less or more diffused in all its waters, it forms the principal ingredient (earth of bone) in the skeletons of the larger animals, and is secreted by many classes of the invertebrates to form their shells, crusts, shields, corals, and other means of protection Economically it is also of vast importance, being used in the manufacture of mortars and cements, in tanning, bleaching, deodorising, and the like, and also in agriculture as a fertiliser or promoter of vegetable decay" (Page)

FORMS OF LIME USED IN INDIA

There are three kinds of lime used in India (a) lime prepared from limestone, (b) lime found on the surface of the ground and known as *kankar*, and (c) lime prepared from fresh-water or marine shells.

LIMESTONE
490

(a) LIME FROM LIMESTONE

Speaking of the distribution of limestone and marble, Mr Ball in his "Economic Geology" says "Limestones can hardly be said to be absent from any of the formations in India, though in some they are either rare or so impure as hardly to deserve the title In the metamorphic series, bands of crystalline limestones occur locally in some abundance,

found in the Bhaner group, where they sometimes attain as great a thickness as 260 feet, and are used both as a building stone and for lime

"In the Gondwana series, limestones are rarely met with, and then chiefly in the Talchir and Raniganj groups, where they occur as lenticular or concretionary masses

"In the rocks of cretaceous age, within the peninsula, limestones of both sedimentary and coral reef origin occur The other sources of lime are principally sub recent and recent tufaceous deposits of *kankar*, travertine, &c

"In the extra peninsular regions the principal formations containing limestones are of carboniferous, jurassic cretaceous, and nummulitic ages Another source of lime is recent coral On the whole it

rk, a de-
to prov-hinopoly,
open ng

Indian Lime

CARBONATE OF LIME

of the railways, have largely replaced the <i>kunkir</i> formerly employed for building purposes in the Presidency	LIMESTONE
<i>In Bengal</i> , although of India, workable stone supplies are practically and Lohardaga. In the peculiar interest because of their proximity to iron ore	492
<i>In the Central Provinces</i> , limestones occur at Simlulpur, Raipur, and Jabalpur, the latter consisting of the famous marble rocks of that name. Limestones also occur throughout the Vindhya range, the most accessible being in the neighbourhood of Warora. At Raipur a stone suitable for lithography has been found	493
<i>In Kutch</i> , limestones of different ages are met with, but those most esteemed belong to the lower Jurassic group	494
<i>In Southern Afghanistan</i> limestones of cretaceous age abound, and in Baluchistan nummulitic limestones are found in the eastern frontier as well as in Northern Afghanistan. In the latter the Safed Sang takes its name from a beautiful Statuary marble	495
<i>In the Panjab</i> , marbles and limestones in considerable variety and from different geological formations are met with	496
<i>In the North-West Provinces</i> and along the Tarai to Darjiling limestones are not infrequent. An account of these may be found in Atkinson's <i>Report on the Geology of the North-West Provinces</i> . <i>Mallet</i> <i>Speaks of Hima</i> <i>Tal, at</i> <i>Bageswar and Almora, at</i> <i>Baitalghat, and Dhikuli for</i> <i>inagur. Lime is also made</i> <i>at</i> <i>Two kinds of limestone</i>	497
are used in the Tarai for the purpose of building the foot of the Kur. the other is the tufa. this latter kind, however, stone costs at the quarry by the Forest Department averaged at half a rupee per maund for a 100 maunds. Thus the stone is landed at most points in the district for Rs 30 per 100 maunds and including the expense of burning, a maund of lime costs 10 to 12 annas. This lime will bear two or three portions of pounded brick or <i>surki</i> . Second class lime ready for use now costs Rs 25 and delivered in Naini Tal Rs 50 to Rs 100 per 100 maunds, it will, however, only bear a proportion of one part of pounded brick to two parts of lime.	
<i>In Central India</i> at Gwalior an abundant supply of flaggy limestones occurs	498
<i>In Rajputana</i> the Arvali group of transition rocks includes many varieties of marble, some of them being of great beauty. The Jhirri quarries of Alwar afford hard white marble. Black marble is met with at Mandla, near Ramghur, white as well as pink and grey marbles at Raigarh in Taipur. But the most extensive marble quarries of Rajputana are at Makrana in Jodhpur. This marble has been celebrated for ages, the Taj of Agra being built of it.	499
<i>In Bombay</i> , there are numerous localities where limestone occurs but no marble. In the Panch Mehals, good building limestones are obtained but not hydraulic, and in Guzerat more or less calcareous rocks are met with	500

CARBONATE OF LIME.

Indian Lime

LIMESTONE

501

In Assam, in the Brahmaputra Valley, nummulitic limestones occur at several localities, the southern face of the Khásia and Jaintya Hills affording an inexhaustible source of supply, known in trade as Sylhet lime

502

district.

503

In the Andaman Islands, an important supply of lime, for Calcutta, is afforded by the coral reefs

The writer has been favoured, by Mr H. B. Medlicott, with the following brief account of the important commercial limestones of India.—

Lime is a scarce article in many parts of India. Much of the lime used in Calcutta is carried many hundred miles by river and railway. The want of a pure limestone flux at moderate cost has been the chief difficulty in working the iron furnaces in the Raniganj coal-field. The most general source of building lime in India is *kankar* or *kunkur* (meaning gravel), a granular or nodular stone found on the surface and in the sub-soil. It is purely of secondary origin being formed on the spot by the evaporation of the ground-water, containing in solution more

in North Western India, the lumps of kankar often coalesce into a continuous mass, fit for use as building stone. A stone so formed must of

504

505

506

507

508

Port Blair which may prove of economic importance, as it is at about the same distance from Calcutta as Katm, and the lime is of equally good quality

"Other localities where limestone is known are numerous but at present of merely local importance, or in most cases of no value whatever. A full list of them, as far as they are known, will be found in the *Manual of the Geology of India*, Vol III, p 449, *et seq*"

Indian Lime

CARBONATE OF LIME.

(b) KANKAR OR CONCRETIONARY LIME.

KANKAR (KUNKUR).—"Throughout the plains of Upper India the principal source of lime is the *kankar* which is found in nodules and layers of various sizes in the clays of the Gangetic alluvium. It yields an excellent but somewhat hydraulic lime" (*H. B. Medlicott. See also the remarks under Limestone.*)

KANKAR.
509

"*kankar*" (which really means any kind
! for concretionary carbonate of lime,
"all kinds of the country and

and externally of a mixture of carbonate of lime and clay. The more massive forms are a variety of calcareous tufa, which sometimes forms thick beds in the alluvium, and frequently fills cracks in the alluvial deposits or in older rocks

"In the beds of streams immense masses of calcareous tufa are often found, forming the matrix of a conglomerate, of which the pebbles are derived from the rocks brought down by the stream. There can be no

"As a flux for iron, *kankar* has been tried on several occasions, and opinions are somewhat divided as to its applicability to the purpose; but owing to the uncertainty of its composition, it is distinctly less well adapted than rock limestones which have a well-defined average composition, even though in the latter the proportion of carbonate of lime may average something less.

510

"Block *kankar* has been largely employed as a building-stone, more particularly in connection with the Ganges Canal Works" (*Ball*)

Most of the roads in Northern India, and indeed in India generally, are metalled with *kankar*.

(c) SHELL-LIME.

SHELLS.—Ainslie, in his *Materia Indica*, mentions lime produced by burning the sea-shells, called in Tamil *kullingie chunambu* Dr. U. C.

SHELL-LIME.
511

CARBONATE OF LIME.

Indian Lime

SHELL-LIME.

that I have visited by burning the shells of the genus *OSTREA*, which
abo
in a

aba
globosa.

LIME ESSENTIAL TO VEGETATION.

AGRICUL-
TURAL USES
512INDUSTRIAL
USES.

Dye—Lime is universally used by the Māmpuris to assist in the transformation of green into blue indigo and to deepen the blue colour of indigo, and a small piece placed in the mouth of a vessel containing indigo is also supposed to preserve the dye (See *Strobilanthes*). Lime is employed in the Rajshahye district for dyeing thread dark blue, of this

Dye adjunct
513Calico
printing
514

Tans of the North-West Provinces, gives a preparation of blue printing ink of permanent colour. A mixture of 4lb of shell-lime, 10lb of stone lime, and 15lb of impure carbonate of soda (*reh*), with 3 gallons of water, is strained through grass, to this is added 1lb of sulphurate of arsenic and 1lb of indigo, the mixture is then boiled "till it assumes the metallic greenish blue lustre of the peacock's tail. It is then thickened with *babul* gum and is then ready for printing." Sir Edward further remarks

A paint
515Tanning
516

Encycl., II, 1221)

C. 516

Indian Lime.

CARBONATE OF LIME.

MEDICINAL USES

Medicine—According to Dutt, in the *Hindú Materia Medica* (p 82) lime is used internally in dyspepsia, enlarged spleen, and other enlargements in the abdomen, and externally as a caustic. A mixture of lime, carbonate of soda, sulphate of copper and borax, is applied as a caustic to tumours and warts. It enters into the composition of several prescriptions for different forms of dyspepsia, such as *Amrita vati* and *Agnikumara rasa*.

Ainslie says the Vytians prescribe lime water mixed with gingelly oil and sugar in obstinate cases of gonorrhœa. "Mixed with gamboge, quicklime is applied externally to painful and gouty limbs. It is also used as a caustic in the bites of rabid dogs" (*S Arjun, Bomb. Drugs*). The exhaustive account of the medicinal properties of lime given by Dr. Waring in his *Bazar Medicines* (p 85) may be here quoted, since by doing so it will practically be unnecessary to refer to other authors —

MEDICINE.
517

518

lime is deposited at the bottom. In cases of emergency, as burns, &c., half an hour is sufficient for this purpose, otherwise it should be allowed to stand for twelve hours at least before being used. It is only the clear water which holds a portion of lime in solution, which is employed in me-

milk.

519

The dose of the clear water is from 15 to 20 drops or minims in milk, twice or thrice daily.

"In acidity of the stomach, in heart-burn, and in those forms of in-

is best given in milk.

"In diarrhœa arising from acidity, lime water frequently proves useful; it is best given in a solution of gum arabic or other mucilage, and in obstinate cases 20 drops of laudanum with each dose increase its efficacy; it may also be advantageously combined with Opium water. In chronic dysentery the same treatment sometimes proves useful. Enemas

trial in the vomiting attendant on the advanced stages of fever; it has

C. 519

CARBONATE OF LIME.

Indian Lime.

MEDICINE.

been thought to arrest even the black vomit of yellow fever. It is also a

520

charges have in some instances been mitigated and even cured by the use of vaginal injections of a mixture of 1 part of lime water and 2 or 3 of water

"In scrofula, lime water in doses of $\frac{1}{2}$ ounce in milk, three or four times a day, proves beneficial in some cases, it is thought to be especially adapted for those cases in which abscesses and ulcers are continually forming To be of service, it requires to be persevered in for some time. Scrofulous and other *For syphilitic ulcers or chancre* water $\frac{1}{2}$ pint and calcium 30 grains, this, commonly known as black

water either pure or conjoined with oil is sore or cracked nipples it proves very serviceable Diluted with an equal part of water or milk, it forms a useful injection in discharges from the nose and ears occurring in scrofulous and other children

"In Consumption, lime water and milk has been strongly recommended as an ordinary beverage The same diet-drink has been advised in Diabetes, but little dependence is to be placed upon it as a cure, it may produce temporary benefit

or
ful
of

521

effective in preventing a rising in small pox.

LIME AS A CONDIMENT

FOOD
In pan.
522

523

alluding to the use of lime in pan, says, "when used for any lengthened period, it considerably modifies the natural condition of the mucous covering of the mouth, and alters the appearance of the tongue so as to render it useless or fallacious as a means of diagnosis in disease Its use in moderate quantities does not appear to act prejudicially on the system, but when largely indulged in, it lays the foundation of much visceral disease."

Indian Lime

CARBONATE OF LIME.

DOMESTIC AND OTHER USES

Manure.—As a manure, lime plays an important part. It is largely

DOMESTIC.
Manure
524

are not so diversified as is desirable. A dressing from 1,000 to 5,000 lbs of lime may be applied per acre, according to the price at which the lime can be obtained' (H. R. Robertson, *Agriculture*, 13)

Lime is often employed as a deodorising agent. "It is mixed with decaying vegetable matter and with animal bodies, with the view of hastening their destruction and preventing the escape of offensive and noxious effluvia. This effect lime produces by its tendency, in common with the other caustic alkalis, to carry the decomposition through the intermediate stages of putrefaction at once to the ultimate products" (*Morton, Cyclop., Agriculture, Vol II, 266*)

Soap.—Lime is used in preparing soap according to Lunge's method, which is described thus: "A flat-bottomed pan is preferred for making this soap, into which is introduced any given quantity of water and slaked lime equal to 12 per cent of the weight of fatty matter. The whole is to be boiled and stirred when an insoluble hard lime soap and a solution of glycerine are produced, when the latter may be drawn off from the bottom of the pan. A certain quantity of water and commercial carbonate of soda (the latter being slightly in excess of the quantity of lime used) are next added, and the boiling and stirring continued, when the hard carbonate flakes on sufficient the separ. important

Soap,
525

Mortar and Cement.—The use of lime in the preparation of mortars and cements is too well known to require any special description. The following paragraph from *Miller's Chemistry, Part II, 462*, is, however, quoted here, as it will be found instructive: "The great consumption of lime in the arts is for the purpose of making mortars and cements. Pure lime, when made into a paste with water, forms a somewhat plastic mass which sets into a solid as it dries, but gradually cracks and falls to pieces. It does not possess sufficient cohesion to be used alone as a mortar, to remedy this defect and to prevent the shrinking of the mass, the addition of sand is found to be necessary."

Cement.
526

burnt lime, a suitable quantity of water is afterwards worked into it, and it is then applied in a thin layer to the surfaces of the stones and bricks which are to be united. The bricks or stones are moistened with water before applying the mortar, in order that they may not absorb the water from the mortar too rapidly. The completeness of the subsequent hardening of the mortar depends mainly upon the thorough intermixture of the lime and sand."

CARBONATE OF POTASH.

Sources of,

the feet, now employ for *sarkhi* grinding steam power to drive heavy rollers which work in a strong iron basin. For further information see Cement.

527

Carbonate of Potash.

POTASHES, PEARL-ASH; CARBONATE DE POTASSE, *Fr.*; KOH-LENSAURES KALI, *Germ*

Vern — *Sarjika*, *BENO*, *Yon khur*, *ivak chhar* or *ouk chhar*, *HIND*;

Spons Encyclop, p 253, *Balfour's Cyclop*

The mon-oxide of the metal Potassium is known commercially as

Potashes

528

Pearl-ash.

529

Conf. with A

759.)

rapidly absorbs moisture if exposed to the atmosphere, forming thereby a thick oily liquid known as *Oleum tartari per deliquium*. If subjected to dry heat it melts at 800°, but loses a portion of its carbonic acid at still higher temperatures. And decomposes at a higher temperature.

Source of carbonate of po-
Alkali carbonate

aceous annuals contain more pearlash than woody arborescent plants but even of the same plant the succulent young parts are more highly charged than mature tissues. Of different plants pines contain on an average only 0.45 per cent, oaks 0.75 to 1.5 per cent, vine shoots 5.50, ordinary straw 5.8, ferns from 4.25 to 6.26, Indian corn stalks 17.5, nettles 25.03, wheat straw before earing 4.0, wormwood 73.0, and beet about the same amount.

These facts naturally suggest the plants best suited for the preparation of pearlash and the —

Indian Manufacture of

CARBONATE OF POTASH.

clarified and the crystallizable sugar extracted, the remaining liquor is permitted to ferment, that the uncrystallizable sugar may be turned into alcohol and so utilized, but in the stills there will yet remain a waste liquor, and it is in this that abundance of potash salts occur. By evaporating this liquor in a long trough divided across into an evaporating and a calcining section, a salt is finally obtained, consisting of a mixture of potassium chloride, sulphate, and carbonate (together 50 or 60 per cent) with insoluble matter and a good deal of sodium carbonate. The potassium carbonate forms about one-third of the weight of the calcined mass, and arises in a great measure from the destruction, during the calcining process, of the potassium oxalate, tartrate, and nitrate which occur naturally in the beetroot, and, consequently, in the liquor from the still" (*Prof Church in British Manuf Ind.*) This instructive account of the extraction of carbonate of potash from the waste of beet-root has been reproduced here because of its direct bearing on many of the native contrivances employed in India for the preparation of pearlash. It would be almost impossible to over-estimate the extent to which a crude carbonate of potash is employed by the people of India. In another volume under *Alkaline Ashes* (A. 759, also A. 1626) will be found an enumeration of the principal plants used by the natives of India for that purpose, and these should be compared with the plants given under *Berilla* (B. 163) as employed in the manufacture of carbonate of soda. Although in India immense tracts of mountainous land are injuriously covered with various species of wormwood (see *Artemisia*), except as a manure, the ashes of these plants are not apparently utilized. From the high percentage of carbonate of potash which the wormwoods contain, the preparation of pearlash might be confidently recommended to the poorer inhabitants of these regions as a useful new industry. A large export trade might reasonably be anticipated from the Himalayas to the plains of India, if not to foreign countries.

While this is possible, an equally profitable industry might also be organised in preparing the carbonate from the injurious amount of saltpetre

SOURCES OF

Wormwood
Ash.
530

The
Carbonate
from
Saltpetre.
531
from the

Rectification of spirit, Bleaching, and in

Turkey-red
Dyeing.
537
Rectification
of Spirit
538
Bleaching.
539

CARBUNCLE.

Carbonate of Soda: Carbuncle.

CARBONATE
OF POTASH.

wood on the hills and from saltpetre on the plains seems, therefore, worthy of consideration

Yearly Production.—The world's annual production is about one million hundredweights

MEDICINE
540

Medicine.—Carbonate of potash is antacid, then alterative and diuretic, and in over doses poisonous. It is described in Hindu works on medicine "as stomachic, laxative, diuretic. It is used in urinary diseases, dys-

emucous remedy (U C Dutt, *Mai Alea 111na*, 6/)

Special Opinions.—§ "An impure carbonate of potash (*papāda khara*) is also sold in the Bombay bazārs, and is used in the preparation of *papada* (*papun*), or little cakes made with the meal of the different sorts of *dhal* and a little quantity of *asafoetida*, these are given as a digestive, but more as an article of food than medicine, the cakes are roasted over the fire and taken with rice" (C T Peters, *M B, Zandra, South Afghanistan*)

For further information see ALKALINE EARTHS, BARILLA, POTASH, REH and SALTPETRE

541

Carbonate of Soda.

Vern.—*Sajji*, *sajji-mitti*, *sajji khar*, HIND, *Sājji*, BENG, *Chour ki-matti*, *chour ki namak*, DUK, *Sajjekhara*, MAR, *Shach chi karam*, TAM, *Lota sach chi*, TEL, *Qili*, *milhul-qili*, ARAB, *Shikhar*, *tine-gaur*, PERS, *Saryikakshara*, SANS

References.—*Pharm Ind*, 322, *S Arjun*, *Bomb Drugs*, 160, 161, *U. S Dispens*, 1331, *Ure, Dict of Arts and Manufactures*, 854.

MEDICINE.
542

Medicine.—A substance too well known to require any special description (See remarks under the preceding and under BARILLA, SAJJJI, and REH) It is antacid and then alterative. "A paste made of equal parts of *yavakshira* and *sajji kakshara* with water is applied to abscesses for the purpose of opening them" (U C Dutt)

Special Opinions.—§ "Carbonate of soda (impure), *bangada khara*, being the residue left during the manufacture of glass bangles. A second form, which appears to be a purer carbonate of soda, is called *Surati khara*, both are used in the treatment of dyspepsia" (C. T Peters, *M B, Zandra, South Afghanistan*).

CARBUNCLE.

543

Carbuncle.

"The Carbuncle of the ancients is garnet cut, as it is called, *en cabuchon*. The art is still practised in India, and the stones, when of good

Calcutta.
544
South India.
545
Bombay.
546
Burma.
547

The garnet when cut as a Carbuncle is convex above and hollowed out below, so as to leave but a thin layer of the stone through which the light passes, revealing the bright colour. The finest carbuncles are said to come from Pegu and Ceylon. Conf. with *Carnelian*.

C. 547

Heart Pea or Winter Cherry.

CARDIOSPERMUM
Halicacabum

CARCHARIAS, Muller and Henle, Day, Fishes of India, 710

548

Carcharias.—Several species of sharks are employed by the natives of India in the preparation of a medicinal oil. It seems probable that the sharks specially selected for that purpose belong to the genus *Carcharias*. Of these *C. gangeticus* is the most ferocious; it ascends the rivers to about the limits of the tidal influence. *C. hemiodon* also goes up the river specimens having been caught near Calcutta. Several other species are frequent in the Red Sea and Indian Ocean, particularly on the coast of Sind. (See SHARKS AND SHARK FINS.)

CARDAMINE, Linn; Gen Pl., I., 70

Cardamine hirsuta, Linn, Fl Br Ind, I, 138, CRUCIFERÆ

549

References.—*Taxila* Fr Ceylon Pl, 14, Dals & Gibs, Bomb Fl, 7, Steuart, Pb Pl, 13, Treasury of Botany

Habitat.—A herb found in all the temperate regions of India, very abundant in Bengal during the cold weather

Food.—The leaves and flowers constitute an agreeable salad, resembling water-cress

FOOD.
550

Cardamom, see *Amomum subulatum*, Roxb, —the Greater Cardamom, and *Elettaria Cardamomum*, Mason—the Lesser Cardamom

Cardamom seed oil, see *Amomum subulatum*, Roxb

CARDIOSPERMUM, Linn., Gen Pl, I, 393

Cardiospermum Halicacabum, Linn, Fl Br Ind, I, 670, Wight, Ic, t 508, SAPINDACEÆ

551

BALLOON VINE, HEART PEA OR WINTER CHERRY

Vern. *Lataphatki*, *nayaphatki*, *noaphutki*, *sibhul*, BENG, *Hasul kakkal* (seed) PB, *Karolio* GUJ, *Kanpluti*, *bodha*, *sibhal* BOMB, TEL, *Bhal*

1, II,
Dutt,
lyma k,
Arjun
Iraa
fason s

Habitat.—A climbing herbaceous plant plentiful in the plains of India, chiefly in Bengal and the North West Provinces, is distributed to Ceylon and Malacca. Tendrils are modifications of portions of the flower bud fruit triquetrous inflated

Medicine.—The root is used in medicine as an emetic, laxative, stomachic, and rubefacient.

MEDICINE
Root,
552

"It is used as a

C. 553

CAREYA.

The Thistle.

MEDICINE.

Leaves

554

tonic in fever, and a diaphoretic in rheumatism" The dried LEAVES are said to bring on the secretion of the menses The following prescription is given by Dr Dutt as a Hindu cure for amenorrhœa Equal parts of *Jyautishmati* leaves, *sarsikâ* (impure carbonate of potash), *Acorus Calamus* root (*vacha*), and the root-bark of *Terminalia tomentosa* (*asana*) reduced to a paste with milk, taken in doses of about a drachm for three days (*Mat Med Hindus*) "On the Malabar coast the leaves are mixed with castor oil, are inter- Mixed with jaggery and es The whole PLANT, body in bilious affec- it is applied to rheu- matism and stiffness of the limbs The plant, steeped in milk, has

Plant.

555

Juice.

556

Dutt, Drury, S Arjun)

FOOD.

Leaves.

557

Seeds.

558

Food—"In the Moluccas the LEAVES are cooked as a vegetable"

CARDUUS, Linn, Gen Pl, II, 467.

559

Carduus nutans, Linn; *Fl. Br. Ind*, III, 361; COMPOSITE

THE THISTLE

Vern—*Kanchari*, *tiso*, *bâdaward*, *Pa*, *Guli bâdawurd*, KASHMIR

References—*Stewart*, *Pb Pl*, 123, *Baden Powell*, *Pb Pr*, 356, *Dymock*, *Mat Med W Ind*, 396, also 2nd Ed, 466

Habitat—A tall stout thistle, found in the Western Himalaya, from Kashmir to Simla, at an altitude of 6,000 to 12,000 feet, also at Hazara in the Panjab, and in Western Tibet, at an altitude of 13,000 feet.

MEDICINE.

Flowers.

560

FODDER

561

DOMESTIC

562

for *Cratægus*)

Domestic.—Murray remarks that the leaves are employed to curdle milk.

CAREYA, Roxb., Gen. Pl, I., 721

num.
inns
ann
numerous

merous *Stamens*
filaments filiform,
d, crowned by an
absorbed, seeds

A genus, containing only 3 species and these confined to India, named in honour of the Rev. Dr Carey—one of the distinguished Serampore Missionaries—a distinguished botanist and a contemporary of Dr. Roxburgh's

C.562

CARICA
Papaya

The Papaya or Papaw

TIMBER.
577

Strength of the Wood — Sapwood — light brown, heart and
 , beauti-
 ght from
 brought
 Mishmi
 being cut

Drury
 says "the cabinet-makers of Monghur use the wood for boxes. It takes
 a polish, is of a mahogany colour, well veined" It is being tried for
 railway sleepers on the Eastern Bengal and Northern Bengal State Rail-
 ways, but the results of the experiment are not yet known Kurz remarks
 that it is used in Burma for gun stocks, house-posts planking, carts, furni-
 ture, and cabinet-work but is too heavy for such purposes It stands
 well under water and is much admired for axles "It is frequently em-

DOMESTIC
Slow-match
578Tinder
579

ing sawdust (Burm Gaz, 1, 129). The timber was formerly used for
 making the drums of sepoy corps" (Drury, U Pl)

580

Careya herbacea, Roxb, Fl Br Ind, II, 510; Wight, Ic, t 557

Vern — *Bhus dalim*, BENG, Chuwa, NEPAL, *Bhumi darimba* SANS

References — Brandis, For Fl, 237, Kurz, For Fl, I, 499 Gamble,
 Man Timb, 197

Habitat — A small undershrub with pink flowers which appear from
 February to March Common in the Tarai from Kumaon to the Kha a
 Hills and Chittagong Also plentiful throughout the plains of Bengal,
 Oudh, and the Central Provinces

CARICA, Linn; Gen Pl, I, 815

581

Carica Papaya, L, Fl Br Ind, II, 599, PASSIFLOREÆ

THE PAPAW OR PAPAYA TREE

Vern — " " " " " "

Kai au-du, COCHIN CHINA

References — Roxb, Fl Ind, Ed C B C, 736 Brandis, For Fl, 244,
 Kurz For Fl Br Ind, I, 522 Gamble, Man Timb, 197

The Papaya or Papaw.

CARICA
Papaya.

Habitat.—A sub-herbaceous, almost branchless tree, commonly cultivated in gardens throughout India; from Delhi to Ceylon. Fruits all the

by the modern Indian names being evidently derived from the American word *papaya*, itself a corruption of the Carib *ababai*. Ainslie says it is a native of both Indies, an opinion held by many popular writers, but not supported by modern botanists. Atkinson regards it as introduced into India by the Portuguese. Brandis tells us that its Burmese name, *thumbarthi*, means fruit brought by sea-going vessels. In 1626, seeds were sent from India to Naples, so that the tree must have been introduced into India at an early date or shortly after the discovery of America. It is generally dioecious, the female flowers sessile, and the male on long peduncles. Sometimes, however it is monœcious or the flowers even hermaphrodite.

Resin.—Exudes a white resin (*Kurs*).

Fibre.—Dr Dymock recommends the fibre from the stem to be examined.

is r
juic
pos

RESIN.

582

FIBRE.

583

MEDICINE.

Jules.

with good M. Charpentier. Cossignani supports its anæsthetic virtues. Further confirmatory evidence has more recently been added by M. Bouton (*Med. Plants of Mauritius*, 1857, p 65), and it may justly be con-

quired. The above is a dose for an adult; half the quantity may be given to children between seven and ten years of age, and a third, or a tea-spoonful, to children under three years. If it cause griping, as it occasionally does, enemata containing sugar have been found effectual in relieving it. Taking the dose above named as correct, the statement of Sir W. O'Shaughnessy (*Bengal Disp*, p 352) that he had prescribed the

CARICA
Papaya.

The Papaya or Papaw.

MEDICINE.

Juice useful
in Lumbrici.
Seeds.

586

Useful as an
Emmena-
gogue.

milky juice as an anthelmintic, in doses from 20 to 60 drops, without obvious effect, is fully explained. It is principally effectual in the ex-

that they assert that if a pregnant woman partake of them, even in moderate quantities, abortion will be the probable result. This popular belief is noticed in many of the reports received from India. In them it is also stated that the milky juice of the plant is applied locally to the os uteri with the view of inducing abortion' (*Pharm. Ind.*, pp 97, 98)

The opinions so liberally contributed for this publication, by the Indian medical officers (see below), give so much of personal experience regarding the properties of this drug that it is scarcely necessary to abstract an account of it from the publications usually consulted. The following passages may, however, be found useful

A writer in the *Ceylon Observer* (30th July 1884) says: "Papain," papainum, or vegetable pepsin, may be prepared from the juice of the green fruit of *Carica Papaya* by adding alcohol, which precipitates papain. This precipitate is dried and powdered and is then quite ready for use. Brunton considers that, in its peptonising powers, it is superior to the ordinary animal pepsin, and it has the additional advantage of neither requiring the addition of an acid nor an alkali to convert the contents of the stomach into peptone. It is used in France, Italy, and Germany, and has been is an invaluable remedy in the

The author of *Atakela* mentions it as a remedy for hæmorrhoids.

b

p

(

h

r

d

Mat Med)

Leaves

587

The Papaya or Papaw.

CARICA
Papaya.

believed to be the cause of disease Firminger (*Van Gard. Ind.*) says that the tree comes into flower during the rains and emits at times a fine fragrance

CHEMISTRY.
588

larger particles and colored juice yellowish brown
weight of beef in or Below
the boiling put the meat into several pieces, and at the close of the experiment it had separated into coarse shreds In the control experiments made without the juice the boiled meat was visibly harder. Hard boiled albumen, digested with a little juice at a temperature of 20° C., could after twenty-four hours be easily broken up with a glass rod 50 grammes of beef in one piece, enveloped in a leaf of C. Papaya during 24 hours at 15° C. after a short boiling became perfectly tender, a similar piece wrapped in paper and heated in the same manner was hard Some comparative experiments are given in the following table

(1) The milky juice of C. Papaya is (or contains) a ferment which has an extraordinarily energetic action upon nitrogenous substances and like many other

1878) The active principle has since been separated and given the name of *Papaine*, it is now an article of commerce in Europe for medicinal purposes and is said to be capable of digesting 200 times its weight of fibrine, it has been used as a solvent of diphtheritic false membrane in the throat, in old standing cases of chronic gonorrhoea, and where other applications in the following cases of gonorrhoea have been used

MEDICAL
OPINIONS

Gonorrhoea, with great benefit, I had a patient at the grounds of Bankura jail and the milky juice collected 24 hours or so, a dull white preparation for internal use,

The Papaya or Papaw	CARICA Papaya
<p>properties, (P H B, Dacca) The juice has the power of dissolving anthelmintic A is said to be- "The juice lar character" "The flesh"</p>	MEDICAL OPINIONS
<p>it beneficial" (Surgeon Roderick Macleod, Gaya) "Introduced by me in the treatment ruary 1873, is very effect digestion, al geon Major J M Zorab Balasor) "The milky juice of the unripe fruit"</p>	
<p>unripe fruit in effective remedy drachm three tit Provinces) resorted to by irritant and is M.</p>	<p>John Gomes, Esq, Medical Store- internally it produces abortion produce abortion Fruit eaten" J, Salem) "The unripe fruit made into a curry, is eaten by women to excite secretion of milk It also has the property of making meat of any kind tender when cooked with it (Honorary Surgeon P Kinsley Chicacot, Ganjam Madras)</p>
<p>20 grains for dyspepsia (Apothecary Thomas Wurd, Madanapalli, Cud dapa) "The peculiarities of this fruit and its effects as a solvent of meat require to be scientifically investigated" (Surgeon General William D R Thomson M D, C I E, a poultice have an excellent The inspissated juice of the</p>	<p>an tree in of a North dose 5 to</p>

CARICA
Papaya.

The Papaya or Papaw.

MEDICAL
OPINIONS.

other methods were used the matter is open to doubt" (*Surgeon W. G. King, M B., Madras*) "The leaves are used externally for nervous pains. The leaf may be either dipped in hot water or warmed over a fire and applied to the painful part" (*Surgeon-Major W. Nolan, M D., Bombay*) "The seeds are considered to be anthelmintic" (*Surgeon Major F. Robb, Ahmedabad*).

The above opinions show how widely and uniformly the properties of the *papaya* are believed in by Native and even by European Medical Officers.

FOOD
Ripe fruit
580
Green fruit
Curries and
pickles.
590
Other modes
of preparation
591

Food—When ripe the fruit attains the size of a small melon, the interior is soft, yellow, and sweetish, eaten by all classes and esteemed innocent and wholesome. When green it is cooked by the natives in their curries and also pickled. The ripe fruit has a flavour peculiar to itself, the better qualities are eaten without sugar, and by many persons are ranked among the first of eastern fruits. By others the *papaya* is eaten with pepper and salt. The seeds have a pleasantly pungent taste, not unlike mustard, hence in all probability the idea occasionally alluded to that this is the mustard tree of the scriptures. Lisboa says the fruit has a sweetish taste and makes an excellent tart. When boiled in slices it is eaten as a vegetable. Don says that in South America the fruit after being boiled and mixed with lime juice and sugar is used in place of apple sauce. Sloane remarks that the unripe fruit is cut into slices and soaked in water till the milky juice is removed. It is then boiled and eaten as turnips or baked as apples. A few drops of the milky sap of the papaw is said to render meat tender. The author of the *Makhzan* recommends that for this purpose the juice should be mixed with fresh ginger. In Barbadoes the flesh of animals is reported to be hung on the tree over night in order to soften it. This idea prevails all over India and is doubtless often resorted to by domestic servants. Drury confirms this and states that he has personally tested the accuracy of the popular notion. Dr. John Davy (*Edin Ph I, 1855*) declares that this is due to accidental causes. According to some writers the best plan to soften meat is to wrap it overnight in the *papaw* leaves, or to drop a little of the fresh juice into the vessel in which the meat is being cooked. Brandis mentions another process, namely, to wash meat with water impregnated with the milky juice. It is even stated that meat is rendered tender by causing the animals to eat the seeds before they are killed. The best qualities of *papaw* are said to be obtained from Singapore and Moulmain stock. "The green fruit, when peeled, boiled, cut into small pieces, and served with sweet oil, vinegar, salt and pepper, serves as a very palatable vegetable, and is very similar to squash in taste" (*Mr. L. Liotard*).

Juice.
592

Structure of the Wood—The stem of this fast-growing tree is too spongy and fibrous to be regarded as affording timber. Gamble describes it as soft wooded.

TIMBER
593

Domestic—The juice is used by freckles. It is also exceedingly applied to the skin (*Treasury of* by the Negroes in washing linen as a substitute for soap (*Unsungness*)).

DOMESTIC.
594

C. 594

The Blistering Papaya of Brazil

CARISSA
Carandas

Carica spinosa.

A branching tree met with in Guiana and Brazil, has a much more acrid juice than the other species. If dropped on the skin it causes disagreeable blisters. The fruit is not eaten, and its flowers have a cart
dru;

MEDICINE
Juice.

595

CARISSA, Linn, Gen Pl, II, 695

A genus of densely branched, spinous, erect shrubs, belonging to the APOCYNACEÆ. There are some twenty species African Asiatic, and Australian. Sir J. D. Hooker remarks of the five Indian species that they are probably mere forms of one or two very variable plants.

axillary, pedun-
thers included,
very 2-celled, 2
Ovules 1 4 in
tely attached to

the septum without a wing or pencil of hairs

Carissa Carandas, Linn, Fl Br Ind, III, 630, Wight, Ic, t 426, APOCYNACEÆ

596

Syn —C CONGESTA, Wight, Ic, t 1289, Bedd, Fl Sylv, Man, 156, Anal t 19, fig 6

Vern —Karaunda, karunda, or karonda, garinga karrona, timukhia,

R

Botany, Firminger, Man Gard, 256

Habitat —A dichotomously branched bush, cultivated for its fruit in most parts of India, said to be wild in Oudh, Bengal, and South India.

DYE.

Fruit

597

MEDICINE

Fruit

598

Root

599

used in the form of curry and chutney by the natives" (Assistant Surgeon Anund Chunder Mukerji, Noikhalley). "Antiscorbutic, expector-

C. 599

CARISSA
spinarum

The Karunda.

MEDICINE

ant" (Surgeon W Barren, Bhuj, Cutch). "The juice is irritant and capable of producing ulcers on the skin when used with food, and has, I

of M. Zorab, Balasore

much used at the cor

P N Mukerji, Cuttack, Orissa)

FOOD

Pickle

600

Preserves

601

TIMBER

602

DOMESTIC

Fences.

603

Food —The fruit is made into pickle just before it is ripe, and is also employed in tarts and puddings; for these purposes it is superior to any other Indian fruit (Firminger). When ripe it makes a very good jelly (equal to red currant), for which it is cultivated in the gardens owned by Europeans. The natives universally eat the fruit when ripe, and ex-

number of
or b)

Carissa diffusa, Roxb, Fl Ind, Ed C B C, 231, Syn for *C. spinarum*, A. DC, which see

604

C. macrophylla, Wall, Fl Br Ind, III, 631.

Syn —CARISSA LANCEOLATA, Dals, C DALZELLII, Bedd, Fl Sylv, Man, 157

References —Dals & Gibs, Bom Fl, 143, Lisboa, U Pl of Bom, 166.

Habitat. —A large shrub with very strong, curved thorns, common on the Deccan peninsula, Coorg (Heyne), Konkan at Ramghat (Dilzell); Courtallum (Wight). The flowers are much larger than those of the other species.

FOOD

Fruit

605

606

Food —The fruit is eaten, it is about the size of a plum and ripens in May. Beddome says it is superior to that of *C. Carandas*.

C. spinarum, A DC, Fl Br Ind, III, 631, Wight, Ic, t 427

Syn —C DIFFUSA, Roxb

The Flora of British India regards this species as probably only a state of

V

Vern —Karaunda HIND, Gan, garunda, garna PB, San karunda, anka koli, URIYA, Karamadika, SANS, Wakoulu, TEL, Kanuwan, ORAON

References —Roxb Fl Ind, Ed C B C, 231 Brandis, For Fl, 321, Thwaites, Pl, 116, Rod, Bal, U. Pl of

Linn, 166

Habitat —A small, thorny, evergreen shrub, wild in most parts of

thence northward to the mouth of the Hugli (*C. diffusa*).

C. 606

The Carnelian.

CARNELIAN

Medicine.—This plant is mentioned by Baden Powell amongst his

MEDICINE.

Wood.
607

Food.—The fruit is eaten in tarts. The leaves are greedily devoured by goats and sheep.

FOOD.
Fruit
608
FODDER.
600
TIMBER.
610
DOMESTIC.
Fences.
611
Fuel
612

Domestic Uses.—Largely used for dry fences, but spreads so rapidly where clearances have been made that it may impede the reproduction and growth of the forest. It coppices freely and makes excellent fuel.

CARMINE.

Carmines and Carminic Acid.

CARMIN, *Fr.*; KARMIN, *Germ.*; CARMINIO, *It.*

References.—*Balfour's Cyclopæd*; *Use's Dictionary of Arts, Manuf., and Mines.*

A pigment of a bright red colour, made from cochineal and alumina or bichloride of tin. This is prepared by throwing into a decoction of cochineal a certain proportion of the base employed. A salt is produced which is allowed to precipitate in shallow basins. The colourless liquid is decanted and the powder carmine dried and preserved. By the old German process carmine is prepared with alum.

The uses of Carmine have recently been greatly extended. It is employed for making fine red inks and for silk-dyeing. It is the finest red the water-painter, and more especially the miniature painter, possesses. The French carmine and rouge is preferred to the English. See Cochineal.

Carnation. See Clove.

CARNELIAN.

614

The Carnelian and the Amethyst, perhaps, the best known members of the Chalcedony series, are the only ones which are crystalline. The silicic anhydride, but hydrated silica also exists. This has led to the classification of the quartzose minerals into—

1st—Transparent Crystallised Quartz or Anhydrous Quartz, as represented by the ROCK CRYSTALS. These, when violet, are known as the Amethyst, and when yellow or sherry-coloured as the Cairngorm, but numerous intermediate shades also exist from red to black.

2nd—Uncrystallised or Crypto-Crystalline Anhydrous Quartz—This corresponds to the Chalcedony series, but by most writers this is also made to include JASPER, an opaque rock of undefined nature rather than a definite mineral. The term AGATE is sometimes given generically to denominate this series, or Agate and Chalcedony are used as synonymous terms.

3rd—Uncrystalline Semi-transparent to Opaque Hydrated Quartz—The OPAL may be given as the type of this group.

C. 614

CARNELIAN.

The Carnelian.

QUARTZ.

The quartzose stones referable to the above sections are extensively used in India for ornamental purposes, in the lapidaries' art, in decorative architecture, and in the manufacture of cheap jewellery. They are popularly assigned a position with the "inferior gems"—the diamond,

were apparently not known to the ancients, and when first brought to their attention obtained fabulous prices. Pliny mentions that fragments of a small Cambay cup were exhibited in the theatre of Nero, "as if," adds Pliny, "they had been the ashes of no less than Alexander the Great himself." Balfour remarks with much truth that "amongst the people of India the inferior g for its intrinsic price, i in which the chief valt so the trade in these extensive than it is to definitely express. Indeed, the utmost that can be done in this direc-

tion, is to remind the reader of the elaborate decorations of the Taj Mahal of A — — — — —

bay and Broach hold their own carbuncles, carnelians, and agate factures in rock crystals, and Ja Rajputana and the Panjab hav industry in ornamental stones of the foreign trade in certain o of the Indian lapidary indus known under the generic name of *ma-hu-ya*

EXPORTS
615

EXPORTS FROM INDIA OF INFERIOR GEMS—Under the heading JADE STONE Burma is said to have exported, since the beginning of the present decade, the following quantities and values —

YEARS.						Quantity.	Value.
						cwt	Rs
1880-81	3,371	8,03,890
1881-82	7,788	23,01,800
1882-83	4,159	9,00,900
1883-84	3,849	8,12,960
1884-85	3,738	5,60,050
1885-86	3,842	5,00,050
1886-87	2,890	5,61,000
TOTAL						29,637	64,40,650

Thus during the past seven years, British Burma has exported over half a million of pounds sterling worth of jade, an amount which has gone

Exports of Inferior Gems.

CARNELIAN.

EXPORTS

wood 7 64 per cent, cutch 2 56 per cent, and jade-stone 3 51 per cent. From the table given above it will be seen that the exports of jade during that year were exceptionally high, but it may safely be added that jade still holds a position as the fourth or fifth most important article of export from Burma, and that, with the extension of the British frontier, the trade in jade, in rock crystals, and in the nobler gems may in the future be considerably extended. The exceptional development of the trade in 1881-82 was due to the discovery of a new mine and the decrease that followed accounted for by the jade thus sent into the market having proved much inferior to the stone usually exported.

An inferior quality of jade-stone is also found at Mirzapur, and a very considerable trans-frontier trade is done in the Panjáb in Karakash jade from Turkistan, and in jade and imitations of jade or false jade from Kashmir. (See on a further page, under AGATE, variety *plasma*.)

We have alluded to jade in the present connection, not from an established belief that it belongs to the quartzose group of minerals with which we are at present dealing, but because it is one of the so-called inferior gems. The chalcedony and rock crystal gems, however, are even as extensively employed in India as jade-stone, yet it has been found difficult to furnish definite facts regarding the extent of the internal and foreign trade in these. Perhaps the most interesting of the early accounts of the Cambay trade and industry in "Cambay stones" and

during the first few years of the present century

The following figures give some idea of the trade:—

The exports were valued in—

	R
1804 at	49,140
1808 at	54,240
Passing over 70 years they were in	
1874 valued at	84,370
1878 at	50,970
but the returns for the five years ending 1878 show	
an average of	70,000

CARNELIAN.

The Rock Crystal.

ROCK
CRYSTAL.
616

We must now describe, as briefly as possible, the principal quartzose inferior gems —

1st—ROCK CRYSTAL, *Mallet, Mineralogy, 62.*

Vern — *Bilaur*, HIND., *Phatak*, GUJRATI, *Tansala* (smoky Cairngorm),
PB The Burmese name for an Amethyst signifies "egg plant, Sapphire"

References — *Ball's Econ Geol.*, 502, *Balfour, Cycl of India; Bomb Gaz*, VI, 201 *Mason's Burma* (1890), p. 579, *Calcutta Jour Nat Hist*, II *Madras Jour*, Lt and Sci, VII, 172 *Mysore Gaz*, I, 20; *Central Prov Gaz*, 506, *Oldham, Jour As Soc*, Beng, XXIII, 271

CHARACTER OF — When pure this mineral consists chiefly of silicic acid, it is an oxide of the carbon-silicon group. The differently-coloured forms of rock-crystal owe their tints to the presence of small quantities of foreign minerals. These coloured crystals are known by various names such as the Amethyst, Cairngorm, Rose-quartz, Pellucid-quartz, False-topaz or Citrine, Smoky-quartz, Milky-quartz, Prase, Aventurine-quartz, &c

are, how-
ignorant
account
ed from
nged re-
o a tinc-
ture of red sandal, it takes a deeper red tint, into tincture of saffron, a yellow, like the topaz, into a tincture of turnesol, a yellow like the topaz, into a mixture of tincture of turnesol and saffron it becomes an imitation of the emerald. Crystals coloured red are known in France as *rubaces* or false rubies.

PROVINCES WHERE MET WITH — Rock Crystals are very abundantly met with in South India, as, for example, at Vellum in Tanjore, in the Godavery basin, and at Hyderabad. In the Bombay Presidency they are found at Tankara in Morvi. Blocks from one to twenty pounds are found as clear as glass and capable of taking a high polish (rock crystals are also imported into Cambay from Ceylon and China). They are by no means uncommon at Sambalpur in the Central Provinces. Agates and quartz in great quantities are found in the Mahál hills in Bengal. Bannu, Sháhpur, and size have been found d crystals as rubies. large crystals are found in their country. Milky-quartz occurs in Mergui.

stones, the value being about the same as garnets. The crystals of Sambalpur are not worked and they have accordingly no local value. At the foot of the Delhi palace a number of vases, pitchers, drinking cups, &c, cut in transparent quartz were found. These are supposed to have been cut out of large crystals found at the Arvali quartzites in the neighbourhood. The Shans of Upper Burma are said to be experts at making imitation gems from rock crystals.

The Agate.

CARNELIAN.

2nd — AGATE, Mallet, *Minerology*, 70AGATE.
617

The name Agate is supposed to be derived from the *achates* (ἀχάτης) river in Sicily, or from *akik*, a river, in Arabic AGATE Fr, *Achat*, GERM, *Akik*, ARAB, *lamni*, HIND (agate), *Chakmak* (a flint), HIND, *Manka*, HIND (cut agates and beads brought from Kandahar), *Asshar*, HIND (Silica), *Pathannu*, HIND (blood-stone)

They are commonly known to Europeans as Cambay stones or Godavery pebbles

References — Hamilton, Capt (1691), *New account of the East Indies*, I, 143, Hove, Dr (1787), *Explorations in Bombay*, *Sel Rec Bomb Govt*, VI, 1, pp 49 to 51 Kennely, Dr (1826), *Trans Med & Phys Soc*, Calc III, 425 Wallace, Major (1854), *Sel Rec, Govt, Bomb*, V, 111, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935, 936, 937, 938, 939, 940, 941, 942, 943, 944, 945, 946, 947, 948, 949, 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, 984, 985, 986, 987, 988, 989, 990, 991, 992, 993, 994, 995, 996, 997, 998, 999, 1000

SOURCES — Indian Agates are mainly obtained from the mines of Rewa Kantha in the Bombay Presidency, but they exist also in Bengal in the Rajmahal and Singbhum districts, in Hyderabad, and in the Central Provinces at Jabulpur

c
f
s

jamo The colour varies, but is generally a greyish white Both kinds come from north east Kathiawar, near Mahedpur in Morvi, three miles from Tankara Of the stones which lie in massive blocks near the surface, the most perfect do not exceed five pounds in weight, while those of inferior quality, in many cases cracked, weigh as much as sixty pounds These stones are brought to the Cambay market

Like the common agate the moss agate, *sua bhays*, comes from Bud Kotra, three miles from two feet under the surf. a pound to forty pound the common agate W on a base of crystals, s dark green or red-brown moss

CARNELIAN.

The Agate.

AGATE

showing either a dark ground with white streaks, or dark veins on a light black ground."

CHARACTER OF—Agates are concretionary masses or nodules, which occur usually in hollows or veins in volcanic rocks. When cut across the sections show layers. "The colour markings are often in concentric rings of varying forms and intensity, or in straight parallel layers or bands. The colours are chiefly grey, white, yellow or brownish red." The composition of most of the forms of agate and carnelian is from 70 to 96 per cent of silica, with varying proportions of alumina, coloured by

by the more porous layers of the stone; it subsequently becomes carbonised, and thus the contrast of the various colours is heightened. The

clear greyish
various shades.
es are found in
m as found in

2 "Moss agates are such as contain arborisations or dendrites of oxide

blood drops

4 "Plasma, a grass-green stone, found engraved in ruins at Rome, on

employed
d in the
entions a

ple-green

m known

chiefly by its zigzag pattern.

Uses of—Agates are much used for ornamental and domestic purposes
sword h
paper-cl

The Carnelian.

CARNELIAN.

ing in marble and to a certain extent are so employed at Agra and made Agates are also book-binders, they are t, as well as employed for

AGATE.

erial of which the murrade Professor Muller seems to be of opinion that it was flourspar, but Ball very properly comments upon this opinion "if it was obtained at Ujein or Oujein, or any other locality within the trappean area, it was almost certain to have been one of the chalcedonic minerals, viz, carnelian or agate. Flour spar is not known to occur in the trap "

3rd—CARNELIAN (from *Caro-nis*, flesh, in allusion to the colour), Mallet, *Mineralogy*, 72.

CARNELIAN
618

CORNALINE, *Fr*, KARNEOL, *Germ*, CORNALINA, *It*

Vern—*Sang* : *ākh* ' as from
Kandahar) *Pb*, up in
Gujarati One o s Kyut-
three or four s bl red car
nelans

References—Ball, *Econ Geol*, 506 Balfour, *Cycl*, I, 555 & 583, *Encycl Brit* I, 277, *Ure's Dict*, *Arts*, &c, I, 656, Baden Powell *Pb Prod*, 97 Copeland, *Bomb Researches*, Thomson, *Mad Jour*, Lit and Sci, V, 161

be consulted

CHARACTERS OF—Dana defines the carnelian as a reddish variety of chalcedony, generally of a clear bright tint, but it is sometimes of a yellow ish red

Ratan-
es come
rbadda,
Burma,

Mergui, and abundantly so in Japan

ARTIFICIAL COLOURING OF AGATES INTO CARNELIANS—While collecting the pebbles the miners divide them into two primary classes—those that are not improved in colour by burning, and those that are Of the former there are three chief varieties (1) the Onyx, known as *mora* or *bawa ghori*, (2) the Cat's-eye, *cheshamdar* or *dola*, and (3) a yellow half clear pebble called *rori* or *lasania*. All other stones are baked to bring out their colour "During the hot season, generally in March, and

CARNELIAN.

The Onyx and the Jasper.

CARNELIAN

carried to the Nerbadda and floated to Broach Here they are shipped in large vessels for Cambay, and are offered for sale to the Carnelian dealers.

"By exposure to the sun and fire, among browns the light shades brighten into white, and the darker deepen into chestnut Of yellows, maize gains a rosy tint, orange is intensified into red, and an intermediate shade of yellow becomes pinkish purple Pebbles in which cloudy clear bands of the palest flesh id even red, free stone, the more rge, thick, even and variegated

stones are worth little "

Uses of —Carnelians are extensively used for seals. Many of the antique gems are engraved on carnelian

ONYX.
619

4th—ONYX, *Mallet, Mineralogy*, 73

ONYX, ONICE, *Fr*, ONYX, *Germ*, ONIQUE, *Sp*

References —*Ball's Econ Geol* 503, *Mason's Burma*, 581 *B Heyne, Indian Tracts*, p 265, *Newbold, Jour Royal Asiatic Soc*, IV 37

The Onyx resembles the agate very closely, differing only in the fact arranged a flat hexagonal planes This is no as

and such like articles

JASPER.
620

5th—JASPER, *Mallet, Mineralogy*, 76

JASPE, *Fr*, JASPISS, *Germ & Dutch*, DIASPRO, *It*, JASCHNA, *Russ*

References —*Mason's Burma*, 581, *Ball, Econ Geol*, 503

present position
classification It is
ner occurs among
found in Tenas-

serim
and kn
says
I have

Toungoo Mountains" Jasper is abundant in the transition rocks of Kadapah, ribbon jasper is said by Mr. Foote to be largely produced in the Sandur hills in Bellary Bright red jasper is also reported to be abundant in the transition rocks of the Narbada and Sone Valleys. Nodules of jasper are also common in conglomerate rocks

The Opal and the Cat's Eye

CARNELIAN

Uses or — Sometimes employed for seals

JASPER
HELIETROPE

6th—OPAL, Mallet, *Mineralogy*, 80.

OPALE Fr, OPAL, Germ, OPALO, It, Dhudia pathar, Hind
Chalcedony and Opal are sometimes known as Gomed
sannidh, Hind

OPAL
621

This is a compact uncrystalline semi-transparent to opaque hydrated silica. When of milky white colour, opalescent, and exhibiting a rich play of colours, it is the *Noble Opal*. When not opalescent it is the *Common Opal*. The former are obtained chiefly from Hungary and

While
ig come
na, and
ore and

Sitabaldi

On being first dug out of the earth opal is said to be soft, and to harden and diminish in bulk on being exposed to the atmosphere.

7th—CAT'S EYES, Mallet, *Mineralogy*, 69

CAT'S-EYES.
622

This stone is perhaps closely allied to Onyx, but by some writers it is placed nearer rock crystal. It is a translucent quartz, presenting a peculiar opalescent reflection, said to be due to the presence of asbestos. It bears to the eye of a cat, and, their name for the stone, the stones are common and are found are not known. Malabar Coast is generally accepted as a form of cat's eyes. They are sent from Cambay to Bom

and *Lussuria* are names given to a much valued pebble, found scantily with cat's eyes in the Rājppla mines of Bombay (*Select Records, Bomb., New Series, No IV, 31*)

LAPIDARIES' ART

It is not proposed to deal with this subject in the present article, it having been deemed desirable to give in one place under "LAPIDARY" an abstract of all that is known regarding this industry, not merely as practised with the inferior gems but with all gems and ornamental stones. For convenience the reader may, however, be referred to the following works which deal more immediately with the cutting, &c., of the inferior gems —

Bom Gaz, VI, 201. Hoey, *Trade and Manuf of Northern India*, pp 54 and 119. Baden Powell, *Pb Manuf*, 192. Kipling, *Cat Cat Internu Exh*, Pb Section, 28. Burma Admin. Rep., 1887-83, p 64. Hendley, *Indian Art Journ*, Part 2, 28

The above account of the inferior gems was in type before the writer received Mr. Mallet's Vol IV of the "Manual of Geology of India"

CARPETS AND RUGS.

Carpets

CAT'S EYES

He has therefore been unable to do more than give references to Mr. Mallet's account of these minerals, but the reader is referred to that work for fuller particulars.

See "CARBUNCLE," "DIAMOND," "JADE," "GARNET," "LAPIDARY," "PRECIOUS STONES," and "RUBY."

Carob tree. See *Ceratonia Siliqua*, Linn; LEGUMINOSÆ

CAROXYLON, *Thumb*, *Gen Pl*, III., 71.

623 Caroxylon foetidum, *Mog*, *Fl Br Ind*, V, 18, CHENOPODIACÆ
Syn for *SALSOLA FOETIDA*, Del, which see, also under CAMEL FODDER, 39

C. Griffithii, *Mog*, *DC Prodr*, XIII 2, 175

An Afghanistan plant, supposed by Stewart and several other writers to be the botanical name for the Sind and Panjab *lana*, from which *Khar-sajji* is made. This is *Haloxylon recurvum*, Bunge, or the *Salsola lana*, Stocks. *Fl Br Ind* V, 15. See also under Camel Fodder 21, and *Haloxylon recurvum*. Correct the mistake of *Caroxylon Griffithii* into *Haloxylon recurvum* in *BARILLA*, B 163

CARPESIUM, Linn, *Gen Pl*, II, 336

624 *Carpesium abrotanoides*, Linn, *Fl Br Ind*, III, 301, COMPOSITÆ

Syn — *CARPESIUM RACEMOSUM* Wall

Vern — *Wotiangi*, KASHMIR, *Hukmandas*, Pb

Reference — *Baden Powell*, Pb Pr, 357

Habitat — A stout herb met with abundantly in Kashmir, extending along the Himalaya to Sikkim, altitude 5000 to 10000 feet. Some of the specimens so named by Wallich belong to *Rhynchospermum verticillatum* Reinw., a plant which extends to the Khasia Hills and Burma, descending to lower altitudes than *Carpesium*.

DYE
625

medi-
cines

CARPETS

626 Carpets and Rugs

TAPIS, *Fr*, TEPPICHE, *Germ*, TAPYTEN, *Dutch* TAPPETI
It, TAPETES, ALFOMBRAS, ALCITIFAS *Sp*, KOWRU, KILIMI,
Rus

The term Carpet is probably connected with the Latin *tapetes* from whence tapestry

Vern — *Dari* (small rug), *satranji* (large carpet) cotton, *Adlin* (large carpet), *galicha* or *kalicha* (small rug) woollen HIND, *Ghalichah* PERS, *Tanikalam TAM*, *Jamca so*, TEL, *Jemkhani* (in Belgaum), BOMB, *Parmadani*, MALAY

C. 626

Carpets.

CARPETS AND RUGS.

References.—Birdwood, *Memo*, 29th Sept 1879, *Indian Arts*, 284, Vincent J. Robinson, *Eastern Carpets*, also *Journ. Soc. Art* (1886), p. 47; Baden Powell, *Manuf. and Arts, Panjáb*, pp. 10 & 26; Dr Forbes Watson's *Rep.*; Col. Davidson in *Rep.*, Hyderabad Com-

It is not contemplated in the present article to do more than draw attention to the main facts regarding the Indian Carpet Industry, the object being more to indicate the nature of the carpets made, the materials of which they are woven and the dyes employed in their coloration, than

there are carpets woven by the warp horizontal, and others in which it is vertical. The former are chiefly cotton carpets and the latter nearly always woollen, although it is frequent in both classes to use cotton or hemp for the warp and wool or hair for the woof. The warp, with the single exception of the so-called Jabbalpur *dari*, is not coloured, but the woof is so manipulated that in both these classes of carpets it covers the warp. The Jabbalpur *dari* are almost precisely of the same character as the Kidderminster or Scotch carpets—a certain proportion of the pattern being developed by the coloured warp which may be either in bands of different shades or of one uniform colour. In such carpets longitudinal or checked patterns are produced, whereas in the ordinary *dari* or cotton carpet the patterns run across the warp.

Popularly the terms *dari* and *satranji* are applied synonymously to cotton carpets, but in more precise language, the former is a rug or small cotton carpet and the latter a large one. *Dar-finder* is a

vertical warp

the following extracts from the *Bowles' Carpetry* (Vol. VIII, 1881)

1st, **DARIS**—"The cotton carpet loom which lies horizontally along the floor passes round stout poles at either end which are secured by ropes

DARIS.
627

The striped cotton carpet loom differs from the coarse cloth-loom only by

CARPETS AND RUGS

Carpets

CATS EYES

He has therefore been unable to do more than give references to Mr. Mallet's account of these minerals, but the reader is referred to that work for fuller particulars

See "CIRBUNCLE," "DIAMOND," "JADE," "GARNET," "LAPIDARY," "PRECIOUS STONES" and RUBY

Carob tree. See *Ceratonia Siliqua*, Linn; LEGUMINOSÆ

CAROXYLON, *Thumb*, *Gen Pl*, III., 71.

623 Caroxylon foetidum, *Moq*, *Fl Br Ind*, V, 18, CHENOPODIACEÆ
Syn for *SALSOLA FOETIDA*, Del, wh ch see, also under CAMEL FODDER, 39

C. Griffithii, *Moq*, *DC Prodr*, XIII, 2 175

recurvum in BARILLA, B 163

CARPESIUM, Linn, *Gen Pl*, II, 336

624 Carpesium abrotanoides, Linn, *Fl Br Ind*, III, 301, COMPOSITÆ

Syn —CARPESIUM RACEMOSUM Wall

Vern —Woliangil, KASHMIR, *Hukmandus*, Pb

Reference —Baden Powell Pb Pr, 357

Habitat —A stout herb met with abundantly in Kashmir, extending along the Himālaya to Sikkim altitude 5000 to 10000 feet Some of the specimens so named by Wallich belong to *Rhynchospermum verticillatum* Reinw, a plant which extends to the Khasia Hills and Burma descending to lower altitudes than *Carpesium*

DYE

625

are quite unknown to the Hind people

CARPETS.

626 Carpets and Rugs

TAPIS *Ir*, TEFFICHE *Germ*, TAPYTEN *Dutch* TAPPETI *It*, TAPETES ALFOMBRAS, ALCITIFAS *Sp*, KOWRU, KILIMI, *Rus*

The term Carpet is probably connected with the Latin *tapetes* from whence tapestry

Vern —Dars (small rug) *satranjs* (large carpet) cotton, *Kālin* (large carpet) *galcha* or *idicha* (small rug) woollen HIND, *Ghalichah* PERS *Jan kalam* TAM *Jamca* is, TEL, *Jemkhani* (n Belgaum), BOMB, *Parmadani* MALAY

C. 626

Carpets.

CARPETS AND RUGS.

References.—Birdwood, *Memo*, 20th Sept 1879, *Indian Arts*, 284, Vincent F. Robinson, *Eastern Carpets*, also *Journ Soc Art* (1886), p. 447; Baden Powell, *Manufact and Arts, Panjab*, pp. 10 & 26; Dr Forbes Watson's *Rep.*, Col. Davidson in *Rep.*, Hyderabad Comm.

It is not contemplated in the present article to do more than draw attention to the main facts regarding the Indian Carpet Industry, the object being more to indicate the nature of the carpets made, the materials of which they are woven and the dyes employed in their coloration, than to treat of the historic and artistic features of the manufactured articles. Indian carpets may be classified either according to the nature of the materials of which they are made or the manner in which they are woven. There are cotton, woollen, silk, goat's-hair, yak's hair, and pashm carpets, or mixed carpets of any two or more of these materials. Then again, there are carpets woven by the warp horizontal, and others in which it is vertical. The former are chiefly cotton carpets and the latter nearly always woollen, although it is frequent in both classes to use cotton or hemp for the warp and wool or hair for the woof. The warp, with the single exception of the so-called Jabbalpur *darî*, is not coloured, but the woof is so manipulated that in both these classes of carpets it covers the warp. The Jabbalpur *darî* are almost precisely of the same character as the Kidderminster or Scotch carpets—a certain proportion of the pattern being developed by the coloured warp which may be either in bands of different shades or of one uniform colour. In such carpets longitudinal or checked patterns are produced, whereas in the ordinary *darî* or cotton carpet the patterns run across the warp.

Popularly the terms *darî* and *satranjî* are applied synonymously to cotton carpets, but in more precise language, the former is a rug or small cotton carpet and the latter a large one. *Darîs* (=dar, a door,

the rug is not any speak of *darîs* for all cotton carpets and *carpets* for woollen carpets, but more particularly pile carpets or those woven on a vertical warp.

The following extracts from the *Ramhau Gazetteer* (Vol. VIII) express clearly same time they variations, thro

1st, *DARIS*—"The cotton carpet loom which lies horizontally along the floor passes round stout poles at either end which are secured by ropes tied to strong wooden pegs driven into the ground. The weavers crouch on a broad wooden plank placed across the warp. This plank rests on stones at the side of the loom, and as the work goes on is moved forward. The door is formed by the

Persian carpets—by strands of the warp, being cut off, these e instrument called hidden by the weft, which forms the colouring of the carpet. The loom has only two heddles. The striped cotton carpet loom differs from the coarse cloth loom only by

DARIS
627

CARPETS AND RUGS

Carpets

DARIS.

being broader and having a stronger reed or *phani*. The chief aim of the carpet-weaver is to hide completely the white warp-yarn, leaving

the well yarn home thus using a greater length of well yarn than the breadth of the carpet

"A cotton carpet costs from $3\frac{1}{2}d$ to $7\frac{1}{2}d$ ($2\frac{1}{2}$ annas to 5 annas) a square foot

Mr Ba shuttle and
with it the issage of the
woof the w by placing

a long pole, supported at either
whole width of the warp. This
means 'mare,' and so called from
from the 'gori' are hung two

of threads, which are attached to the under and upper threads of the web respectively. When it is desired to cross the threads of the warp, it is simply necessary to pull up one of the bamboos and lower the other, as the bamboos are merely hung to the 'gori' by ropes at each end, the raising and lowering is easily done by tightening or loosening the suspending string by means of a stick attached. No regular shuttle is used. A number of workmen sit in a row, on that part of the *durree* (*durs*) which has already been completed, and pass the thread along between the lines of the warp, from hand to hand. The thread is wound in a long egg shape on an iron skewer or needle

and so on, the threads as they are passed through the threads of the warp are kept close together and the work is rendered compact and

Woollen *daris* are, however, also made in many parts of India, as in the Panjáb and Bombay. Those woven by the aboriginal races are small in size, thick in texture, and even painfully uneven in quality but

Carpets.

CARPETS AND RUGS.

the fact that in India they are often spoken of as Persian carpets, Indian carpets "Carpet-making all the parts of the loom, seems is almost entirely in the hands

PILE
CARPETS.

"Persian carpet-looms differ from plain carpet-looms in having the warp fastened vertically, instead of horizontally, in the absence of heddles and treddles, and in the absence of the reed *phans*. The loom consists of two uprights, from fifteen to twenty feet high and from ten to fifteen feet apart, supporting two beams, one fixed to the lower ends of the uprights and the other moveable. The warp-yarn is passed round these beams forming a continuous loop. On one side of the frame there are three to five uprights, and on the other side there are three to five uprights. A sketch out to it

that have to be taken up for the first row. The workmen repeat in chorus what the overseer says, and fix up the loops, tie a knot, and cut the pieces off. As soon as the first row is ready, a weft-yarn is passed between the two sets of the warp, and is fixed tightly in its place by the aid of a fork like instrument called the heckle. In this manner row after row is laid up, till the whole of the carpet is woven, when it is taken down from the loom, spread on the floor, and sheared.

"Persian carpets vary in price, according to texture and design, from 14s to £1-8s (R7—R14) the superficial square yard. There are (1882) seventy five Persian carpet weavers" (*Bomb. Gaz.*, XIII, District 401)

PRESENT POSITION AND FUTURE PROSPECTS OF THE INDIAN PILE CARPET INDUSTRY.

cotton threads, which are soft in texture and not made hard and tight by over-twisting and sizing. On these wool thread is tied and the allowance of wool is very liberal. The looms are large enough to make any size of carpet, and there are, therefore, no seams. For ordinary English carpets the warp is of the same size as the weft. There is very frequently an under-weft on the back or front of the fabric. Into the weft is woven, a long needle drawal of the knife with which the carpet is sheared. The demand for cheap

cuts away. In an Indian carpet, the whole fabric sinks together under the foot.

"Moreover, very few of the English jacquered power looms are more than three-quarters of a yard wide. Hence the necessity for seams, which are the first places to wear thread bare.

"So it may be said that it is more economical, when buying a carpet, to give three or four times the English price for an Indian hand woven fabric. It is not, of course, contended that bad Indian carpets are

Carpets

CARPETS AND RUGS.

ments of chemical laboratories with their processes introduced, and such a system of organised work set up as completely transformed not only the trade but actually the carpets themselves which were the foundation of

PILE
CARPETS.

Panjab was known beyond its border for the production of carpets, and then only by the productions of the Lahore jail executed for a London firm. There exist no specimens to show that the Multan industry, the

Vincent Robinson's address to the Society of Arts, he is reported to have said—'At one time I attributed this degeneration almost exclusively to the influence of the Government Schools of Art and the jails but at present I feel that it is chiefly due to the influence of English commerce on the historical handicrafts of India.' This seems a much more likely

to as follows in the *Gazetteer for Cambay* —

"Cambay carpets had once a great name. Among the articles mentioned in the proclamation of 1630 'for restraining the excess of private trade to the East Indies,' are rich carpets of Cambay. Later on a chief part of the Senior Factor's duty at Cambay was to buy carpets. This trade has greatly diminished since the British took possession of the place, and the Nawab is now paying the Nawab a

Pile carpets are made of cotton at Hyderabad and at many other places, tufts of cotton yarn being used in place of wool. In the same way

higher prices than the others'

Pile Carpets ARE MADE at a limited number of jails in each Presidency and Province and by a few private manufacturers scattered here and there over the country. The references given to the Gazetteers convey some idea of the distribution of the industry, but it may be concluded that

CART AND CARRIAGE BUILDING. Woods used for.

PILE
CARPETS.

abad and Benares are best known
"Cotton,"
used in carpet
kinds of car-
pets, cotton and woollen, are made can be obtained from the authorities of
the Indian Museum in Calcutta

629

CARPINUS, Linn , Gen Pl, III, 405

Carpinus faginea, Lindl., DC Prodr, XVI, 2, 127, CUPULIFERÆ.

Vern — Shirish, Umar, bijawri Pb Gish, N W P
References — Brandis, For Fl, 492, Gamble, Man Timb, 390

Habitat —A moderate sized tree of the Himálaya, from Kumaon (and
Nepal?) eastward altitude 4 000 to 7,000 feet
Structure of the Wood —Similar to the next species

TIMBER
630
631

C. viminea, Wall , DC Prodr, XVI, 2, 127

INDIAN HORNBEAM
Vern — Charkhri, kái, Pb , Pune, gorá, chamkharak, N W P, Chukissi,
konikath NEPAL
References —Brandis For Fl, 492; Kurs, For Fl Burm, 477, Gamble,
Man Timb 390 Stewart, Pb Pl, 200, Baden Powell, Pb Pr, 572,
Balfour, Cyclop

Habitat —A moderate-sized tree of the Himálaya, from the Ravi east-
ward, from 5 000 to 7,000 feet frequent near water Also met with in the
Martaban Hills, altitude 5,000 to 6,000 feet, and according to Brandis,
on the Khasia Hills

Structure of the Wood —White, shining, no heartwood, warps in
seasoning Weight 50lb per cubic foot, growth moderately slow The
stem is irregular in section, like that of the European Hornbeam, which
it much resembles both in bark, wood, and general appearance Cleghorn
states that it is much esteemed by carpenters

Carrot. See Daucus Carota, Linn , UMBELLIFERÆ

632

CART AND CARRIAGE BUILDING—Woods used for—

During the Colonial and Indian Exhibition two conferences were held
to examine the timbers shown in the Imperial Indian Section Mr.
Hooper, the well-known London Coach Builder, remarked "That a wood
was much wanted in the carriage trade for a light but strong material"

C. 632

The Safflower.

CARTHAMUS
tinctorius.

hot dry weather of the north seasoned the wood in a way very much superior to the artificial methods employed in Europe." The following are the timbers used in India for these purposes, more especially those marked* —

WOOD USED
FOR CART
AND CARRI-
AGE BUILD-
ING

<i>Acacia ferruginea</i> (carts)	* <i>Lagerstrœmia parviflora</i> (buggy)
<i>A. melanoxylon</i> (coaches, railway)	<i>Melia Azadirachta</i> (carts) [shafts]
<i>Albizia amara</i> (carts) [carriages]	<i>Michelia Champaca</i> (carriages).
<i>Barringtonia acutangula</i> (carts)	<i>Milusa velutina</i> (carts)
<i>B. racemosa</i> (carts)	<i>Mimusops Elengi</i> (carts)
<i>Bassia longifolia</i> (carts)	<i>Prosopis spiciagera</i> (carts)
<i>Berrya Ammonilla</i> (carts)	* <i>Pterocarpus indicus</i> (carts, gun-
<i>Briedelia montana</i> (carts)	<i>P. Marsupium</i> (carts) [carriages]
<i>B. refusa</i> (carts)	<i>Pterospermum suberifolium</i> (carts)
<i>Calamus Rotang</i> (carriages)	<i>Sandoricum indicum</i> (carts)
<i>Careya arborea</i> (carts).	<i>Sapindus emarginatus</i> (carts).
<i>Cassia Fistula</i> (carts)	<i>Schleichera trijuga</i> .
<i>Chloroxylon Swietenia</i> (carts)	<i>Shorea robusta</i>
<i>Cynometra ramiflora</i> (carts)	<i>Strychnos Nux-vomica</i> .
* <i>Dalbergia latifolia</i> (wheels, gun car- riages)	<i>S. potatorum</i>
* <i>D. Sissoo</i> (felloes naves, carts).	<i>Tectona grandis</i> (railway car-
<i>Diospyros melanoxylon</i> (carriage)	<i>Terminalia Arjuna</i> . [riages]
<i>Eugenia Jambolana</i> (carts). [shafts]	<i>T. belerica</i>
<i>Ficus bengalensis</i> (cart yokes)	<i>T. Chebula</i> .
<i>Gmelina arborea</i> (carriages, palan- quins)	<i>T. tomentosa</i>
* <i>Heritiera littoralis</i> (buggy shafts)	<i>Thespesia populnea</i> (carts and car- riages)
<i>Hymenodictyon excelsum</i> (palan- quins)	<i>Ulmus integrifolia</i> (carts).
* <i>Lagerstrœmia Flos Regiæ</i> (carts, gun-carriages)	<i>Vitex altissima</i> (carts)
	<i>Xylia dolabriformis</i> (carts).
	<i>Zizyphus xylopyra</i> (carts).

CARTHAMUS, Linn, Gen Pl, II, 483

Carthamus oxyacantha, Bieb, Fl. Br Ind, III 386, COMPOSITÆ

633

Vern — *Kantiari kandiara*, *poli*, *khâresa karar*, *poliyan* P^b

References — *Stewart P^b Pl*, 123, *Aitchison Cat*, *P^b Pl*, 80, *Baden Powell*, *P^b Pr*, 356, *Cooke, Oils and Oilseeds*, 34, *Balfour, Cyclop*

Habitat — Wild in the North-West Provinces and the Panjab, most common in the more arid tracts. Mr C B Clarke thinks this may be the wild form of Safflower

Oil — Dr Stewart says that near Peshâwar and elsewhere in the Panjab, an oil is extracted from the seeds which is used for illuminating purposes, as well as for food. Dr Stocks probably alludes to this when

OIL
634

MEDICINE
635
FOOD
636
637

C. tinctorius, Linn, Fl Br. Ind, III, 386

THE SAFFLOWER, WILD OR BASTARD SAFFRON, AFRICAN SAFFRON, AMERICAN SAFFRON, CARTHAMINE DYE, Eng, CARTAME, SAFRAN BATARD, Fr, DER SAFFLOR, FARBERDISTEL, FALSCH

C. 637

CARTHAMUS
tinctorius.

The Safflower.

SAFRAN, *Germ.*; ZAFFRONE, CARTAMO, *It. & Sp.*; POLERROI, *Russ.*

Vern.—*Kusum, kasumba, kar* (the seed), *barre*, HIND.; *Kusum, kasam-phul, kajirah*, BENG.; *Galap machu*, MANIPUR; *Kásam, kurtam, ku-*
kasdi, RAJ.;
(seed),
a-virai,
nbe (or
kusambi), kasumba, NAN; *jisoon, su, mshu, supan, suoon*, BURM.;
Qurtum, qirtum, usfar, ARAB.; *Kashirah, muasfir, kasakdanah*, PERS.;
Kusumbha, kamalottara, kúshumbha, SANS.; *Kurtum*, EGYPT. The
κῠῆκος, κῠίκος of the Greeks

In Sind the seeds are called *Kardai* (*kurtum*), and in Panjáb *Khar, polian*

References.—*Doct. Et. Ind. Et. C. R. C. Soc. Stenogr.*, Ph. Pl. 12.

gra), 154

the origin of this plant. It has never been found in a wild state, but botanists assign to it an origin in India, Africa, or Abyssinia. De Candolle (*Origin, Cult. Pl.*) says that the grave-cloths found on Egyptian mummies are dyed with carthamine. The Chinese received the plant only in the second century B C, when Chang-kien brought it back from Bactriana. The Greeks and Latins were probably not acquainted with it, although Birdwood and other writers give *κῠῆκος* as its Greek name. As

vated in India.

CULTIVATION.

A few years ago Safflower was an exceedingly important substance, but recently the aniline colours have driven it almost entirely out of the European market. "It still, however, holds its place with the natives as a brilliant though evanescent dye, and as they employ it largely for home use, it must still rank among the industries of the country, as

(*E. James*)
 account of the
 source of oil.
 p, safflower is

CULTIVATION
638

C. 638

The Safflower.

CARTHAMUS
tinctorius.

chiefly grown as subsidiary to some other crop, participating, therefore, in the treatment given to its associate. On this account it is extremely difficult to obtain trustworthy details as to the area under safflower, the method and cost of cultivation, nature of soil necessary, or value of the crop.

(a) *In Bengal* it is chiefly grown in the Eastern division, where even still it constitutes a crop of some considerable value, although greatly decreased through the introduction of aniline dyes. In fact, the Indian safflower

CULTIVATION.

BENGAL.

639

Sown

Oct. to Dec.

rod of sowing

, for example,

are, as a rule,

been left fal-

- - - - -

the

the

res,

ture,

hree

cul-

- - - - -

Gathered
March to May.

even till May. In removing the florets, the flower-heads are not much injured, and as they are fecundated before the time of removal, the seeds continue to mature within their small, white, angular, one-seeded fruits, and are ripe in April to May. They are then collected for the oil crop (*Agric. Hort. Soc. Journ.*, VII, 191)

Area.

£
£
£
£

under this crop in Bengal, but the following figures are quoted from Dr. McOann's work (which is taken from the official returns sent to the Economic Museum): Dacca, 11,500 acres; Gya, 2,260 acres; Monghir,

N.-W. P. AND
OUDH.

640

CARTHAMUS
tinctorius
The Safflower.

CULTIVATION	38 per cent is irrigated land. The mode of cultivation is very similar to what has already been described for Bengal. Light soils are preferred, the plant is rarely grown alone, but is generally sown in the gram fields and disposed like rape in lines. It is extensively grown along with carrots near wells, participating in the rich cultivation bestowed on the latter. It is also associated with cotton, wheat or barley. In the North-West
Sown Oct. to Nov.	
Price	attempts of which have not been ascertained (<i>Outhie and Buller</i>). In a report on the dyes and processes of dyeing in Ajmir it is stated that about 20 000 maunds of safflower are annually received from Delhi, the
BOMBAY. 641	
Area	grain &c., to which last the cultivator looks for his profits. Probably
Sown Oct. gathered March.	
Production	
Varieties Sadhi 642 Kusambyachi. 643	chiefly for its oil-seeds <i>kusumbyachi</i> , a slenderer plant grown for its dye yielding flowers (<i>Bomb. Gaz. XII, 164</i>). In Gujarât the ' <i>kabri</i> ' or so! The land is ploughed. The seed is thrown broad reaped in February. The
PANJAB 644	ited sown were,
CENTRAL PROVINCES 645	during the settlement, 288 acres under the crop and in Hoshiarpur 6,722 acres, especially in the northern part of the Garhshankar <i>Tahsil</i> . It is generally grown as a mixed crop in lines with gram and requires a sandy soil. It is sown in September.
	(c) <i>In the Central Provinces</i> , a little over 6 000 acres are annually under this <i>rabi</i> crop and Raipur is stated to export the dye stuff to about Rs 10,000 a year.
Area.	The half of the crop is sown in the safflower of Bengal, the North may and the Central Provinces, remain the official reports for the incorr either incomplete or quite are than 10 000 acres are under this crop in the remaining provinces of India.

The Safflower.

CARTHAMUS
tinctorius.

(f) In Berar, safflower, however, appears to be cultivated to a very considerable extent. Mr. Liotard informs us that the area under it is over 40 obvious cate a is not seems t of the (g) only in small patches, and there is no export trade.

CULTIVA-
TION
BERAR.
646MYSORE.
647
BURMA.
648

besides, Burma, instead of exporting safflower, receives annually a small

VARIETIES.

Spiny Form.
649

the Deccan alluded to above.

(6) Almost spineless form. This is known as *bhulī* in Patna, *bod-ki* in Berar, *murīā* (or shaved) in Azamghar and the *kusumbyāchi* in the Deccan. A superior quality of dye is derived from this form.

Spineless
Form
650
OUTTURN.

Bengal as from R3 to R15 a bigha.

PRESENT POSITION OF THE SAFFLOWER INDUSTRY.

Simmonds in his *Tropical Agriculture* says: "The cultivation of safflower, known as *Coosumban* in Bengal, is receiving attention at the hands of the local Government. The prosperity of Bengal, though it mainly depends upon the jute trade, is in some measure attributable to the demand for safflower." The writer proceeds to state that the value of the exports from Dacca alone "would be from nine to ten lakhs of rupees—690 lakhs of rupees—tending, in Bengal, a Simmonds' from all India they were F established.

TOTAL
651

**CARTHAMUS
tinctorius****The Safflower****TRADE**

ing" The total exports for 1886 87 were only R83 819 The following table gives the exports from Ind a for the past fourteen years —

<i>Exports</i>		
YEAR	SAFFLOWER	
	Quantity	Value
	Mds	R
1873-74	13 206	7 58 906
1874-75	14 222	6 50,827
1875-76	4 080	1,63 528
1876-77	7,662	3 04 672
1877-78	3 698	1,48 806
1878-79	4 977	1 86 711
1879-80	2 411	1 8 ,456
1880-81	6 675	3 51 157
1881-82	2 293	94 754
1882-83	3 008	92 038
1883-84	2 333	64 497
1884-85	2 167	83 083
1885-86	1 898	68 991
1886-87	2 149	83 819

report in June 1883 that "there is no land under safflower cultivation in

**DYE
652
Preparation****THE DYE**

superior to another—a fact accountable for either by the more favourable nature of the soil or the care bestowed in cultivation If intended for export, after having been dried as above, the florets are either placed in a bag or on a basket or other contrivance permitting of the easy escape of a supply of water which is kept poured on them while beaten or trodden on This process is continued until the water passes through quite

Yellow
653
Red
654

C. 654

The Safflower.

CARTHAMUS
tinctorius.

DYE.

water (if clean) is re-
of mud or other im-
colouring matter is
care must be taken

carefully dried, they are ready for the market.

The *Gazetteer* for the district of Karnal in the Panjáb describes the

"Stripped
Safflower."
655

delay in the preparation injures the dye" This process is so very defec-

Reason of
lower price
paid for
Panjab
Safflower.

Originally
grown for
yellow dye.

Adulteration.
656

to the continuance of even the present greatly reduced trade The

Cowdung.
657
Rice flour.
658
Turmeric.
659

water, it is employed by fraudulent dealers in the adulteration of shag tobacco" (*Morton's Cycl, Agri*)

Tobacco.
660

CARTHAMUS
tinctorius.**The Safflower.****DYE.**Estimation of
Quality.
661

The quality of safflower cake is estimated by dyeing a known weight of cotton; about 4 ounces of safflower will dye 1lb of cotton cloth light pink; 8 ounces will dye it full rose-pink; and from 12 ounces to 1lb will dye it a full crimson. In order to take up this quantity, the cotton must be several times dyed in fresh solutions of the colouring matter.

Chemical History.—It is scarcely necessary to go into great detail re-

Two yellows
and one red.**Carthamin.**

36 per cent. of the florets, while from 0.3 to 0.6 per cent. is the usual amount of Carthamin. The proportion of Carthamin present varies, however, in the inverse ratio to the amount of the soluble yellow principle. The second yellow colour is soluble only in an alkaline liquor.

If the dye-stuff, after the removal of the soluble yellow principle, be acidulated with acetic acid, filtered, and first acetate of lead and next ammonia added, the second yellow colour will be precipitated along with

of the florets). In India pearl-ash is most frequently used, especially that prepared by incinerating *bajra* (*Penicillaria spicata*) or of *chir-chira* (*Achyranthes aspera*), (impure potassium carbonates), but the natural earth carbonate of soda or *sajji-matt* is also frequently employed for this purpose.

EUROPEAN
DYE
SOLUTIONS.
663**EUROPEAN DYE SOLUTIONS.**

Preparation of Dye Solution and European Methods of Dyeing with

pletely alters the colouring matter.

The Safflower.

CARTHAMUS
tinctorius.

DYE.

"Carthamin in a pasty state, as obtained by the process just described, is met with in commerce suspended in water for direct use. The paste is dried upon suitable vessels—porcelain saucers, plates, or even upon polished cardboard

lowing passage may prove useful to Indian dyers or persons interested in the safflower industry. "Carthamus from which the yellow matter has been extracted and whose leaves have been broken down, &c."

in ca
chert
the
long
it is
throu

and passed through fresh baths, continuing to wash and dry it between each operation, till it has acquired the depth of colour that is desired. When it has reached the proper point, a brightening is given it by turning round the sticks seven or eight times in a bath of hot water, to which about half a pint of lemon-juice for each pailful of water has been added.

When silk is to be dyed *ponceau* or poppy-colour, it must be previously boiled as for white, it must then receive a slight foundation of *arnatto*. The silk should not be alumed. The *nacarat*s and the deep cherry-colour are given precisely like the *ponceaux*, only they receive no *arnatto* ground, and baths may be employed which have served for the *ponceau*, so as to complete their exhaustion. Fresh baths are not made for the latter.

"The lightest of all these shades, which is an extremely delicate flesh-colour, requires a little soap to be put into the bath. This soap lightens the colour, and prevents it from taking too speedily and becoming uneven. The silk is then washed, and a little brightening is given it in a bath which has served for the deeper colours.

"All these baths are employed the moment they are made, or as speedily as possible, because they lose much of their colour upon keeping, by which they are even entirely destroyed at the end of a certain time. They are, moreover, used cold, to prevent the colour from being injured. It

CARTHAMUS
tinctorius.**The Safflower.****DEY.**

must have been remarked, in the experiments just described, that caustic alkalis attack the extremely delicate colour of carthamus, making it pass to yellow. This is the reason why crystals of soda are preferred to other alkaline matters

"In order to diminish the expense of carthamus, it is the practice in preparing the deeper shades to mingle with the first and the second bath about one-fifth of the bath of archil" (*Ure's Dict of Arts, Man, and Mines, Vol. I., 661*).

INDIAN DYE
SOLUTIONS.
664**INDIAN DYE SOLUTIONS.**

Indian Method of dyeing with Safflower.—As already stated, the

appear to be known to the natives of India. The dye stuff, after the

the tamarind is employed in place of lime-juice. In Mánipur the fruits of *Garcinia pedunculata* are viewed as superior to lime-juice, and have

Combinations
665

The Safflower.

CARTHAMUS
tinctorius.

DYE.

N W P) With *Terminalia Chebula* or *T. citrina* and protosulphate of iron, safflower gives a dark neutral tint, with safflower, sappanwood, and alum a purplish brown, and with indigo and safflower, greens and purples (*McCann, Dyes and Tans of Beng*)

An almost indefinite series of colours are obtained in India by various combinations with safflower. It should be carefully observed, however,

Use of acids
and alkalis.
666

latter case are
employed also
precipitating
been given it
fabrics, alkali
condition can
peculiarity be

accurate account of the indigenous modes of dyeing with safflower

Fixing Safflower Dye.—It is much to be regretted that no one has as yet discovered a mode of preventing the decoloration of safflower dye, its fleeting property appears to depend on the oxidation of the particles of carthamin held mechanically in the fabric. The inhabitants of different parts of India boast of possessing a secret of effecting this purpose and careful observation on the part of local officers may help to throw some light on the subject. All that is necessary to re-establish the carthamine dye as an important industry is the discovery of some mode of preventing this oxidation of carthamin. The fruit of *Garcinia pedunculata*, a common tree in Assam, has already been alluded to

FIXING
668

the property
extensive use
justifies this
us that the
dyers of Chittagong district claim to be able to produce a "semi-permanent" safflower dye. This is done by adding safflower to water in which

actually made use of now and then as a discharge, so as to produce a yellow pattern upon a pink ground, weak acids do not affect the colours, but chlorine and sulphurous acid destroy the colour at once" (*Crookes*) Safflower dyed fabrics should not be washed with soap, as the colour is removed by the alkali of the soap

Rouge.—It is necessary to refer here very briefly to an important purpose for which safflower is employed, viz, the manufacture of rouge

ROUGE
669

CARTHAMUS
tinctorius.**The Safflower.****DEY.**

must have been remarked, in the experiments just described, that caustic alkalis attack the extremely delicate colour of carthamus, making it pass to yellow. This is the reason why crystals of soda are preferred to other alkaline matters

"In order to diminish the expense of carthamus, it is the practice in preparing the deeper shades to mingle with the first and the second bath about one-fifth of the bath of archil" (*Ure's Dict. of Arts, Man., and Mines, Vol. I., 661*).

INDIAN DYE
SOLUTIONS.
664**INDIAN DYE SOLUTIONS.**

Indian Method of dyeing with Safflower.—As already stated, the

appear to be known to the natives of India. The dye stuff, after the

the tamarind is employed in place of lime-juice. In Manipur the fruits of *Garcinia pedunculata* are viewed as superior to lime juice, and have

Combinations.
665

The Safflower.

CARTHAMUS
tinctorius.

N. W. P.) With Terminalia Chebula or T. citrina and protosulphate of

DYE.

Use of acids
and alkalis.
666

employed along with the alkaline dye solution may have the power of

peculiarity be fully appreciated, otherwise the observer cannot give an

667

FIXING
668of preventing this oxidation of carthamin. The fruit of *Garcinia*
pedunculata, a common tree in Assam, has already been alluded toRouge.—It is necessary to refer here very briefly to an important
purpose for which safflower is employed, *vis*, the manufacture of rougeROUGE.
669

CARTHAMUS
tinctorius

The Safflower.

DYE

végétale. This trade is unaffected by the aniline imitations of safflower, and constitutes an article of considerable importance. The dry carthamine precipitate is sometimes called India or China lake, and this mixed with finely pulverised talc constitutes *rouge végétale*. (See Carmine; also Carnelian—the coloration of inferior gems)

OIL
670

THE OIL

There are two kinds of seeds, or, to be more accurate, of fruits—one

account of the little heat which it gives out (*Baden Powell*) It is used locally for culinary purposes, and is said to form an ingredient of the

Prices.

"In Bulandshahr the safflower yields about 7 maunds of seed per local *bigha*. The oil-cake is supposed to be the perquisite of the oil-presser in lieu of wages. A maund of seed yields 7 seers of oil, 14 seers of oil-cake, and 19 seers of husk or *bhusa*, and the oil sells at from 4 to 5 seers for the rupee, the cake at 36 seers, and the *bhusa* at 4 maunds" (*E. T. Atkinson*)

"The pure oil is seldom offered for sale. Though it lowers the quality of the oil, the outturn is generally increased by mixing its seeds with gingelly seed" (*Bomb Gas*, 153). Although the oil is apparently not exported from India a considerable trade is done with Liverpool and London in the seeds

EXPRESSION
Dry cold
671

Expression of Oil.—"The oil is expressed in the same manner as the

Dry Hot.
672

Dry Hot extraction of Oil.—"There is also another way of extracting the oil which is, I think, so peculiar that I will attempt to describe it. It

it, in fact, but this sing his well ropes, used for exposure

A hole is dug in the jar or *gurrak* of any capacity, then plate with a hole of about a centre. Above this is placed *dhurra* or *kussum* seed inverted three is luted with clay, and essels,—

Dried The life is kept in ignition for about half an hour, when it is removed. The

C. 672

Process of
extracting
the oil after
the Dry Hot
method.

The Safflower.

ARTHAMUS
tinctorius

upper inverted vessel is found to be about half full of the charred seed, and the lower one, which was imbedded in the ground about one third full of a black sticky oil. B charred, but the natives assert preservation of leathern vessels is worth the while of chemists. This kind of oil would be of any commercial value at home. The yield of oil by this process is more than a fourth larger than by the press" (*R W Bingham Jour Agri Hort Soc, XII, 340*)

OIL

THE MEDICINE.

MEDICINE.

"This plant is the *Kutumbhu* of Sanskrit writers, who describe the seeds as purgative, and mention a medicated oil which is prepared

OIL
673

A fixed oil is prepared from it which the *Vytians* used as an external ulcers, which into an of the dried flowers taken internally cures jaundice (*Hort Jamaica, I, 72*). Loureiro says that the seeds are considered as purgative, or *ecoprotic*, resolvent and emmenagogue. In South America as well as in Jamaica, the flowers are much used for colouring broths and ragouts.

Flowers.
674
Seeds
675

resemble in colour, but from which they may be distinguished by their tubular form, and the yellowish style and filaments which they enclose. In large doses carthamus is said to be laxative, and administered in warm infusion, diaphoretic. It is used in domestic practice as a substitute for saffron in measles, scarlatina and other exanthematous diseases to promote the eruption. An infusion made in the proportion of two drachms to a pint of boiling water is usually employed, and given without restriction as to quantity" (*U S Dispens*)

OIL
677

"The seeds are laxative. The oil is used as a dressing for ulcers" (*Surgeon W Barren, Bhuj, Cutch*)

Decoction
678
FOOD
Seeds
679
Leaves
680

CARUM
Carul.

The Caraway.

681

CARUM, *Linn.*; *Gen. Pl.*, I., 890.Carum Carui, *Linn.*; *Fl. Br. Ind.*, II., 680; UMBELLIFERÆ.CARAWAY; FRUITS OU SEMENCES DE CARVE, *Fr.*; KÜMMEL, *Germ.*Vern.—*Shia jird* (U. O. Dutt), *sira*, HIND; *Jira*, BENG.; *Zira silyah*,References.—*Stewart, Pb Pl.*, 104; *DC. Prodr.*, IV., 115; *Pharm. Ind.*,*Cyclop. Agri.*

Habitat.—A herbaceous plant cultivated, for its seeds, as a cold-season crop on the plains of India and frequently on the hills, as a summer crop, as in Baltistán, Kashmir, and Garwhal, &c, at an altitude of between 9,000 to 12,000 feet. Distributed to Western and Northern Asia and Europe. The Greek and Latin names of the plant are said by some writers to be "derived from Caria, the native country of the plant" (*Bird-*

CONDIMENT.
682

Per-
is now
lists of

Camgium The existence of the name *Isiayati-sira*, that is, European *sira*, should not by itself be viewed as excluding the true Caraway from an

C. 682

The Caraway.

CARUM
Carui.

oriental origin since such a name might simply mean that in that part of the country it was first brought to the attention of the natives by the Europeans. Indeed, the facilities of trade offered by the Persian Gulf can easily be understood to have made the people of Bombay more familiar with an imported article than with a wild or even cultivated plant of the Panjáb Himálaya. Authors are about equally divided in the restriction of the word *sira* to Carum Carui on the one hand, and to Cuminum Cuminum on the other (Conf. with C. nigrum).

Dr Dymock says that Caraways are brought from the Red Sea Ports to Bombay where they are sold at Rs per pound. Dr Stewart alludes to a considerable trade from Afghánistan, Kashmir, and other parts of the Panjáb Himálaya to the plains of India. The imports of Caraway into Great Britain are about 20,000 cwts a year and chiefly from Holland. It is also largely grown in Kent and Essex.

Oil.—A valuable essential oil is obtained from the seeds, called Caraway Oil. This oil is colourless or pale yellow, thin, with a strong odour and flavour of the fruit. It is used in medicine and more extensively as a perfume for soaps (*Spons*).

Perfumery.—Piesse, in his book on perfumery, remarks that the odorous principle obtained from the seeds by distillation, when dissolved in spirit, may be combined with lavender and bergamot for the manufacture of cheap essences in a similar way to cloves.

Medicine.—As a medicine the dried fruit possesses stimulant and

CONDIMENT

TRADE.
683OIL.
684PERFUMERY.
685MEDICINE.
686

water

"Muhammadan writers describe the fruits as aromatic, carminative, and astringent, from them they prepare an eye-ash which is supposed to strengthen the sight, they are also used as a pectoral, and considered diuretic and anthelmintic. A caraway bath is recommended for painful swelling of the womb, and a poultice for painful and protruding piles" (*Dymock's Mat Med W Ind*, 304).

Fruit
687CHEMISTRY
688

absolutely deprived, perfectly pure carvone would no doubt prove no longer to possess the specific odour of the drug. By distilling it over sodium, it acquires a rather pleasant odour, its specific gravity at 15° C is equal to 0.861.

The Bishop's Weed.

CARUM
copticum.

MEDICINE

(*Waring's Bazar Med*) They are administered in flatulence, flatulent colic, atonic dyspepsia, and diarrhœa, and are often recommended for cholera. They are used most frequently in conjunction with asafœtida, myr and an aror native doctors as a stomachic, caridac, and stimulant, and also by the veterinary practitioners in India in the diseases of horses and cows. Dr. Bidie is strongly in favour of the extended use of this medicine. "As a topical remedy it may be used with advantage, along with astringents, in cases of relaxed sore-throats. For disguising the taste of disagreeable drugs and obviating their tendency to cause nausea and griping, I know of no remedy of equal power." The seeds have come into special notice in England and Germany for the manufacture of *Thymol*, enormous quantities of which are now made and used as an antiseptic (*Smith*)

Thymol.
694OMUM.
695

Flatulence and as an antispasmodic in hysterical pains. Of late, it has been extolled as a powerful antiseptic superior to carbolic acid (*Home*)

drunkenness and dipsomania, omum seems worthy of trial" (*Waring's Bazar Med*). Dr. Stocks was the first to draw attention to a crystalline substance sold in the bazars of the Deccan and Sind, known as *Ajwain-ka-phul*. This is prepared from the fruits of *Carum copticum* or forms spontaneously on the surface of the distilled water (*Pharm Ind*)

Chemical Composition.—The authors of the *Pharmacographia* say

CHEMISTRY.

(1856) to be identical with thymol, C_8H_8 , $\left\{ \begin{array}{l} C_8H_8 \\ C_8H_7 \end{array} \right\}$, as contained in

Thymus vulgaris

on, first rectified the oil deposited much or more in to a cold some. We found the somewhat larger fete fusion. On ystallizes when a

CARUM
copticum

The Bishop's Weed.

CHEMISTRY

"C. C, the
the same
however,
ol, either
alcohol,
specific gravity 0.830, and saturated with sulphuretted hydrogen, crystals
of $(C_{10}H_{14}C)_2SH_2$ are at once formed as soon as a little ammonia is
added." (*Pharmacog*)

Special Opinions—§ "Stimulant and laxative. The white variety is
lactagogue" (*Assistant Surgeon Nehal Singh, Saharanpore*) "Have
used it to increase the flow of milk with no decided effect" (*Surgeon
D. Picachy, Purneah.*)

Food—The seed is used parched and powdered, or raw and entire

FOOD
Seed
689
Roots
690

(*Assistant Surgeon Shib Chunder Bhattacharj, Chanda, Central Prov-
inces.*)

691

Carum copticum, *Benth* ; *Fl Br Ind*, II, 682 ; *Wight, Ic*, t 566.

THE BISHOP'S WEED, LOVAGE, AJAVA SEEDS, AMTZAD,
Dutch ; *Sison, Fr.* ; *AMEOS, Port.*

Syn—AMMI COPTICUM, *Boiss* ; *LIGUSTICUM AJAWAIN, Fleming* ; *L.
AJOWAN, Roxb* ; *PTYCHOTIS COPTICA, DC.* ; *P. AJOWAN, DC* ; *SISON
AMMI, Jacq* ; *BUNIUM AROMATICUM, Linn*

Vr--

Ri

Med Hind, 172, 173, 324 ; *Dymock, Mat Med*, v 124, 224, 225, 226,

Prescrip

OIL
692
MEDICINE
693

Black Caraway.

CARUM
Roxburghianum.

these seeds he gave the name *Carum nigrum*, without apparently having either seen the plant or ascertained anything more about them. Stewart seems to have gone into the subject for he reduces Royle's *C. nigrum* to *C. Carui*. In this view he appears to be supported by Mr. C. B. Clarke in the *Flora of British India*, since Royle is by that author quoted as having found the true caraway in Kashmir and Garwal. In what has been already said under *C. Carui* this opinion has been supported, but at the same time it must be added that Dr. Dymock and many other writers continue to allude to a black form of caraway. Dr. Dymock says "*Sajira* or *Siah sirah* (Bomb.) has more slender and darker-coloured fruits than the true caraway, a transverse section shows a similar structure. The flavour approaches that of Cummin, and the Persian name which it bears signifies black cummin. It is probably the article described in Persian works on *Materia Medica* as *Kirmani* or black cummin."

698

aver
into
expo

Under *C. Carui* it has already been stated that a considerable trade is done between the North-Himalayan and trans-Himalayan regions with the plains of India in what has been accepted as the true caraway. These two seeds are distributed all over India, the Europeans using the

forcibly draw attention to the fact that recent writers have, as it would appear, been confusing two very distinct seeds under one botanical name.

It is thus probable that the vernacular names given under *C. Carui* and *C. nigrum* (the former being the name of the true caraway and the latter of the black form) are not the same.

MEDICINE.

699

FOOD.

700

701

Carum Roxburghianum, Benth., *Fl. Br. Ind.*, II, 682, Wight, Ic., t. 567.

Ajmod, *bodajamo* Guj.
References — *Roxb., Fl. Ind.*, Ed. C. B. C., 273, *Dalgl. & Gibs, Bomb., Fl.*, t. 1, p. 100.

Carum, and Ajmod

C. 701

CARUM
nigrum.

Black Caraway.

CHEMISTRY

"Thymol is more conveniently and completely extracted from the oil by shaking it repeatedly with caustic lye, and neutralizing the latter

"The oil of ajwain, from which the thymol has been removed, boils at about 172° , and contains cymene (or cymol) $C_{10}H_{14}$, which, with concentrated sulphuric acid, affords cymen sulphonic acid, $C_{10}H_{13}SO_3OH$. The latter is not very readily crystallizable, but forms crystallized salts with baryum, calcium, zinc, and lead, which are abundantly soluble in water. In the oil of ajwain no constituent of the formula $C_{10}H_{16}$ appears to be present, mixed with alcohol and nitric acid, it at least produces no crystals of terpin

"The residual portions of the oil, from which the cymene has been distilled, contains another substance of the phenol class different from thymol"

Special Opinions.—§ Sometimes used by the natives for colds, useless as far as my experience goes (*Surgeon Major C J McKenna, Cawnpore*) "Much used in flatulence diarrhoea, and with other drugs in dyspepsia. Very useful in flatulence and with dyspepsia, especially administered in powder mixed with other antispasmodics" (*Surgeon G Price, Shahabad*) "contains, and which is in Madras famine relief I don't think it was of a

dr (G B Madras).
used in dyspep-
si (Hospital Assis-
ta stomachic, mixed

with black pepper and salt and taken in empty stomach, relieves flatulence and colic and promotes digestion" (*Assistant Surgeon Shih Chunder Bhattachary, Chanda, Central Provinces*) "The water distilled from the seeds is very useful as a carminative, and is largely used by the natives, being administered to newly born infants as a carminative and stimulant. T

ly used as a
Central Prov
mixtures for i
in powder, an
to newly bor
vinces) "A
Negapatam)
W A Barren Belgaum, Bombay)

Food—The seeds are aromatic, and form an ingredient of the preparation known as *pan*

Carum nigrum, ? Royle, *Him Bot*, 229.

BLACK CARAWAY

Syn—Stewart Baden Bonn 1824

kirmāni, siyah sirah, PERS

References—*Pharm Ind*, 99, Baden Powell, *Pb Prod*, 351, Moodeen Sheriff, *Supp Pharm Ind*, 90, Dymock *Mat Med W Ind*, 305, S. Arjun, *Bomb Drugs*, 63, Birdwood, *Bomb Drugs*, 39

Habitat—Royle mentions that seeds under the name of *Zeera steeh* are imported from Kunawar, and that these are "a kind of caraway" To

Cloves.

CARYOPHYLLUS
aromaticus

593 U C Dutt, *Mat Med Hind*, 184 307, Dymock, *Mat Med*

Habit.

in the 12th year, when the average annual produce may be estimated at 6-7lb of marketable fruit from each tree. There is usually a crop every year, but in Sumatra the trees often bear only twice in 3 years. When past its prime, the tree has a ragged appearance. Its existence in Sumatra is supposed to be limited to a duration of about 20 years, except in very superior soil when trees perhaps last 30 years. At Ambon it does not bear years. F old trees to through it very a winds. The harvesting of the flower buds (cloves) commences immediately they assume a bright red colour. The best and

matting near a slow wood fire, and very rarely they are scalded in hot water before smoking. They are ready for packing when they break easily between the fingers" (*Spon's Encycl*)

OIL.
707

of spirit

Description of the Drug—"The varieties of cloves occurring in commerce do not exhibit any structural differences. Inferior kinds are dis-

CARYOPHYLLUS
aromaticus

Cloves

MEDICINE
702

Habitat—A herbaceous plant extensively cultivated throughout India, from Hindustan and Bengal to Singapore and Ceylon

Medicine—The seeds of this species are useful in hiccup, vomiting, and pain in the bladder. They form an ingredient of carminative and stimulant preparations, and are useful in dyspepsia

Special Opinions—§ 'Carm native. It is an essential ingredient of native cookery and is generally called *Randhuni*' (*Assistant Surgeon Shib Chunder Bhattachary, Chanda Central Provinces*)

Food—Often raised in gardens during the cold season for the seed which is used in flavouring curry, also used by the Europeans as a substitute for parsley (*Royle*). Extensively cultivated in Gajarat (*Lisboa*). Leaves though of an unpleasant smell are now and then used by Europeans as a substitute for parsley (*Voigt*)

FOOD
Seeds
703
Leaves
704

705

Carving, Fancy work, Images, &c—

Timbers used for—

Berberis nepalensis, *Spreng* (useful for inlaying)

Buxas sempervirens *Linn* (carving)

Cedrela Toona, *Roxb* (carving)

Celastrus spinosus, *Royle* (carving and engraving)

Chickrassia tabularis, *Adr Juss* (carving)

Cocos nucifera, *Linn* (fancy work)

Crataeva religiosa, *Forst* (models)

Cupressus torulosa *Don* (images)

Dalbergia cultrata, *Grah* (carving)

D latifolia, *Roxb* (carving and fancy work)

D Sissoo *Roxb* (carved work)

Diospyros Ebenum, *Konig* (used for inlaying)

D melanoxylon, *Roxb* (fancy work and carving)

Euonymus grandiflorus, *Wall* (carving)

E. Hamiltonianus, *Wall* (carving into spoons)

Givotia rottleriformis, *Griff.* (carving figures)

Gmelina arborea, *Roxb* (carving images)

Hardwickia binata, *Roxb* (ornamental work)

Holarrhena antidysenterica, *Wall* (carvings)

Kydia calycina, *Roxb* (carving)

Melia Azadirachta, *Linn* (idols)

Pistacia integerrima, *J L Stewart* (carving, ornamental work)

Premna tomentosa, *Willd* (fancy work)

Santalum album, *Linn* (carving)

Stephegyne parvifolia, *Korth* (carved articles)

Symplocos cratægoides, *Ham* (carving)

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

—

idols)

Wrightia tinctoria, *R Br* (carving)

W. tomentosa, *Rom & Sch* (carved work)

CARYOPHYLLUS, *Linn*, *Gen Pl*, I, 719

706

Caryophyllus aromaticus, *Linn*, *DC Prodr*, III, 262.

CLOVES

[MYRTACEÆ

Syn.—EUGENIA CARYOPHYLLATA, *Thunberg*

Ve

Ve

Ve

Ve

Ve

Ve

Ve

Ve

Ve

Ve

Ve

Ve

Referenc

71 C

Ind,

Ed,

C. 706

m, I,

1 arm

15th

d, I,

Cloves.

CARYOPHYLLUS
aromaticus.

503, U. C. Dutt, *Mat Med Hind*, 1864, 307; *Dymock, Mat. Med*
W. Ind., 2nd Ed., 328, O'Shaughnessy *Rep.*, 1820, 111.
Pl and Drugs of Sind, 193,
Bazar Med., 44, *S. Arjun*,
Pr., 35, *Lisboa, U. Pl of B. Ind.*
34; Spens. Encyclop., 1807.
Treasury of Botany, Ajmir Me

Habitat.—A native of the Moluccas. Cultivated in Southern India. The Dutch tried to restrict its cultivation to the Island of Amboyna, but in the course of time it got introduced into India and other tropical countries. The flower-buds of this plant yield the cloves of commerce.

Cultivation and yield.—The tree naturally selects a volcanic soil, and a sloping position. The Dutch tried to restrict its cultivation to the Island of Amboyna, but in the course of time it got introduced into India and other tropical countries. The flower-buds of this plant yield the cloves of commerce.

The tree naturally selects a volcanic soil, and a sloping position. The Dutch tried to restrict its cultivation to the Island of Amboyna, but in the course of time it got introduced into India and other tropical countries. The flower-buds of this plant yield the cloves of commerce.

When past its prime it has a ragged appearance. Its existence in Sumatra is supposed to be limited to a duration of about 20 years, except in very superior soil, when it may not bear till the 12th-15th years. Hence it is never

winds. Sometimes, however, they are beaten off by long bamboos, and caught in cloths spread below. The plucked cloves undergo a process of drying, which renders them easily for packing when they break easily between the fingers" (*Spens' Encycl.*)

Oil.—Every part of the plant abounds with essential oil. The process is a colourless or a pale yellow. It easily dissolves in alcohol, and is often adulterated with dissolving oil of clove of spirit.

Description of the Drug.—"The varieties of cloves occurring in commerce do not exhibit any structural differences. Inferior kinds are distinguished by being less plump. The cloves met with in the Inc Those suited for medical use"

OIL
707

CARYOPHYLLUS
aromaticus.

Cloves.

DESCRIPTION
OF EHT
DRUG

spicy, pungent taste, and should emit a trace of oil when pressed with the nail (*Waring's Basar Medicines*) "The Americans have introduced into commerce an imitation in a solution of true natives, are largely of mixed spice and cloves or fruits are *Encycl*, 1808)

MEDICINE
Buds
708

Medicine —The dried flower-buds which constitute the cloves of com-

grain pill made of equal parts of jalap and powdered cloves generally opens the bowels "Cloves are much used in Hindu medicine, as an excellent effect in debility, loss of appetite, and in convalescence after fevers. "The oil, *Lavanga-tela*, is used externally in rheumatic pains,

tonic, and digestive qualities They have a curious superstition to the effect that one male clove eaten daily will prevent conception" (*Dymock's Mat Med W Ind*, 329)

Chemical Composition —"Few plants possess any organ so rich in essential oil as the drug under consideration The oil known in pharmacy as *Oleum Caryophylli*, which is the important constituent of cloves, is obtainable to the extent of 16 to 20 per cent. But to extract the whole, the distillation must be long continued, the water being returned to the same material

"The oil is a colourless or yellowish liquid with a powerful odour and taste of cloves, sp gr 1.046 to 1.058 It is a mixture of a hydrocarbon and an oxygenated oil called *Eugenol*, in variable proportions The for-

ceous odour.

of eugenol is given by the formula $C_6H_5\left\{\begin{smallmatrix} OH \\ CH \end{smallmatrix}\right\}CH_2$. It belongs

CARYOTA
urens

TRADE

Sago Palm

Imports for 1884 85

Pres dency to wh ch imported	Quant ty	Value	Country from wh ch imported	Quantity	Value
	lb	R		lb	R
Bombay	4 598 4 9	10 50 680	Zanz bar	4 776 842	11 05 877
Bengal	190 526	58 283	Aden	11 767	2 908
Br tsh Burma	1 283	425	Other Countries	2 397	1 056
Mad as	773	453			
TOTAL	4 791 006	11 09 841	TOTAL	4 791 006	11 09 841

Exports for 1884 85

P es dency from wh ch expo ted	Quant ty	Value	Country to wh ch expo ted	Quant ty	Value
	lb	R		lb	R
Bombay	1 618 465	3 55 692	Un ted K ngdom	1 112 224	2 32 739
Bengal	29 65	10 090	Ch na—Hongkong	349 698	84 966
Madras	1 390	1 462	Stra ts	124 01	33 543
S nd	20	5	Turkey in Asia	15 137	3 887
			Aden	7 000	1 790
			F ance	7 000	750
			Other Count es	33 830	8 574
TOTAL	1 649 040	3 67 249	TOTAL	1 649 040	3 67 249

Very little can be said regarding the present position of the new industry of cultivating cloves in South India. Good samples were, however shown at the Colonial and Indian Exhibition.

CARYOPTERIS, Bunge Gen Pl, II 1157

Caryopteris Wallichiana, Schauer DC Prodr XI 625;

[VERBENACEÆ

Vern.—*Moni mohani* KUMAON Shechin NEPAL Malet LEPCHA

References.—*Brand s For Fl* 370 *Gamble Man Timb* 299

Habitat.—A large shrub with thin grey papery bark peeling off in vertical strips met with on the outer Himalaya from the Indus to Bhutan ascending to 3000 feet

Structure of the Wood.—Dark grey, moderately hard with the scent of cherry wood

CARYOTA, Linn Gen Pl III 918

Caryota urens, Linn Gamble Man Timb 420 PALMÆ

KNOWN IN BOMBAY AS THE HILL PALM also SAGO PALM

Vern.—*Mar* HIND *Runghong s mong* LEPCHA *Bara flawar* ASS
Salopa URIYA *Mari ka jhar* DEC *Bherawa beri* bh ris mahad beri

CARYOTA
URENS
711

Sago Palm.

CARYOTA
urens.

Handwritten text in Devanagari script, likely a title or description of the plant.

Handwritten text in Devanagari script, likely a description of the plant's characteristics or uses.

Handwritten text in Devanagari script, likely a reference or source.

References.—Roxb., *Fl. Ind.*, Ed. C.B.C., 668; Brandis, *For. Fl.*, 550, *Kurz, For. Fl. Burm.*, 11., 530; Voigt, *Hort. Sub. Cal.*, 637; Thwaites, *En. Ceylon Fl.*, 329; Dalz. & Gibs., *Bomb. Fl.*, 278; *Pharm. Ind.*, 248;

Habitat.—A beautiful palm, with smooth, annulated stem, met with in the forests of the western and eastern moist zones. On the Western Ghats, it extends to near Mahabaleshwar. In the Settlement Reports of the Chanda district it is stated that this palm abounds in the south-eastern corner of Aheree, and might with advantage be extended to all parts of the district, for it thrives well wherever it is planted. It is common in Burma, Bengal, and Orissa, ascending in Sikkim to 5,000 feet.

Fibre.—"The leaves give the *Kittul Fibre*, which is very strong and is made into ropes, brushes, brooms, baskets, and other articles; the fibre from the sheathing petiole is made into ropes and fishing-lines" (*Gamble*), and is said to be suitable for paper manufacture.

At the Colonial and Indian Exhibition (1886-87) much interest was taken in *salapa* fibre sent from Orissa, Burma, and Kolaba in Bombay. A corset manufacturer applied at the office of the Indian section for a fibre which might take the place of whalebone in corset-making. He was shown the *salapa* (*kittul*) fibre and also the similar cord like fibres from the interior of the stems of the cocoanut and palmyra palms. It

FIBRE.
712

Ceylon. At the Colonial and Indian Exhibition he pointed out to the writer a sample of the much inferior *kittul* like fibre from *Arenga saccharifera* (see A. 1336) as the *kittul* he had formerly seen as sent from India. He admitted that the sample of *salapa* shown him at the Exhibition was

C. 712

CARYOTA
urens

Sago Palm

as good as any he had ever seen from Ceylon, and seemed confident a large trade could be done in the Indian fibre

It is commonly reported that in Ceylon the black fibre from the leaf-stalks is manufactured into ropes which are of great strength and

Tomentum
stem fibres

palis is employed as a bow string or as a fishing line (see 1067)
(*Royle Fib Pl*)

MEDICINE
713

Medicine — "An excellent spirit is obtained by the fermentation and distillation of the toddy obtained from this elegant palm, which is not uncommon on the west coast of the Madras peninsula. It is well adapted for pharmaceutical purposes. 'A glass of the freshly drawn toddy, taken early in the morning, acts as a laxative' (*Pharm of India*). 'The nut is used as an application to the head in cases of hemiplegia, from an idea of the supposed efficiency of the half nut in curing the affected half of the head' (*S Arjun, Bombay Drugs*)

FOOD
714

Food — Roxburgh writes "This tree is highly valuable to the natives of the countries where it grows in plenty. It yields them, during the hot season, an immense quantity of toddy or palm wine. I have been informed that the best trees will yield at the rate of 100 pints in the 24 hours. The sap in some cases continues to flow for about a month. When fresh, the toddy is a pleasant drink, but it soon ferments and when distilled becomes arrack, the gin of India. The sugar called jagery is obtained by boiling the toddy. The pith or farinaceous part of the trunk of old trees is said to be equal to the best sago, the natives make it into bread, and boil it into thick gruel, these form a great part of the food of these people and during the late famine (1830?), they I have reason to believe this substance eaten the gruel and think it fully get from the Malay countries

(111111)

'The trees are tapped when they are from fifteen to twenty five years old. Besides bruising and binding it, the spathe, which is called *kote*, is heated to make the juice flow. Every three or four days a white

TIMBER
715

value of the juice the big trunked palm differs little from the palmyra. Since 1879 when the tree tax was raised from 1s 6d to 6s (*annas 12 to Rs 3*), the number of trees tapped has greatly fallen" (*Bomb Gas* (Kolaba), XI, p 30)

and durable
wood
conduits,
"Is in

General use for field tools (*Bomb Gas*, XV, 1, 65)

716

Cascarilla bark, the bark of *Croton Eluteria*, EUPHORBIACEÆ

A native of the Bahamas. The bark is imported into India

C. 716

Casearia

CASEARIA
tomentosa.

CASEARIA, Jacq., Gen Pl, I, 796

Casearia esculenta, Roxb, Fl Br Ind, II, 592, SAMYDACEÆ

717

Syn — *C. LÆVIGATA*, Dals, in Hooker's Jour Bot, IV, 107, *C. CHAM*
PIONII and *C. ZEYLANICA* ThwaitesVern — *Amunda jungura*, TEL, *Wal wareka*, SINGReferences — *Roxb, Fl Ind, Ed C B C, 377, Drury, U Pl 119, Dals*
& Gibs, Bomb Fl, 11, Thwaites, En Ceylon Pl, 19

Habitat —

Coorg, comr
to SingaporeMedicine — "The roots are purgative, and as such used by the hill
people" (*Roxb*)Food — "The leaves are eaten in stews by the natives" (*Roxb*)

MEDICINE

718

FOOD

719

720

C. glomerata, Roxb, Fl Br Ind, II, 591

Vern — *Lurjur*, SYLHET, *Burgonli*, NEPAL, *Sugvat*, LEPCHAReferences — *Roxb, Fl Ind, Ed C B C, 376, Kurr, I, 530, Gamble,*
*Man Timb, 205*Habitat — A shrub or (in the interior of Sikkim) a tree 20 to 30 feet
in height Frequent in Bhutan and on the Khasia Hills at an altitude
of 3,000 feetStructure of the Wood — Yellowish white, moderately hard, rough,
weighing between 45 and 48 lb per cubic foot Used for building,
charcoal, and occasionally for tea boxes

TIMBER.

721

C. graveolens, Dals, Fl Br Ind, II, 592

722

Vern — *Chilla, naro, aloal, kathera, pimprī*, HIND, *Rari*, KOL, *Beri,*
Kharwar, Newri, SANTAL, *Girchi, tundri*, GOND, *Rewat*, KURKU,
Moda, MARReferences — *Brandis, For Fl, 243, Gamble, Man Timb, 205, Dals &*
*Gibs, Bomb Fl, 11, Lisboa, U Pl of Bomb, 81 and 265*Habitat — A shrub or small tree, 20 feet in height, found in Garhwāl
and Kumaon, Sikkim at an altitude of 1,500 feet, Deccan Peninsula and
in Burmagh, weight
purpose
sion of the

TIMBER

723

DOMESTIC

724

C. tomentosa, Roxb, Fl Br Ind, II, 593, Wight, Ic, t 1849.

725

Syn — *C. ANAN*

Vt

72, 243
Lisboa,
es, En

Ceylon Pl, 19

Habitat — A shrub or small tree, attaining a height of 25 feet, common
throughout India and CeylonMedicine — The bark is bitter and used as an adulterant for the
(*Mallotus philippinesis* or) *Kamela* powder "The pounded fruit yields a

MEDICINE

726

CASSIA
Absus

Senna

MEDICINE

milky, acrid juice employed to poison fish" (*Brandis*) The leaves are used in medicated baths and the pulp of the fruit is a very useful diuretic (*Lindley*)

Special Opinion — § "Bark applied externally in dropsy" (*Rev A Campbell Santal Mission, Bengal*)

TIMBER
727

Structure of the Wood — Yellowish white, moderately hard, rough, close-grained, weight 41 lb per cubic foot, used to make combs.

Cashew-nut. See *Anacardium occidentale*, *Linn*, ANACARDIACEÆ.

Cassareep, and

Cassava Bread, and Tapioca, see *Mamhot utilitissima*, *Pohl*, EUPHORBIACEÆ

CASSIA, *Linn*; *Gen Pl*, I, 571

The word Cassia is taken from the Latin and the Greek *Kassia*, and from this has been derived *CASSIA* the Italian, and *CASSE*, the French. In the Scriptures two or three different things appear all to be rendered as Cassia. The genus is of considerable importance from a medical point of view.

728

Cassia Absus, *Linn.*, *Fl Br Ind* II, 265

Vern — *Tashmisaj* chashmisaj *PERS*,
chashum cheshmak *PERS*,
Mulaspal-virai karunka nam
vittulu *TEL* Karin kolla *M*
Chimar or chimr, chinól, *GUJ*
SING

References — *Roxb Fl Ind Ed C B C* 351 *Gamble Man Timb*,
136 *Thwaites, En Ceylon Pl* 96 *Stewart Pb Pl* 61 *Aitchison*,
Cat Pb Pl 51 *Pharm Ind* 78 *Moodeen Sher ff Supp Pharm*
Ind 92 *Dymock*
Beng Dispens 30
account of Sind B
Drugs 45 *Drury*
Him Dist 731
Treasury of Botany 232

Habitat — An erect annual 1-2 feet high having grey, bristly, viscose hairs. Found growing at the foot of the Western Himalaya and from thence distributed to Ceylon.

History — The seeds of this plant were used by the ancient Egyptians in the treatment of ophthalmia, and through them the Roman and the Greek, and from the latter the Muhammadan writers became aware of their properties. Dioscorides speaks of them under the name of *Akakalis*. Their

MEDICINE
Seeds.
729

trial to this treatment, and the results were on the whole confirmatory of its alleged efficacy. Dr. G Smith, Superintendent of the Eye Infirmary at Madras, in his report, characterises it as a dangerous

Alexandrian Senna of Commerce

CASSIA
alata.

MEDICINE.

Extract,
730

application in catarrhal ophthalmia and granular lids, adding that its application causes great pain. As met with in the bazárs, these seeds are of a black, shining colour, somewhat flat, of an oval or oblong form, pointed at one extremity, about one-sixth of an inch long, having a bitter taste" (*Pharm Ind*). They are very bitter, somewhat aromatic and mucilaginous, and, as such, have been found very useful in mucous disorders. An extract is prepared from them and used to purify the blood. Dr. Irvine, in his *Materia Medica of Patna*, says that the receptacle of the seed possesses stimulant and diuretic properties (dose 5 grains to 1 scruple). According to some authors, a plaster made from the seeds is a useful application to wounds and sores, especially of the penis.

Special Opinions — § "Seeds are found efficacious in ring-worm" (*Surgeon C F W Meadows, Burrisal*). "Cathartic, dose $\frac{1}{2}$ to 3 drachms, used in habitual constipation, or in constipation caused by pregnancy, with confection of rose and liquorice, have proved effective. In dyspepsia, flatulent colic, and bilious headache, it is given as a compound powder, containing ginger, black rock-salt, amla and bury, and chotty hus" (*Hospital Assistant Abdulla, Civil Dispensary, Fubbulpore*).

According to Dr. Dymock, the Bombay supply comes from Sind and Cutch, value, R4 a Surat maund of 37 $\frac{1}{2}$ lb.

Cassia acutifolia, Delile

European
Senna
731

THE ALEXANDRIAN SENNA of Commerce.

Syn.—C SENNA, β Linn. C LANCEOLATA, Nestoux, non Forsk nec. H & A, C LENITIVA Bisch. SENNA ACUTIFOLIA, Batka. See also the remarks under C LANCEOLATA, Forskhal.

Habitat — A native of Nubia (at Sukkot, Mahas, Dongola, Berber), of Kordofan and Sennaar, and other parts of Africa.

For Indian Senna see C. angustifolia, C. Burmanni, and C. obovata.

C. alata, Linn., Fl. Br. Ind., II., 264.

732

Vern.—*...*

TAN
Bark.
733

Dyes and Tans. The numerous samples of this bark, shown at the late Colonial and Indian Exhibition, were highly commended by the tanners.

MEDICINE
Leaves.
734

Indian or Tinnevely Senna.

CASSIA
angustifolia.

shóna makhi, MAR , Nattu nila virai nila virai, nila-vakai TAM , Nela
tangédu TEL , Nila vaka MALA , Nelavarike KAN , Sa ta kola nild
vari, nela-vari SING , Puve kain yoe, BURM

References — D H I A S E S n Pl D n H nht S A

many parts of India. The *Flora of British India* says *C. angustifolia* 'has no claim to be considered indigenous to India'. *C. lanceolata*, Forsk., is a native of Arabia. It seems probable that the mistake made by Dr Brandis gave origin to the statement (see *Pharmacographia*, also *Bentley and Trimen, Med. Pl.*) that *C. angustifolia* is indigenous to Sind and the Panjáb.

The cultivated plant as met with in India is the Tinnevely Senna of commerce and the uncultivated the Bombay Senna or *Senna Mekki* or *Sana miki* *Sona maki* of the East. The last mentioned is imported into India from Arabia. In Bombay it is cultivated at Poona to supply the requirements of Government Hospitals and not as an article of commerce. Stocks say it is grown in Sind.

Botanic Diagnosis — This species is closely related to the preceding, but the leaflets are usually 5-8 jugate, are narrower, being oval, lanceolate, tapering from the middle towards the apex, they are longer, often nearly 2 inches long, and are either quite glabrous or furnished with a very scanty pubescence. The legume is narrower (7-8 lines broad), with the base of the style distinctly prominent on its upper edge.

Description of the Drug — This plant thus affords two of the commercial forms of senna —

1st TINNEVELLY SENNA — This is the leaf obtained from the plant carefully cultivated in South India and (at Poona) in Bombay. Owing to greater care in its collection, Tinnevely senna is of better quality than the Arabian article. The leaves are also larger, being 1-2 inches long of

Tinnevely.
738

Dr Dymock says that large quantities of Tinnevely senna are now sent to Bombay and that so successfully does this Indian article compete in the market, that the importation of Arabian senna is rapidly declining, Tinnevely senna being exported to Europe in its place.

Arabian.
739

CASSIA
angustifolia

Arabian Senna

MEDICINE
Leaves
740

Medicine — Senna was first made known by the Arabs in the ninth century it is extensively employed as a simple and active purgative. The Alexandrian is generally regarded as more powerful than Tinnevely and the Arabian or Moka much inferior to either of these. The object

of potash, tartrate of potash or sulphate of magnesium along with an
Dr
"and
at the

Dr Waring (*Bazar Medicines*) says "The imported senna met with in

CHEMISTRY

decoction for fevers and also to cattle

Chemical Composition — The purgative property is considerably increased by combination with bitters. This fact has been confirmed by many observers. The purgative properties are due essentially to a glucoside acid named *Cathartic Acid*. This which is almost insoluble in ether or chloroform. In senna and magnesium and in this form insoluble in alcohol. The objective decoction, although the senna yields rapidly one or minutes after partaking the by being reddened on the addition of ammonia. Senna taken by wet nurses with equal rapidity influences the milk, purging the sucking infant. If injected into the blood senna acts as a cathartic.

For further particulars see 'Alexandrian Senna' under *C. acutifolia*, and for Senna substitutes see *C. obovata*

purchased at one anna a lb" (*Surgeon Major W. Dymock, Bombay*)

Powdered leaves are used in secondary syphilis" (*Surgeon Major F. L. Ratton M.D. Salem*) "Senna leaves are always purchased in the bazars and esteemed for their cathartic properties" (*A Surgeon*) "An efficient purgative commonly taken by the natives as a cold infusion, causes griping and abundant flow of mucus" (*Assistant Surgeon Shib Chunder Bhattacharya, Chanda Central Provinces*) "Not much used in these days" (*Brigade Surgeon S. M. Shircore, Moorshedabad*)

C. 740

Tanner's Cassia	CASSIA auriculata
Cassia auriculata, Linn, Fl Br Ind, II, 263	741
THE TANNER'S CASSIA	
Syn — SENNA AURICULATA, Roth	
Vet — " " " " " " " " " "	
Ref — " " " " " " " " " "	
" " " " " " " " " "	
<i>Exhib</i>	
Habitat — A tall shrub with the virgate branches and under side of the leaves finely grey downy. Wild in the Central Provinces, the Western Peninsula South India, and Ceylon, often planted elsewhere.	
Gum — It is said in <i>Spons' Encyclopaedia</i> to yield a medicinal resin, very scarce, but Dr Dymock informs the writer he has never seen this supposed resin although he has frequently handled the bark. In Bengal a brownish sap hardens on the surface of wounds on the bark, this may be the so-called resin.	GUM 742
Tan and Dye — The bark is one of the most valuable of Indian tans, and is also, like myrabolans, used to modify dyes. It is said to give a buff colour to leather. Bidie remarks that "when the Government Tannery existed at Hunsur, this bark was used almost exclusively for tanning purposes." This bark was highly commended by the Tanners who attended the conference on tanning materials held at the Colonial and Indian Exhibition in London. It was regarded as a little too dark-coloured but the leather shown as tanned by it was admired. It was recommended that an effort should be made to have an extract prepared from this bark for export to Europe similar to Cutch. Mr Wardle in his recent report says "The bark does not produce much dye, only light	DYE & TAN Bark 743
matter, apparently not used economically	
§ "Skins of animals are tanned by soaking them in water in which the bark of this shrub has been infused for several days" (<i>Honorary Surgeon P Kinsley, Chicacole, Ganjam</i>)	Flowers 744
Fibre — Specimens of the bark were sent to the Calcutta Exhibition	
	FIBRE 745
this plant (Koxo)	

CASSIA
Burmanni.

The Tanner's Cassia.

MEDICINE

Seeds

746

Medicine.—“The SEEDS of this common Indian plant, like those of

Bark

747

obtusely pointed at one extremity, and varying in colour from brown to dull olive-green they are tasteless and inodorous. The BARK is highly astringent, and Dr Kirkpatrick states (*op. cit.*, No 475) that he has employed it in the place of oak bark for gargles, enemata, &c., and found it a perfect substitute for the imported article. Both the seeds and bark appear worthy of further trials. A spirituous liquor is prepared in some parts of India by adding the bruised bark to a solution of molasses, and allowing the mixture to ferment” (*Waring, Pharm Ind.*, pp 78, 79)

Leaves

748

A decoction or infusion of the LEAVES of this plant is much esteemed as a cooling medicine by the Singhalese, and also as a substitute for tea (*Thwaites Murray*). Ainslie says that the Vytians reckon the

Plant

749

into the eyes

Special Opinions.—§ “Bark substituted for oak-bark. Seeds powdered a good local application for ophthalmia” (*Apothecary Thomas Ward, Madanapalle, Cuddapah*). “Antiscorbutic, antibilious, *trifala*, which is made up of dry *awala*, *gall*, and *hirada*, is used as a diuretic and also as an expectorant” (*Surgeon W Barren, Bhuj, Cutch*). “The whole plant, or any part of it, is used in diuresis and diabetes with fair results. The decoction of the flower-buds is an agreeable form in which it is taken in

Flower-buds

750

FOOD

Leaves,

751

Bangalore)

Food.—The leaves are eaten as a green vegetable in times of famine (*Lisboi*).

Domestic Uses.—The branches are largely used by natives as tooth-brushes, and are esteemed as preferable to those of any other plant for this purpose. The root is of great use to workers in iron for tempering the metal (*Ainslie*).

DOMESTIC
Tooth brushes

752

Root

753

754

Cassia Burmannii, *Wight* (in *Madras Jour.*, VI, t 5)

Vern.—The same as those of *C. angustifolia*, *Vahl*

References.—*Brand*, *F. & C. V. M. T. I. & D. L. B.*

Habitat.—A glabrous, shrubby plant, 14 feet in height, often procumbent, pod much curved into a kidney-shape, with a crest in the middle of the valve opposite each seed, leaflets 4-8 pairs. Frequent in the Panjáb (Salt Range, ascending to 2,500 feet, where it is known as *sanna*) and Trans-Indus (where it is called *jñan*), according to *Brandis*; it

C. 754

CASSIA
Fistula.

The Purging Cassia.

GUM.
757

Exhibition from Travancore.

DYE AND TAN
Bark
758

Dye and Tan—The bark is used in tanning, chiefly along with *Terminalia*. Dr. McOann reports that in the district of Lohárdagá, in Bengal, a light-red dye is obtained from the bark, with alum as a mordant, 2 chittacks of bark with 2 tolas of alum being boiled together. The colour is deepened by the use of pomegranate rind. Mr. Wardle reports that the bark contains only a very small quantity of colouring matter. It yielded yellowish drab with tussar silk, light fawn with corah and eri silks, and light yellow-brown with wool. The wood ash is used as a mordant in dyeing. In Dacca and in Cuttack the bark is used as a tan. McOann describes the process of tanning as follows: "Skins, after being treated with lime and cleaned, are soaked in the water, and then are pounded; *tomentosa*, for 24 hours (now Sir E. Bijnor). Experiments were tried at the Government factory, the result being that *amaltas* bark was pronounced a very valuable tanning material. The North-Western Provinces do a small trade in exporting the *amaltás* bark."

MEDICINE
Pulp
759
Root bark
760Flowers
761
Bark.
762
Leaves
763
Root
764

CASSIA
obovata.

Country or Italian and Jamaica Senna.

772

Cassia lanceolata, Forskhal

This species is, by the majority of authors, viewed as quite distinct from either *C. acutifolia* or *C. angustifolia*. It is a native of Arabia, and doubtless to a certain extent is used as a substitute or adulterant for the Mecca senna. It differs chiefly from *C. acutifolia* in having glandular petiolets, the plants are, however, very nearly allied, and, as Forskhal's description is anterior to Delile's account of *C. acutifolia*, both might be reduced to one, which in that case would have to receive the name *C. lanceolata, Forskhal*. Most Indian authors give *C. lanceolata, Forskhal*, but in the writer's opinion incorrectly, as a synonym for *C. angustifolia, Vahl*.

C. Lignea See *Cinnamomum Tamala, Nees*, LAURINEÆ.

773

C. marginata, Roxb, Fl Br Ind, II, 262, Wight, Ill, t 83Syn — *C. ROXBURGHII DC*Vern — *Urimidi uskiamen, TEL, Ngoomce, BURM, Ratoo-waa SING*References — *Roxb Fl Ind Ed C B C, 350 DC Prod II 499, W & A Prod 286 Gamble, Man Timb, 137, Thwaites, En Ceylon Pl, 95 Bedd, Fl Sylt, t 180*

Habitat — A small deciduous tree, with deeply cracked brown bark, found in the Western Peninsula, and in Madras Ceylon and Burma (Thoungyeen forests)

Structure of the Wood — Heartwood light brown very hard. The wood is well adapted for turning, naves of wheels, and handles of tools

TIMBER

774

775

C. mimosoides, Linn, Fl Br Ind, II, 266Vern — *Patwa ghas, SANTAL*

Habitat — Grows on the Himalaya, ascending 5000 to 6000 feet in Kumaon, and on the hills of Bengal and of the Khasia, to Ceylon and Malacca

Medicine — “Root given for spasms in the stomach (*Rev A Campbell, Santal Mission, Pachamba*)”

MEDICINE

Root

776

777

C. nodosa, Ham, Fl Br Ind, II, 261Vern — *Gnu-theing, BURM*References — *Mason's Burm, 404 770*

Habitat. — A common species in the Eastern Himalaya, Manipur, and Burma

It has the properties assigned to most of the wild species

778

C. obovata, Colladon, Fl Br Ind, II, 264; Wight, Ic, t 575Syn — *CASSIA SENNA LINN, SENNA OBTUSA, Roxb*

Known in India as COUNTRY SENNA, and as ITALIAN, TRIPOLI, and JAMAICA SENNA, from its being one of the first species made known to Europe, it was cultivated in Italy during the 16th century

Vern — *Bhuu Tarwar, BOMB*

References — *Roxb Fl Ind (Fl C B C) 352 W and A Prod 288 Moodeen Sheriff Supp Pharm Ind, 94 in part Flück and Handb. Pharmacog 218, Bentley and Trim, Med Pl 189 U S Dispens 1296 Ainslie Mat Med II, 249, Treasury of Botany Dymock Mat Med W Ind, 263*

Habitat — The Western Peninsula, Mysore and South India, especially the Coromandel coast. A small shrub, with the leaves smaller (leaf

C. 778

Negro Coffee.

CASSIA
occidentalis.

lets 3-6 pairs) than in *C Burmannii*, and the pods not near so prominently tubercled over the seeds as in that species

The writer is by no means certain that he is correct in regarding the

MEDICINE
Leaves

779

drian Senna being used as an adulteration in the commercial article This habit has now for some time been discontinued, as also the cultivation of the plant (Conf with *C Burmannii*)

Cassia occidentalis, Linn , Fl Br Ind , II , 262

THE NEGRO COFFEE

Vern — *Kasóndi* bar, *kasóndi* or *kasunda* HIND and DUK , *Hikal*, BOMB , *Kasamara* SANS *Kalkashunda*, BENG , *Nattam takarai*,

780

Habitat — A diffuse, sub-glabrous under shrub, scattered from the Himalaya to the Western Peninsula, Bengal, South India, and Burma to Ceylon Probably introduced Distribution cosmopolitan in the tropics

Medicine — The LEAVES, ROOTS, and SEEDS are used medicinally, and by Hindu and Muhammadan writers they are supposed to have the same properties as *C Senega* The

MEDICINE
Leaves

781

Root

782

Seed

783

sion of
In the f
are emp
the form
the root

various parts of the plant is said to destroy the purgative principle in the seeds and make them taste like coffee The whole plant is purgative Dose of the leaves about 90 grains" (*Dr. Dymock Mat Med W Ind*)

"In the West Indies the ROOT is considered diuretic, and the leaves, taken internally and applied externally, are given in cases of itch and other cutaneous diseases, both to men and animals The negroes apply a plaster The root is of the stomach, and in

has analysed the seeds

CHEMISTRY

CASSIA
occidentalis

Negro Coffee

MEDICINE.

previously treated with ether, by means of alcohol of 60 per cent, the alcohol is distilled off, the syrupy residue treated with absolute alcohol,

of various bodies) It is soluble also in weak alcohol, and in acids and alkalies. The colour cannot be fixed upon tissues by any known mordant. This circumstance induced the author to term it *achrosine*, or 'not colouring,' although being coloured itself.

Special Opinions—§ "Leaves pounded and made into a paste are applied to fresh wounds to bring on their healing by first intention" (*Assistant Surgeon Anund Chunder Mukarji, Noakhali*). "The mature" (*Surgeon J. H.* the treatment

FOOD
Seeds
784
NEGRO
COFFEE

the early part of the year a sample of an article imported at the port of Liverpool from Bathurst, River Gambia, under the above name. They were identified at Kew as the seeds of *Cassia occidentalis*. According to Livingstone, these are used under the name of '*Fed-goso* seeds' on the Zambesi as a substitute for coffee. Monteiro, however, states in his '*Angola and the River Congo*' (*Vol II, p 249*) that *Fedegosa* seeds are used only medicinally as a substitute for quinine. The seeds are roasted and ground, and their infusion taken either alone or generally mixed with coffee" (*1877, p 39*).

"These seeds occasionally find their way into the European market. The following extract from a letter from Dr. Nichols of Dombea dated

native plant as coffee, but it is only lately that I have enquired into the

for our good coffee. Afterwards some of the seeds roasted and ground were brought to me, and the aroma was equal to that of the coffee ordinarily used in the island.

"I intend to send you a good quantity of the '*café marron*' in its stages of preparation, in order that you may have an opportunity of undergoing my experience, and afterwards, you will I think be willing to raise *Cassia occidentalis* above the rank of a weed. I may inform you that the plant itself is used by the native 'doctors' medicinally in the

C. 784

Kasondi Senna

CASSIA
Sophora.

FOOD

form of a de-
I will enquir
report the re
to the sugar
in large quantities" (1881, pp 34-35)

Cassia Oil. See *Cinnamomum zeylanicum*.

C. siamea, Lamk, *Fl Br Ind*, II, 264

Syn —C FLORIDA, Vahl SENNA SUMATRANA, Roxb

Vern —Assod, BOMB, Beati manye konne, TAM, Sime tangadi, KAN,
Waa, SING, Maisalee, BURM

Re — — — — —

785

Habitat —A moderate-sized tree, with smooth bark, found in South India, Burma, and Ceylon Distributed to the Malayan Peninsula and Siam

Structure of the Wood —Sapwood whitish, rather large Heartwood

TIMBER
786

C. Sophora, Linn, *Fl Br Ind*, II, 262

Syn —SENNA SOPHERA and S ESCULENTA, Roxb; C CHINENSIS, Jacq,
SENNA PURPUREA, Roxb

Vern.—Banar kasunda bas ki kasondi, HIND, Kal kashundā, BENG,
Sari-kasondi, jangli takla, DUK, Kuwadice, GUJ, Ran tankala,
MAR, Ponnā-virai periya takarai, pēra-virai TAM, Taidi tangēdu,
nuti kashindha, kasa mardhakamu, tagara chettu, TEL, Ponnam-
takara, MALA, Kasamarda, SANS, Ūru tora, SINGH.

References — — — — —

95, Dals

Supp Pha

Mat Mea

Raw Proa

Pl 343

Balfour, Cyclop

787

Habitat —A closely allied species to C occidentalis, from which it

is said to Ceylon and Penang

Medicine —The BARK, LEAVES, and SEEDS are used as a cathartic, and the JUICE of the leaves is viewed as a specific in ring worm, specially when made into a plaster in combination with sandal-wood A paste made from the root is sometimes used instead of the juice of the leaves The powdered seed is used for the same purpose and also for itch The Sanskrit name means "destroyer of cough, it is supposed by Hindus to have expectoran for snake-b in the forr given in a and leave Top of Dalsu,

MEDICINE

Bark

788

Leaves

789

Seeds

790

Juice

791

C. 791

CASSIA
Tora.

The Fœtid Cassia

CHEMISTRY

Chemical Composition—"This plant, like several others of the same genus, owes its medicinal activity to the presence of chrysophanic acid, sometimes called Rhein, form $C_{14}H_6O_2$ (OH_2). This substance belongs to the anthracene group of carbon compounds, and, like alizarin, is regarded as dioxyan thraquinone, $C_{14}H_6O_2 \{ \begin{smallmatrix} OH \\ OH \end{smallmatrix} \}$. It crystallizes in six sided prisms, is tasteless, and may be sublimed without decomposition, it is contained in Goa powder (50 per cent) rhubarb, most varieties of dock, *Lichen orcella*, *Permelia parietina*, *Cassia alata*, *C. occidentalis*, *C. Tora*,

and Vaseline, dissolve readily containing 52 per cent the fixed oils, a considerable amounts direct from Araroba ice, yielding the acid after re

FOOD
Leaves

792

793

TIMBER

794

795

TIMBER

796

797

Is any, Cassia, 'murtia'

Food—The leaves are eaten by men and animals" (*Atkinson*)
The disagreeable smell is removed by boiling

Cassia, sp (?)

Major Ford sent from the Andaman Islands, in 1866, a sample of a hard, durable wood, olive brown, with a structure very similar to that of *Ougeinia dalbergioides*. Evidently some common Andaman wood, and known by the name of *Gnugyi* (*Gamble, Man Timb*)

C. timoriensis, DC, Fl Br Ind, II, 265

Vern—*Arremene*, SING, *Toung maisalee*, BURM

References—*Kurz, For Fl Burm*, 393, *Gamble, Man Timb*, 138, *Thwaites, En Ceylon Pl*, 96

Habitat—A handsome, small, evergreen tree, met with in Burma and Ceylon

Structure of the Wood—Dark brown, nearly black, resembling that of *C. slamea*, used in Ceylon for building and furniture

C. Tora, Linn, Fl Br Ind, II, 263

THE FŒTID CASSIA

Vern—*Chakund* *panevar*, HIND and BENG *Chak oda arak*, SANTAL *Pawar, panwar, pawas, chakunda*, PA, *Panwar*, N W P, *Takdd*

C. 797

The Fœtid Cassia

CASSIA
Tora.

tarota takla, tankla MAR Kanario, korariya, Guj, Tankala, koraria,

Re

and Tans of Beng, 121, 141 Lisbon, O 11 of Domo, 153, 190, 243,
291; Balfour, Cyclop, Wardle, Report on Dyes & Tans of India

Habitat —A gregarious annual under shrub, from 1 to 2 feet in height, found everywhere in Bengal, and widely spread and abundant throughout the tropical parts of India

Dye —Baden Powell, Atkinson, and other writers say that the seeds

DYE
Seeds
798

a species of *Rhamnus*,
The use of Cassia seeds
! chemical examination.

Mr. Wardle, while examining the dyes of India, had occasion to try the seeds of this plant, and found that they afforded a most useful yellow dye suitable for tasar silk. Mr. Wardle does not appear to have investigated the question of their special property, if any, of being used along with indigo, but from his results it is natural to infer that they would produce a green shade with indigo instead of assisting the blue

CASSYTHA
filiformis.

The Fœtid Cassia; Akaswel.

MEDICINE
Leaves.
799
Seeds.
800

Medicine — The leaves are used as an anæsthetic, both leaves and

Root
801rubbed on a stone with lime-juice, the *Vytians* suppose to be one of the"leaves of a Cassia shrub common in
in dhobie's uth" (*Deputy Surgeon-*FOOD
Seeds.
802
Coffee
substitute.
803
Leaves.
804pot-herb, both leaves and fruit (*Campbell*).§ "The seeds are said to yield a decoction which is reported to be in every respect as good as coffee" (*Mr. C. D. Hardinge, Rangoon*) "A kind of coffee is made from this in Arracan" (*Prof. Romanis, Rangoon*).Cassia, see *Rhus nigra*.CASSYTHA, *Linn* ; *Gen. Pl.*, III, 164.

805

Cassytha filiformis, *Linn* ; *Fl. Br. Ind*, V., 188 ; *Wight, Ic.*, t. 1847 ;

LAURINER

Sweet or Spanish Chestnut

CASTANEA
vulgaris.

which
India
r parts
Arabia,

MEDICINE.
Plant.
806

natives in a vapour bath for
being placed under the bed"
Pindi, Panjab) "Sanskrit
and regard it as possessing

the property of increasing the secretion of semen" (U C Dutt, Civil
Medical Officer, Serampore)

Domestic—"A portion of the plant is by the Santal tied round the
neck, arm, and ankles, as a cure for rickets" (Rev A Campbell, Report,
Chutia Nagpur)

DOMESTIC.
Charm.
807

CASTANEA, *Gärtn*, *Gen Pl*, III, 409

[FERE

808

Castanea vulgaris, *Lam*, *DC Prodr*, xvi, 2, 114, 683, *CUPULI-*

THE SWEET CHESTNUT OF SPANISH CHESTNUT, CHÂTAIGNIER,
Fr, *EDELKASTANIE*, *Germ*

Syn.—*C vesca*, *Gärtn*

References—*Brandis*, *For Fl*, 491, *Gamble*, *Man Timb*, 319, *DC*, *Orig*
in of Cult Pl, 353, *Smith*, *Disc*, 110

Habitat—"A large, long-lived, deciduous tree, of rapid growth, more
rapid than the oak, introduced in the Himálaya, and grown in various
localities, and especially in a large number of places in the Panjáb and
the hills of the North-West Provinces, in Darbhing, and the Khásia
Hills" (*Gamble*)

Cultivation—"It has been sown or planted in several parts of the

CULTIVATION
809

state of the species" (*DeCandolle*, *Orig Cult Pl*)

Food—The nuts are eaten When ground into meal they form an
important article of food for the poor Mr Atkinson says the tree was
introduced by Sir John Strachey in Kumáon, and in Dehra by Dr.

FOOD.
810

TIMBER.
811

CASTANOPSIS
tribuloides.

Probable New Tanning Material for India.

vigorously, along the Voeges it is grown for vineyard poles, in Kent and Sussex for hop poles" (Brandis)

CASTANOPSIS, Spach, Gen Pl, III, 409

812

Castanopsis indica, Alph DC, Prodr, XVI, 2, 109, CUPULIFERÆ

References —Brandis, For Fl, 490, Gamble, Man Timb, 388, Kurz, For. Fl, Burm, 478, Balfour, Cyclop

FOOD.

813

TIMBER

814

both in

largely
oiled

and the branches burnt for manure.

815

C. rufescens, Hook f. & Th, Gamble, Man Timb, 389

Vern —Dainé katás, NEPAL, Sirikishu, LEPCHA, Hingori, Ass

Habitat.—A very large evergreen tree of Sikkim Himalaya, from

FOOD.

816

TIMBER.

817

818

C. tribuloides, Alph DC, Prodr, XVI, 2, III, Wight, Ic, t 770

BURN

References.—Gamble, Man. Timb, 389; Brandis, For Fl, 490; Balfour, Cyclop

FOOD

819

TIMBER

820

Structure of the Wood —Grey, moderately hard. Annual rings marked by darker lines Used for planking and shingles, being good and durable.

C. 820

The Bay Chestnut The Ule Tree

CASTILLOA
elastica.

The tree coppices admirably and with *Castanopsis indica*, *Quercus spicata*, and *Engelhardtia* might be grown on the hills wherever firewood and charcoal forests are required

CASTANOSPERMUM, *A Cunn*, *Gen Pl*, I, 556

"A genus of plants so named in consequence of the supposed resemblance of the seeds to the sweet chestnuts of Europe

Castanospermum australe, *A Cunn*, LEGUMINOSÆ

821

THE MORETON BAY CHESTNUT

References—*Drury*, *U Pl* 124 *Balfour*, *Cyclop*, *Smith*, *Dic*, 110
Treasury of Botany

FOOD,
822
TIMBER
823

CASTILLOA, *Cerv*, *Gen Pl*, III, 372Castilloa elastica, *Cerv*, URTICACÆ

824

THE ULE TREE

References.—*Brandis*, *For Fl*, 427, *Kurz For Fl*, *Burm*, II, 419;
Smith Dic, 87-89 *Spons Encyclop*, 1659-61 *Reports of Bot Gar*
dens Nilgiri Hills, for 1881-82, 1882-83, and 1885-86

Habitat—A lofty forest tree of the Bread fruit family, native of America, lately introduced into Ceylon and some parts of India. In *Kew Report for 1877*, p 15, is given an account of the attempts made to introduce this plant into India. Burma, Assam, Ceylon and the lower slopes of the Nilgiris have now been pronounced as suitable for its cultivation.

Mr Lawson reports of the Nilgiri plants "In these days of uncertain coffee crops and low prices, planters are anxious to cultivate any plant

because we have not yet learnt how to tap the trees properly"

Gum—The tree exudes, on tapping, a milky juice which, when thickened, forms what is called the Central American rubber. In some coun-

GUM
825

of the ju ce of *Ipomœa bona-nox*

For further particulars of this gum see under "India rubber"

Castor Oil, see *Ricinus communis*, *Linn*, EUPHORBIACÆ

C. 825

CASTANOPSIS
tribuloides
Probable New Tanning Material for India

vigorously, along the Vosges it is grown for vineyard poles, in Kent and Sussex for hop poles" (*Brandis*)

CASTANOPSIS, Spach, Gen Pl, III, 409

Several species of this genus are met with on the mountains of Eastern India, but none are reported to be used for tanning. This is probably an oversight, since the European members possess this property to a considerable extent, *Castanea vesca* containing 14 to 20 per cent of tannic acid

812

Castanopsis indica, Alph DC, Prodr, XVI, 2, 109, CUPULIFERÆ

Syn — *CASTANEA INDICA* Roxb, *Fl Ind Ed C B C*, 674 *Kurz*, II, 478;

Vt

Cha
khyā,

References — *Brandis For Fl*, 490, *Gamble, Man Timb*, 388 *Kurz For Fl*, *Burm*, 478; *Balfour Cyclop*

FOOD
813
TIMBER
814

and the branches burnt for manure

815

C. rufescens, Hook f & Th, Gamble, Man Timb, 389

Vern — *Dalnē katus* NEPAL, *Sirikishu* LEPCHA, *Hingori*, ASS

Habitat — A very large evergreen tree of Sikkim Himālaya, from

FOOD,
816
TIMBER
817

818

C. tribuloides, Alph DC, Prodr, XVI, 2, III, Wight, Ic, t 770

Syn — *CASTANEA TRIBULOIDES*, *Kurz* (II, 480), *QUERCUS FEROX* and *Q ARMATA*, *Roxb, Fl Ind, Ed C B C*, 673

Vern — *Tāmari katony* KUMAON *Musrē katus kotur, chiss maku, shingali* NEPAL, *Bar hingori, kanta singar*, ASS *Dingsaot*, KHASIA, *Singhara*, TIPPERAH, *Kantia lal batana*, CHITTAGONG, *Kyantsa*, BURM

References — *Gamble, Man Timb*, 389, *Brandis For Fl*, 490; *Balfour, Cyclop*

FOOD
819
TIMBER
820

Structure of the Wood — Grey, moderately hard Annual rings marked by darker lines Used for planking and shingles being good and durable

C 820

The Bay Chestnut. The Ule Tree.

CASTILLOA
elastica.

The tree coppices admirably, and with *Castanopsis indica*, *Quercus spicata*, and *Engelhardtia* might be grown on the hills wherever firewood and charcoal forests are required

CASTANOSPERMUM, *A. Cunn* ; *Gen Pl*, I, 556

"A genus of plants so named in consequence of the supposed resemblance of the seeds to the sweet chestnuts of Europe"

Castanospermum australe, *A. Cunn* ; LEGUMINOSÆ

821

THE MORETON BAY CHESTNUT

References.—*Drury, U Pl*, 124, *Balfour, Cyclop*, *Smith, Dic*, 110, *Treasury of Botany*

Habitat.—A tree of the sub-tropical regions of Australia, occasionally planted for ornament, introduced into India about thirty years ago

Food.—The seeds are eaten by the natives of Australia, but are unpalatable to Europeans (*Smith*)

FOOD.
822
TIMBER
823

Structure of the Wood.—White, with a yellowish tinge, hard

CASTILLOA, *Cerv*, *Gen Pl*, III, 372Castilloa elastica, *Cerv*, URTICACÆ

824

THE ULE TREE

References.—*Brandis, For Fl*, 427 *Kurz, For Fl, Burm*, II, 419; *Smith, Dic*, 87, 89 *Spons' Encyclop*, 1659-61 *Reports of Bot Gardens, Nilgiri Hills*, for 1881-82, 1882-83, and 1885-86

Habitat.—A lofty forest tree of the Bread fruit family, native of America, lately introduced into Ceylon and some parts of India. In *New Report for 1877*, p 15, is given an account of the attempts made to introduce this plant into India. Burma, Assam, Ceylon, and the lower slopes of the Nilgiris have now been pronounced as suitable for its cultivation

Mr Lawson reports of the Nilgiris—
coffee crops and low price
that will return a small profit
localities in the Wynad
suit the Castilloa, and
vator" Colonel Campbell
Calicut "It has been
hope they
this place
either from
because we
Gum—
and it is—

GUM
825

For further particulars of this gum see under "India rubber"

Castor Oil, see *Ricinus communis*, *Linn*, EUPHORBICÆ

C. 825

Beefwood of Australia.

826

Syn —C MURICATA, Roxb, Fl Ind, Ed C B C. 623

Vern.—Yangli saru, HIND , Fan, BENG , Vilayatisaru, wilayati sarī, saraka jhar, BOMB , Furijur, muzzun, SIND , Sarpuhala, sarova, suru,

tions. Conf with Tamarix.

Wye's of
Bomb
Lisboa,
on, in
Balfour.

Habitat—A large, evergreen tree, with leafless, drooping branches

CULTIVATION
827

the vernacular names of that plant

Cultivation—"It has been largely planted in North Arcot, South Arcot, Madras, and other districts of the Madras Presidency, for fuel, for which it is excellent, but it requires to be near the sea-coast and to have water at the roots, at least 10 feet from the surface of the ground. Trees planted in sandy soil often suffer much from drought the first two or three years, the tap-root then finds its way down to about 10 feet, and reaching water the tree begins to thrive. It is of course best near the sea, but fine trees may be seen in places in Northern India, especially at Saharanpur and Amballa" (*Gamble*).

The Ma
cost of cultu
put down a
gross capita
Valuing the
of B181

and in the eighth or ninth year the land may be cleared, the remaining trees, at the lowest estimate, after paying all expenses on the same, would realize R600

Gum.—Reported to yield a good resin

Dye.—The bark is used in tanning (*Birdwood, Bomb Prod.*, and *Bide, Mad Exh List for 1855*). A brown dye is extracted from it according to Balfour. Mr Wardle remarks "The bark contains a small quantity of colouring matter, and produces in dyeing light reddish drab colours on each of the fabrics on which I have experimented." He further adds—"The shades produced by this dye-stuff are very good

CUM.
828
DYE
820

C. 829

Cedrelas or Toon woods

CEDRELA.

though faint, but the dye-stuff contains too small an amount of colouring matter to be of any great value in the dye house. Lisboa says that it is used in Bombay as a mordant.

Medicine — The bark is slightly astringent, and is employed in infusion as a tonic, according to Dr Gibson it is an excellent and at the same time a readily available astringent, useful in the treatment of chronic

e, very hard,
cut, weighs
pice well, and
tremely quick
important trees
wood is used

for fires, as it burns readily, and the ashes retain the heat for a long time. It is much valued for steam engines, ovens, &c." (*Treasury of Botany*). Clubs made of the hard wood are used in Fiji for beating the bark of the PAPPE MULBERRY (*Broussonetia papyrifera*, Vent) for the manufacture of Tapa cloth (*New Official Guide to Museums*, 121). The natives of Australia make their war-clubs from this wood (*Smith*).

Domestic Uses — "The burnt ash is made into soap" (*Smith*).

DYE

MEDICINE.
830TIMBER
831DOMESTIC
Ash
832

Catechu, see—

[A 139] (a) *Acacia Catechu*, Willd., LEGUMINOSÆ (black catechu)

(b) *Uncaria Gambier*, Roxb., RUBIACÆ (pale catechu)

[A 1298] (c) *Areca Catechu*, Linn., PALMÆ (palm catechu)

Cattle and Buffaloes see Oxen

Cat, Civet, see Tigers and Panthers.

Catha. Several species exist in India, but by the *Flora of British India* they have been all reduced to *Celastrus*, which see

Catha edulis yields the *Kat* or *Kafter* of the Arabs, the leaves of which if chewed are said to prevent sleep. Sometimes imported into India, largely so to Aden, where they are used as a substitute for Tea.

Cat's-eyes, see Chalcedony.

Cat's-skins, see Skins.

Cauliflower and Brocoli, see *Brassica (oleracea) botrytis*, B 851

Caustic Potash, see Potassium, also Carbonate of Potash, C. 527.

Caustic Soda, see Sodium, also Carbonate of Soda.

CEDRELA, Linn., *Gen Pl*, I, 339

834

reco
to d
the
C ..

part
branches, from which when in flower a panicle three or four feet long is suspended. This is the characteristic form of the North-Western Himálaya at altitudes from 4,000 to 8,000 feet. It frequents damp shady streamlets, growing so gregariously as to exclude all other trees

C. 834

CEDRELA
serrata.

The Toon woods.

In the Monograph of the Meliaceæ published in 1878 by Oasimir de Candolle, the species of *Cedrela* formerly grouped under the one head of *Cedrela Toona*, *Roxb*, have been separately described.

They are thus distinguished —

Ovary glabrous—

Leaflets petioled

C. serrata, *Royle*

Leaflets subsessile

C. glabra, *C de Cand*

Ovary hairy—

Leaflets acute at the base

C. Toona, *Roxb*

Leaflets round at the base

C. microcarpa, *C de Cand*

Mr Gamble, in his *Manual of Timbers*, XII, remarks that in his "*Trees, Shrubs, and Climbers of the Darjiling District*, three varieties were spoken of and separated as follows —

ber-December, bark light-
es, found in the upper hills

"No 1 is *C. Toona*, *Roxb*. No 2 probably *C. microcarpa*, *C de Cand*., No 3 probably *C. glabra*, *C de Cand*. It would, however, have probably been better to describe No 1 as 'deciduous in the cold season,' and Nos 2 and 3 as 'deciduous in the rains' There is perhaps a fifth species

"They may also be distinguished as follows by the capsule —

Capsule smooth { capsule round

C. Toona.

{ long, pointed

C. microcarpa

Capsule covered with corky tubercles

C. glabra.

"Of the Northern Bengal specimens which we have examined, E 360 and E 2333 will be *C. glabra*, while E 655, E 2332, E 3509, E 3619 and E 3623 will be *C. microcarpa*. Some of the Assam, Chittagong, and

000 feet, is probably *C. multijuga*,
RM, Nee, KAREN (Trade name,

It has a light, soft, pink wood,
with the usual characteristic scent strongly perceptible, and structure
resembling that of the other species of Toon, the pores being perhaps

cepted as indicating
microcarpa, *DC*, as

835

Cedrela serrata, *Royle*, *Ill*, p 144, t 25; *Monog*, *DC*, I, 742,

[*MELIACEÆ*

Syn — *C. Toona*, *Roxb* (*Hook*, *Fl Ind*, 1, 558 in part)

Vern — *Drawi*, *dalli*, *dal*, *dauri*, *khishing*, *khinam*, N.W. H

TIMBER
836

large pores,

C. 836

The Toon woods.

CEDRELA
Toona

Domestic Uses—Used about Simla, for the hoops for sieves for bridges, and for many such purposes. The shoots and leaves are lopped for cattle fodder.

DOMESTIC
837
FODDER.
838

Cedrela Toona, Roxb., *Fl. Br. Ind.*, I, 568, *Wight, Ic.*, t. 161.

THE TOON OR INDIAN MAHOGANY TREE, MOULMEIN CEDAR.

Vet. — T. — 1 — 2 — 3 — 4 — 5 — 6 — 7 — 8 — 9 — 10 — 11 — 12

BURM

References — *Roxb., Fl. Ind.*, Ed. C. B. C., 213, 633, *Brandis For. Fl.*,

and Australia

Gum—It yields a resinous gum, of which little is known at present

M. Nees von Essenbeck has published an account of some experiments with the bark, which indicate the presence in it of a resinous

GUM.
839

DYE
Flowers
840
Seeds.
841

"It was a commoner practice under native rulers than it appears to be now to wear *bisanti*-coloured clothes in the spring, whence its name *bisanti*" or spring time. Safflower and tûn are combined in Tirwa. Dr. McCann

C. 841

CEDRELA
serrata

The Toon woods

In the Monograph of the Meliaceæ published in 1878 by Gasimur de Candolle, the species of *Cedrela* formerly grouped under the one head of *Cedrela Toona*, *Roxb*, have been separately described.

They are thus distinguished —

Ovary glabrous—

Leaflets petioled

Leaflets subsessile

C serrata, *Royle*

C glabra, *C de Cand*

Ovary hairy—

Leaflets acute at the base

Leaflets round at the base

C Toona *Roxb*

C microcarpa, *C de Cand*

Mr Gamble, in his *Manual of Timbers XII*, remarks that in his "Trees, Shrubs, and Climbers of the Darjiling District, three varieties were spoken of and separated as follows —

No 1 Deciduous in the rains

No 2 Deciduous in the cold season

No 3 Deciduous in the cold season

No 4 Deciduous in the cold season

No 5 Deciduous in the cold season

No 6 Deciduous in the cold season

No 7 Deciduous in the cold season

No 8 Deciduous in the cold season

No 9 Deciduous in the cold season

No 10 Deciduous in the cold season

No 11 Deciduous in the cold season

No 12 Deciduous in the cold season

No 13 Deciduous in the cold season

No 14 Deciduous in the cold season

No 15 Deciduous in the cold season

No 16 Deciduous in the cold season

No 17 Deciduous in the cold season

No 18 Deciduous in the cold season

No 19 Deciduous in the cold season

No 20 Deciduous in the cold season

No 21 Deciduous in the cold season

No 22 Deciduous in the cold season

No 23 Deciduous in the cold season

No 24 Deciduous in the cold season

No 25 Deciduous in the cold season

No 26 Deciduous in the cold season

No 27 Deciduous in the cold season

No 28 Deciduous in the cold season

No 29 Deciduous in the cold season

No 30 Deciduous in the cold season

No 31 Deciduous in the cold season

No 32 Deciduous in the cold season

No 33 Deciduous in the cold season

No 34 Deciduous in the cold season

No 35 Deciduous in the cold season

No 36 Deciduous in the cold season

No 37 Deciduous in the cold season

No 38 Deciduous in the cold season

No 39 Deciduous in the cold season

No 40 Deciduous in the cold season

No 41 Deciduous in the cold season

No 42 Deciduous in the cold season

No 43 Deciduous in the cold season

No 44 Deciduous in the cold season

No 45 Deciduous in the cold season

ber December, bark light
es, found in the upper hills

"No 1 is *C Toona*, *Roxb*, No 2 probably *C microcarpa*, *C de Cand*. No 3 probably *C glabra*, *C de Cand*. It would however, have probably been better to describe No 1 as 'deciduous in the cold season,' and Nos 2 and 3 as 'deciduous in the rains' There is perhaps a fifth species

"They may also be distinguished as follows by the capsule —

Capsule smooth { capsule round

 { long pointed

Capsule covered with corky tubercles

C Toona.

C microcarpa

C glabra

"Of the Northern Bengal specimens which we have examined, E 360 and E 333 will be *C glabra*, while E 655 E 2332, E 3599 E 3619 and E 3623 will be *C microcarpa*. Some of the Assam, Chittagong, and Burma specimens are probably *C microcarpa*

"No B 3378 from the Salween, 2000 feet, is probably *C multijuga*, *Kurz*, t, 229 — Vern *Toungdama*, BURM, *Nee*, *HAREN* (Trade name, like the other Toon woods, *Thitktado*) It has a light, soft, pink wood, with the usual characteristic scent strongly perceptible, and structure resembling that of the other species of Toon, the pores being perhaps more scantily distributed. Weight 35 lb per cub c foot"

The preceding remarks may for the present be accepted as indicating the Nepal plant, *C glabra*, *DC*, and the Sikkim *C microcarpa*, *DC*, as distinct from the following —

835

Cedrela serrata, *Royle*, Ill, p 144, t 25, *Monog*, *DC*, I, 742,

[MELIACEÆ

Syn — *C TOONA*, *Roxb* (*Hook*, *Fl Ind*, t 566, in part)

Vern — *Drawi*, *dalli*, *dál*, *dauri*, *khishing*, *khinam*, N W H

" " " " " "

TIMBER
836

large pores.

C. 836

The Toon woods.

CEDRELA
Toona

Domestic Uses —Used about Simla, for the hoops for sieves for bridges, and for many such purposes. The shoots and leaves are lopped for cattle fodder.

DOMESTIC
837
FODDER
838

Cedrela Toona, Roxb., Fl. Br. Ind., I, 568, Wight, Ic., t. 161

THE TOON OR INDIAN MAHOGANY TREE, MOULMEIN CEDAR.

Vern —*Tun, tuni, lum maha nim, maha limbo tunka jhar, tuna, lud,*

BURM

References —*Roxb., Fl. Ind., Fd. C.B.C., 213, 633, Brandis, For. Fl.,*

Ann. D. Ind. 14, Dyer Dyes and Mats, IV - 11, 1, 28, 23, Bird wood, Bomb. Prod., 325, Lisboa U. Pl. Bomb. 46, 241, 258, Balfour, Cyclop. Treasury of Bot. Kew Cat., 29, Fleming's Med. Pl. and Drugs in As. Socy. Res., Vol. XI, 163, Med. Top., IX, 93

Habitat —A large tree, about 50 to 60 feet in height, growing in the tropical Himalaya from the Indus eastward, and throughout the hilly districts of Central and South India to Burma, ascending to 3,000 feet in the N.-W. Himalaya and in Sikkim (?) to 7,000 feet. Distributed to Java and Australia.

Gum —It yields a resinous gum, of which little is known at present.

M. Nees von Essenbeck has published an account of some experiments with the bark, which indicate the presence in it of a resinous astringent matter, a brown astringent gum and a gummy brown extractive matter, resembling *Ulmine* (Balfour).

Dye —The flowers yield a red and a yellow dye (in Bengal generally known as *Gulnari*) said to be used to a small extent only. Madras dyes sent to Paris.

It is known as *basanti* in the north-west provinces. It is fleeting and apparently only used by the poorer classes. In Burma it is used in conjunction with safflower. Sir E. Buck, in his *Report on the Dye-stuffs*,

GUM.
839

DYE.
Flowers
840
Seeds,
841

of Cawnpore is produced.

"It was a commoner

to wear *basanti*-colour

or spring time. Safflower

Dr. McCann

C. 841

CEDRELA
Toona.

The Toon-woods: Moulmein Cedar

DYE.

says the cloth previously dyed yellow is changed into red by the *pán* eaten by Hindus.

colc

MEDICINE
Bark
842

usel

be
Dr.
an
extract of the bark in chronic infantile dysentery Blume attributes valuable antiperiodic virtues to it, and in this character it is favourably noticed by Dr J Kennedy (*Ann of Med*, 1796, Vol I, p 387) Dr Æ. Ross speaks of it as a reliable antiperiodic, and Dr J Newton as a good substitute for cinchona The dose of the dried bark is about an ounce daily in the form of infusion The powder of the bark was found by Dr Kennedy to be of great service as a local astringent application in various forms of ulceration According to Dr Dymock, the native physicians use the bark in combination with bonduc nuts as a tonic and antiperiodic, a fact also mentioned by Ainslie in his *Materia Indica* The FLOWERS are called *Gul-tan* in Bombay and considered emmenagogue "The bark was used in Java by Blume in epidemic fevers, diarrhœa, and other complaints Horsefield gave it in dysentery, but only in the last state, when inflammatory symptoms had disappeared" (*Balfour*)

Flowers
843FOOD
844TIMBER
845

Food—The seeds are used to feed cattle The young shoots and leaves are lopped as cattle fodder.

Structure of the Wood—Brick-red, soft, shining, even but open grained, fragrant, seasons readily, does not split nor warp Annual rings distinctly marked by a belt of large and numerous pores

It is durable and is not eaten by white ants, is highly valued and universally used for furniture of all kinds, and is also employed for door-panels and carving From Burma it is exported under the name

Price

and is used in
ent cases It
or many other

purposes

C. 845

The Deodar or Himalayan Cedar.

CEDRUS
Deodara.CEDRUS, *Loud*, *Gen Pl*, III, 93Cedrus Deodara, *Loudon*, *DC Prodr*, XVI, 2, 409

846

DEODAR HIMALAYAN CEDAR.

Syn.—PINUS DEODARA, *Roxb*, *Fl Ind*, Ed C B C, 677Vern.—*Kilan ka-phér*, *kilan deodar*, HIND, *Dewdār*, *geyar*, *keli*, *kelu*,*hindi*, *nashtar* PERSReferences—*Brandis*, *For Fl*, 516, *Gamble*, *Man Timb*, 400, *Stewart*,*Pb Pl**Sheriff**296 D**Dispen**410, 4**Dist*,*four*,*Irvine*, 28

Habitat—A very large and tall tree, found in the North-West Himalaya, between 4 000 and 10 000 feet, extending east to the Dauli river (a tributary of the Alaknanda below the Niti Pass), in the mountains of Afghanistan and in North Beluchistan

Gum—It yields a true oleo-resin, called *Kelon-ka-tel*. The preparation of this oleo resin is thus described by Mr Baden Powell—

First, an earthen *ghara*, or vessel with a wide mouth, and capable of containing about 4 seers, is sunk into the ground. Next, a large *ghara* of about 12 seers capacity is taken, and three small holes are drilled in its under side, it is then filled with scraps of the pine wood,

GUM
847

be made to do over wood yields about 26 *chistaks* of tar and 4·3 *chistaks* of charcoal. To procure a seer of tar to charge the pot, and 2 maunds (*Pb Prod*, 410) wood by destructive distillation, and resembles crude turpentine. It is used for anointing the inflated skins which are used for crossing rivers, and as a

C. 847

CEDRUS
Deodara.

The Deodar or Himalayan Cedar.

MEDICINE
848

remedy for ulcers and eruptions, for mange in horses and sore feet in cattle" (*Gamble, 406*)

Medicine—The aromatic wood is employed medicinally as a carmi-

FOOD
849
TIMBER.
850

stomach could bear. Its use may be extended to other skin diseases with advantage. Dr Royle states that the leaves and small twigs of the *Deodara* are also brought down to the plains, as they are supposed to possess mild terebinthinate properties (*Plarm Ind*). In Kangra the wood is pounded with water on a stone, and the paste applied to the temples to relieve headache. Assistant Surgeon Sakharam Arjun describes the wood as a bitter stomachic, useful in fever, costiveness, piles, and pulmonary complaints.

Food—The young shoots and plants are eagerly browsed by goats, &c.

Structure of the Wood—Heartwood light-yellowish brown, scented, moderately hard. In each annual ring the outer belt of firmer and

the edge of certain annual rings are frequently found concentric strings of dark-coloured pores or intercellular ducts, which are prominent on a vertical section as dark lines, and in the vicinity of which the wood is sometimes more resinous.

In common with most species of the Order, the *Deodar* has well marked annual rings which, there is little, if any, reason to doubt, each repre-

warm
kes it
from
lly be

the practice to take only for use in any forests, the experiments made on trees in that or neighbouring localities. But the experience we have

The Oleum Nigrum.

CELASTRUS
paniculata

- inner Himálaya, having usually the age of trees 6 feet in girth above 140 years,
 2nd—Those in the intermediate ranges and valleys, having 6 feet in girth for an age of between 110 and 140 years,
 3rd—Those in the outer ranges under the full influence of the monsoon, and having the age of trees 6 feet in girth usually below 110 years.

Deodar wood is extremely durable, being by far the most durable of the woods of the Himálayan conifers. It is the chief timber of North-West India, and is used for all purposes of construction,—for railway sleepers, bridges, and even for furniture and shingles. (*Gamble*.)

TIMBER.

CELASTRUS, Linn.; Gen Pl, I, 364.

851

The *Flora of British India* raised Wight and Arnott's sub-genera (1) *EUCELASTRUS* and (2) *GYMNOSPORIA* to the rank of genera. This was at first followed by the authors of the *Genera Plantarum*, but subsequently (Vol I, page 997) was corrected back to the original position. The former embraces some four species of unarmed climbers, and the latter five.

Celastrus emarginata, Willd.; CELASTRINEÆ.

852

Syn.—*GYMNOSPORIA EMARGINATA*, Roth, in *Fl. Br. Ind.*, I, 621, *CELASTRUS EMARGINATA*, W. and A., *Prod.*, 160; Roxb., *Fl. Ind.*, Ed. C.B.C., 208, *CATHA EMARGINATA*, G. Don.

C. oxyphylla, Wall.

853

Syn.—*GYMNOSPORIA ACUMINATA*, Hook. f.; *Fl. Br. Ind.*, I, 619

C. paniculata, Willd.; *Fl. Br. Ind.*, I, 617; Wight, *Id.*, t. 158.

854

BLACK OIL; THE OLEUM NIGRUM PLANT.

Syn.—*CELASTRUS ALNIFOLIA*, Don.; *C. DEPENDENS*, Wall.; *C. MULTIFLORA* and *NUTANS*, Roxb.

Vern.—*Mal kangni*, *mál kungí*, HIND., *Sankhú*, *sankhú* (leaves, *kotaj*, *kutaj*)

na-young, BURM. The vern. names of *Oleum Nigrum*: *Malkangni*, *ka-jantar*, DUK, *Valuluvai-tailam*, TAN, *Malkanginitailam*, TEL.
 References.—Roxb., *Fl. Ind.*, Ed. C.B.C., 209. Brandis *For Fl.*

CELASTRUS
paniculata

The Oleum Nigrum.

OIL
855

Habitat — A scandent shrub of the outer Himálaya, from the Jhelum to Assam, ascending to 4,000 feet, Eastern Bengal, Behar, South India, and Burma, in Ceylon it is common up to an elevation of 2,000 feet
Oil.—The seeds yield by expression a deep scarlet or yellow oil, used medicinally.
time Its odor
turns of a dar
cation along
leaves and seeds of a

MEDICINE.
OIL
856

... small blue
12 annas to
... for cattle
They are given in rheumat
obtained from the seeds by
externally This oil, under t
forward by the late Dr H
When administered in doses of from ten to fifteen drops twice daily, its
action as a powerful stimulant is generally followed in a few hours by
free diaphoresis not attended by exhaustion It is specially efficacious in

Seeds
857

aphrodisiacal and stimulant, useful both as an external and internal

Leaves
858

yellow and of the consistence of oil The black oil manufactured at Vizagapatam and Masulipatam is the best It is a good diuretic, diaphoretic, and nerve stimulant It is certainly the best remedy for *beriberi* I have seen many cases which did not benefit for weeks or months under the use of other medicines, but began to improve at once when this oil was employed The first good effect of this medicine is generally the increase in the quantity of urine, and with this the dropsical effusion

The Oleum Nigrum

CELASTRUS
senegalensis.

patient except milk and bread—a restriction which is as injurious as un- MEDICINE.

Food for
Quails.

patient is under this treatment he should eat meat roasted I have seen
two or three cases of *beri-beri* cured by this treatment, and have also

diet, while using it, should consist exclusively of wheaten cakes and flesh
of sheep" (*Honorary Surgeon P Kinsley, Ganjam, Madras Presidency*)
"An oil extracted by heat is a specific in the treatment of *beri-beri* with
marked success

Is a stimulant and
diet should be of
and milk, and no

among the people of the Northern Circars, especially of those of the
malarious tracts" (*Surgeon-Major E. W. Levinge, Rajamundry, Godavery
District*). "Said to be useful as an aphrodisiac" (*Surgeon-Major D. R.
Thompson, Madras*).

Structure of the Wood.—Pinkish yellow, soft.

TIMBER.
859
860

Celastrus senegalensis, Lam.

Syn.—

621;

Vern.—

khar

babu

peddi

References.—*Roxb. Fl. Ind., Ed. C.B.C., 208. Brandis, For. Fl. &,
Kurz, Fl. Burm., I, 252; Beddome, Fl. Sylhet., LXVI; Dals. & Gibs,
Bomb. Fl., 48. Gamble, Man. 11mb, 87.*

Habitat.—A profusely-armed tall shrub, common in the northern dry
and intermediate zones of Central, South-Western, and North-Western
India, distributed to Afghanistan, Central Asia, and Australia. The
Flora of British India, Afghanistan, Central Asia, and Australia. The

comprises
the leaves
stems are
narrower

Medicine.—The BARK, ground to a paste and applied to the head, with
mustard oil, is said to destroy *pediculi*.

MEDICINE.
Bark.
861

CELOSIA
argentea.

Celestite; Celosia

862

Celastrus spinosus, Royle.Syn.—GYMNOSPORIA ROYLEANA, Wall., as in *Fl Br Ind*, I, 620Vern.—*Yaliddhar*, HIND, *Dsaral*, TRANS INDUS, *Kandu*, *kamla*, *kandari*, *kander*, *löp*, *patäki*, *lei li*, *phupari*, *badlo*, *kadewar*, Pn, *Kura*, *bagriwala darim*, *gwäla darim*, N-W PReferences.—Boiss, *Fl Orient*, II, 11 Brandis, *For Fl*, 80, Gamble, *Man Timb*, 86, Baden Powell, *Pb Prod*, 582, Stewart, *Pb Pl*, 41.

Habitat.—A thorny, distorted bush, abundant on the outer North-Western Himälāya (Kumaon and Garwhal, altitude 1,000 to 4,500 feet) and distributed to the Concan and thence to Afghānistan, common on the

MEDICINE

Seed

863

TIMBER

864

to be

ained,

on as a

Baden

possible substitute for boxwood, for carving and engraving Powell remarks that it is used in the Panjāb for walking-sticks.

865

Celery. See *Apium graveolens*, Linn, UMBELLIFERÆ

Bombay

866

CELESTITE; Mallet, *Mineralogy*, 141.

Celestite or Celestine is a natural mineral, found in rhombic or

Punjab

867

the Salt Range

CELOSIA, Linn, *Gen Pl*, III, 24.For botanical characters of the genus see under *Amarantaceæ* (A. 914).The name is derived from *kelos*, burnt, in reference to the colour of the flowers in the common garden species

868

Celosia argentea, Linn, Fl Br. Ind, IV, 714, AMARANTACEÆVern.—*Debkoti*, *sufaid murgha*, *sarwari*, HIND; *Sirgit arak*, SINTAL, *Sarmali*, *sirālī*, *ghogiya*, N-W P, *Sarwali*, *salgāra*, *chulchil*, *sil*, *sarpankha*, Pn, *Swet-murgā*, BENG, *Surwali*, *ucha kukur*, SIND; *Lapadi*, GUJ, *Kudhu*, *kurdu*, BOMB, *Kurdā kurada*, MAR, *Gurugu*, *panche chettu*, TEL; *Kirri handa*, SING Several of these vernacular names imply white-cock's-comb

Re

MEDICINE

Seeds

869

Oil

870

FOOD

871

FODDER

872

Habitat.—An abundant weed of the fields in Central and Northern India (from Chutia Nagpur to the Panjāb), occasionally ascending to altitude 5,000 feet in the Himälāya, it is also met with in the warmer parts of Ceylon. It appears very commonly in the monsoon season

Medicine.—The SEEDS are officinal, being an efficacious remedy in diarrhœa. The Rev A Campbell says the Santals extract a medicinal oil from them.

Food.—The plant is used as a pot-herb in times of scarcity, and is eaten by cattle, especially buffaloes.

C. 872

Celosia ; Celsia

CELSIA
coromandeliana

Celosia cristata, Linn , Fl Br. Ind , IV , 715 , Wight, Ic , t 730

873

Ve

References — Roxb Fl Ind , Ed C B C , 229 , Dals & Gibs , Bomb
Fl , 215 , Stewart Pb Pl , 182 Murray Drugs and Pl , Sind , 101 ,
Baden Powell Pb Pr 373 , Balfour, Cyclop , Treasury of Botany ,
Spons, Encyclop , 938

Habitat.—Cultivated as an ornamental plant in the plains, and on the
Himalaya, Kashmir (5000 feet) In Spons Encyclopædia occurs the
remark that this plant is 'Common all over Bengal and Northern India
generally'

Fibre—"It yields a strong flexible fibre, so highly esteemed that rope
made of it sells at five times the price of jute rope' Confirmation of this
fact is much required, and also samples of the plant from which the fibre
has been extracted It is known in Bengali as *Lal-murga*, but Roxburgh
makes no mention of the fibre, indeed, with the exception of the notice in
Spons Encyclopædia quoted above, no author, as far as the writer can
discover, alludes to the fibre

FIBRE
874

Medicine—The FLOWERS are officinal, being considered astringent
they are used in cases of diarrhoea and in excessive menstrual discharges
The SEEDS are viewed as demulcent

MEDICINE
Flowers
875
Seeds
876

Special Opinion — § 'Seeds demulcent and useful in painful micturi-
tion, cough and dysentery' (Dr U C Dutt, Serampore)

Food—Cultivated in gardens—both the red and the yellow forms—on
account of the stem which is eaten as a pot herb Professor Church
(in *Food Grains of India*) is apparently in error when he speaks of the
food properties of the seeds of this plant The writer can find no mention
of the plant being cultivated on account of its seeds nor indeed of these
being eaten Besides three of the vernacular names given by the Pro-
fessor are not names for this plant *Sil* (and names derived from that
word) are more correctly applied to *Amarantus paniculatus*, the seed of
which is eaten so that it seems probable Professor Church's account of
Celosia cristata should be transferred to *Amarantus paniculatus*

FOOD
877

CELSIA, Linn , Gen Pl , II , 929.

Celsia coromandeliana, Vahl , Fl Br Ind , IV , 251 , Wight, Ic ,
t 1406 SCROPHULARINEÆ

878

Vern — *Kukshima koksind* BENG , *Kutki*, MAR , *Kuluhala* SANS

Cyclop

and
durcase
gent (*Pharm of Ind*)MEDICINE
Juice
879

Special Opinions — § "Juice of the whole plant, including the root,
leaves and stem, squeezed out by pounding it, is used in half chittack
doses, morning and evening, in cases of syphilitic eruptions The juice of

R

C. 879

CELTIS
caucasica.
The Honey-berry.
MEDICINE.
Root
880

"The root is used in dysentery and as a cholagogue" (*Brigade Surgeon*
J. H. Thornton, Mongkur).

CELTIS, *Tourn.*; *DC. Prodr.*, XVII., 168.

881

Celtis australis, *Linn.*, *DC. Prodr.*, xvii., 169, 170, 179; **URTICACEÆ.**

THE EUROPEAN NETTLE-TREE, THE HONEY-BERRY TREE.

FOOD.
Fruit.
882
FODDER.
883

Habitat.—A moderate-sized, deciduous tree, found in the Suliman and Salt Ranges, and throughout the Himálaya from the Indus to Bhután, ascending to 8,500 feet, also in the Khásia Hills. Extensively cultivated in South Europe.

Food and Fodder.—The tree is largely planted for fodder; cows fed on the leaves are supposed to give better milk. The **FRUIT** is also eaten. "It is remarkably sweet, and is supposed to have been the Lotus of the

which Herodotus, Dioscorides, and leasant, and wholesome, and which make those who ate it forget their If eaten in Spain, and Dr. Walsh very fond of them" (*Treasury of*

Botany) It is nowhere grown as a fruit tree in India, although, as Atkinson adds, it is eaten by all classes and is esteemed.

A dark-purple form of the fruit is called *roku* and a smaller yellow form *choku*.

TIMBER.
884

Structure of the Wood.—Grey or yellowish grey, with irregular streaks of darker colour. Weight 47½ per cubic foot. It is tough and

DOMESTIC.
885

king
 y of

886

C. caucasica, *Willd.*, *DC. Prodr.*, xvii., 170.

Vern.—Bathar, brámi, brimá, brimla, bigni, biñu, kharg, khark, khirk, karik, kharak, khalk, ku, takhum, tágho, wattamman, kanrak, kirk, kar, kargam, taghum, takpun, kurg, kanghol much (the fruit), Pá., Tughar, Pushru.

C. 886

The Nettle-trees.

CELTIS
cinnamomea.

References.—Brandis, *For. Fl.*, 428, 429; Gamble, *Man. Timb.*, 341; Stewart, *Pb. Pl.*, 209; Hutchison, *Cat. Pb. Pl.*, 139; Baden Powell, *Fl. Pr.*, 574; Balfour, *Cyclop.*

FIBRE.

887

MEDICINE.

Fruit.

888

FOOD.

Fruit.

889

TIMBER.

Fuel.

890

DOMESTIC.

Crafts.

891

SANDALS.

892

893

Pl., 209)

Celtis cinnamomea, Lindl.; Kurz, *For. Fl. Burm.*, II., 472.

Syn.—*C. dysodoxylon*, Thw.

Vern.—Gurenda, Sing.

References.—Gamble, *Man. Timb.*, 343, Thw., *En. Ceylon Pl.*, 71.

stern

also

Vara-

Kya-ka (or Hell's incense), is used as a charm against evil spirits. This was described by Dr. W. Dymock in the 1st edition of his *Flora Medica of Western India* under its vernacular name. The writer's attention having been drawn to this, a correspondence was instituted. Dr. Dymock stated that the Bamba, a well-known name for *Celtis*, was the

MEDICINE.

Wood.

894

name of *Celtis dysodoxylon*.

ing people as *pudacarpa*.

by the Dutch *strunthout*, as

its disgusting odour, which results specially in the thick stem and the larger branches. The smell of it so perfectly resembles that of human ordure, that one cannot perceive the smallest difference between them.

other cutaneous eruptions, the body being at the same time anointed with it externally."

CELTIS
Wightii.
The Nettle-trees.
MEDICINE
 Price
 895

Dr. Dymock states - "The peculiar odour is probably due to the presence of *naphthylamine*. The price of the wood in Bombay is Rs 30 per cund of 24 cund. The Best of 2 cund is P. J. wood and P. 8 "

have been here recorded as a basis of further investigation, since the Indian trade in the wood is of some importance

896

Celtis eriocarpa, Dene ; DC Prodr., XVII, 179.

Vern — *Akaka, katāka*, HIND, *Bathar, bat tamanku*, PB ; *Tagha*, ARG

References — *Brandis, For Fl.* 429, *Gamble, Man Timb*, 343; *Baden Powell, Pb Pr*, 574; *Balfour, Cyclop*

Habita

Salt Ranc
from the

DOMESTIC

Domestic Uses. — The bark is used for making shoes (*baueu r. o. u. e. u.*)

897

C. orientalis, Linn See Sponia orientalis, Planch

898

C. Roxburghii, Planch, Brandis, For Fl, 429.

Syn. — *C. trinervia*, Roxb, *Fl Ind*, Ed C B C, 252

Vern — *Kharak, bathar, brāmaj, brundu*, PB, *Cher, chara, kathaniar*, C P, *Bowmaj*, BOMB.

References — *Bedd, Fl Syle.*, CCCXII, *Gamble, Man Timb*, 343; *Dals & Gids, Bomb Fl*, 273, *Lisboa, U Pl Bomb*, 131

TIMBER

899

900

C. tetrandia, Roxb, DC Prodr., XVII, 179

EUROPEAN MYRTLE TREE

Habitat. — A tall tree of the outer Himalāya, from Kumaon eastward, to the Ava Hills in Burma, also on the Western Ghats

TIMBER.

901

Structure of the Wood — Greyish white, moderately hard Used in Assam for planking and canoes.

C. trinervia, Roxb See C. Roxburghii, Planch.

902

C. Wightii, Planch, DC Prodr., XVII, 184; Wight, Ic, t. 1969

Syn — *SOLENOSTIGMA WIGHTII*, Bl, *Kurz, For Fl Burm*, II, 471

Vern — *Vella thorasay*, TAM, *Tella kaka-mushti*, TEL

References — *Gamble, Man Timb*, 343; *Thwaites, En Ceylon Pl*, 267. *Balfour, Cyclop*

Habitat. — A small evergreen tree of the mountains of South India and the Andaman Islands, is also met with in the hot dry parts of Ceylon

TIMBER.

903

Structure of the Wood — Greyish white, very hard, close-grained Weight 53 lb per cubic foot. Annual rings indistinctly marked by a narrow belt without pores (*Gamble*)

C. 903

Cements.

CEMENTS.

CEMENTS.

904

CEMENTS, *Fr.*; CEMENTE, *Ital.*, *Ger.*

The term "Cement" is applied to a class of substances used for uniting two bodies, and which ultimately harden and bind them together. The following classification of these substances from *Spon's Encyclopædia* may be here given: (a) Calcareous cements, (b) Gelatinous cements, (c) Glutinous cements, (d) Resinous cementing compounds, and (e) Non-resinous cementing compounds. Interesting information regarding the Cements of India will also be found in *Bulfour's Cyclopædia of India*.

Calcareous.
905

from 10 to 25 per cent of alumina, magnesia, and silica, yield a lime, on burning, which does not slake when moistened with water, but forms a mortar with it, which hardens in a few days when covered by water."

cements." (See Cocoa-nut Juice under *Cocos nucifera*)

(b) GELATINOUS CEMENTS—These have their origin in the substance known as "gelatine" obtained by boiling animal tissues in water. It is separated from water by simple evaporation, when it is converted into a dry hard substance called by different names, such as "glue," "size," "isinglass," &c., according to the sources from which they are derived. Of these, "glue" and "size" are employed as cements, and in India a strong and useful glue, made from cartilage obtained from fish, is used by every jeweller and gold-leaf beater.

Gelatinous.
906

(c) GLUTINOUS CEMENTS—The base of this class of cements is a sub-

Glutinous
907

this class of substances are due to the presence of resin, gum-resin, or gum, such as common rosin, india-rubber, gutta-percha, gum arabic, &c. The following are a few of the Indian plants which are known to afford substances used as cements:—

Resinous.
908

Adenanthera pavonina (seeds).
Egle Marmelos (glutinous and tenacious matter).
Artocarpus hirsuta (juice).
A. integrifolia (juice).
Balsamodendron Roxburghii (gum-resin).
Bauhinia retusa (gum).
Borassus flabelliformis (juice).

Crataeva religiosa (fruit).
Dichopsis elliptica (gum).
Euphorbia Cattimandoo (milky juice).
E. Royleana (juice).
Feronia Elephantum (gum).
Tamarindus indica (seeds).
Typha angustifolia (down of the ripe fruit).

Cultivation of Ipecacuanha.

CEPHAELIS
Ipecacuanha

sternutatory Boiled to a paste and applied to the cheeks, it is employed in the cure of tooth ache" (*Murray*).

Special Opinions.—§ "*Nak chikni*, sulphur, vinegar, and the leaves called *chitta*, mixed together, are used for pityriasis versicolor" (*Surgeon-Major C. W. Calthrop, Morar*). "It is used for hemicrania" (*Surgeon-Major J. Robb, Ahmedabad*).

CEPHAELIS, *Swartz.*; *Gen. Pl.*, II., 127.

Cephaelis Ipecacuanha, *Rich.*; *Fl. Br. Ind.*, III., 178; *Bot Mag.*, [t. 4063; RUBIACEÆ

IPECACUANHA ROOT, *Eng.*; RACINE D'IPECACUANHA ANNELEE, *Fr.*, BRECHWURZEL, *Germ.*

Syn.—C. ERETICA, *Pers.*; CALLICOCCA IPECACUANHA, *Brok.*; IPECACUANHA OFFICINALIS, *Arruda*

References.—*Kurs, For. Fl. Burm.*, II, 5; *Gamble, Man. Timb.*, 219; *Pharm. Ind.*

Ind., 543; (

1873, 233,

Papers, 343

Ag. Hort. Soc., Vol. V, p. 47.

MEDICINE.

916

CULTIVATION.

creasing costliness of the drug, have occasioned active measures to be taken for attempting its cultivation in that country. Though known for several years as a denizen of botanical gardens, the ipecacuanha plant has always been rare, owing to its slow growth and the difficulty attending its propagation.

"With
cult has
The first
had been

standing every care, the plants could not be made to thrive. Three plants, which had been sent to the Rungbi plantation in 1868, grew rather better, and by adopting the method of root propagation, they were increased by August 1871 to 300. Three consignments of plants, numbering in all 370, were received from Scotland in 1871-72, besides a smaller number from the Royal Gardens, Kew. From these various

conditions as regards sun and shade, but thus far with only a moderate

Dr. King reported to the Director of the Royal Botanic Gardens, Kew, in 1877, that he had distributed plants from the Calcutta Botanic Garden to Ceylon, Singapore, Burma, and the Andaman Islands, and also stated

CENTIPEDA
orbicularis

White Behen

Resinous

Non-resinous
909

class are too numerous to be mentioned here. The reader is referred to the list given in *Spons' Encyclopædia*, pp 626-627

CENCHRUS, Linn, Gen Pl, III, 1105

Cenchrus catharticus, Del, Duthie, Fodder Grasses, 15, GRAMINEÆ

Syn.—C ECHINATUS Rich.

Vern — *Bhurt*, HIND, *Dhaman*, argana N W P, *Basla*, led *lapla*, *bhort*, PB, *Bharbhunt*, JEPPORE, *Bharout*, AJMER, *Kukar*, BANDA

References — *Stewart*, Pb Pl, 252 *Aitchison Cat Pb Pl*, 163 *Murray*, Pl and Drugs, Sind, 10 13, *Duthie*, List of Grasses, N W P, 9

of the

FODDER
910

ritious
(the)
The

911

C. montanus, Nees.

This fodder grass is known as the *anjan* and *dhaman* in the Panjáb, and is considered by some one of the most nutritious of grasses and makes good hay

912

CENTAUREA, Linn, Gen Pl, II, 477

Centaurea Behen, Linn, COMPOSITÆ

THE WHITE BEHEN OR WHITE RHAPONTIC

Vern — *Bahman safaid* *suffaid bahman* HIND, BOMB; *Behen* (or

Habitat — A native of the Euphrates Valley. The root is largely imported into India, reaching Bombay from the Persian Gulf. It is always to be found in native druggists' shops

CENTIPEDA, Lour, Gen Pl, II, 430

913

Centipeda orbicularis, Lour, Fl Br Ind, III 317, Wight, Ic, [t 1610, COMPOSITÆ

Syn — ARTEMISIA STERNUTATORIA, Roxb Fl Ind Ed CBC 600

in sea

MEDICINE

Seeds

914

Leaves

915

Medicine — T

Hindus also the p

India, but the dry

ed in the druggist

dered LEAVES are used in affections of the head, such as colds, &c, as

C. 915

Cultivation of Ipecacuanha.

CEPHAELIS
Ipecacuanha.

conditions as to soil, moisture, and shade We have not even now a CULTIVA-
TION.

tropical It may, therefore, be found necessary to afford the plants

however, fortunately not been realized, and the drug is now obtainable at pretty much the same price as twenty years ago

In *South India* cultivation seems more hopeful than in Sikkim The late Mr Melvor, in May 1870, planted a few Ipecacuanha plants in the Botanic Gardens at Barhyār These succeeded fairly well, but in 1881 82, Mr Lawson, the present Superintendent of the Botanic Gardens, reported that he did not think the plant could be there grown as an article of commerce Later on, he seems to have attained more confidence in the pos-

been made above, that gentleman says of the South Indian experiments

could not produce the drug in any quantity at the usual market rate (from 4 to 5 shillings per pound), at which it can be bought in London

In an official communication dated May 1887 Dr Bidie writes hopefully

PROPAGA-
TION.

product. There are doubtless, however, many other similar regions where it might be grown The plant grows slowly, and has little in it to attract the attention of the cultivator, so that it may be doubted when private enterprise may be prepared to relieve the Government of its present

OEOPHAELIS
Ipecacuanha
Cultivation of Ipecacuanha.
**CULTIVA-
TION.**

to prevent the culti-
 opean planters The
 s, besides, little calcu-
 seedlings, and in 1870-71
 Some of these were culti-
 sent to Madras. Of the
 the higher regions of the
 Night hills were not found to be suitable. About this stage the Bom-
 bay Government became anxious that a consignment of plants should
 cultivation at the Cinchona planta-
 definite consignment of Messrs
 Mr W. Walton of the Cotton De-
 ase, under the care of that gentle-
 which Dr King, in 1871, reported as
 thy condition. These were sent to
 eral
 both
 ten-

the
 writer has been permitted to peruse, it would appear that the process of

them "The recent success in propagating has been entirely due to the
 discovery that this plant, unlike most others, can be propagated freely
 the plant's growth, materials
 y Propagation has all along
 and at an elevation of about
 have naturally been confined
 nts, so as to get a sufficiently
 large stock for experiment, with the view of determining the conditions

of plants have been put out at different elevations and under different
 C. 916

Cultivation of Ipecacuanha.

CEPHAELIS
Ipecacuanha.

CULTIVA-
TION.

conditions as to soil, moisture, and shade. We have not even now a

"I
a
tw
r

in charge of them the advisability of growing I

pr
to
(fr

PROPAGA-
TION.

product. There are doubtless, however, many other similar regions where
it might be grown. The attention of the
enterprise may be attracted
private
present

CEPHAELIS
Ipecacuanha.
Medicinal properties of Ipecacuanha.
PROPAGA-
TION.

efforts. Dr. King, in his paper read before the Agri-Horticultural Society, indicates clearly the peculiarities and necessities of the plant, and in his more recent communication (the official papers referred to above) he reiterates more strongly the same opinions. "There can be no doubt that the occurrence of a distinctly marked cold season is disadvantageous to the growth of Ipecacuanha. I sent plants of it for trial to the Andaman Islands and Singapore, both being localities where there is no cold season. But at neither place has the cultivation been much of a success. I had an opportunity of seeing, in the Singapore Garden, during the year 1879, the Ipecacuanha plants which I had sent from Calcutta, a year or two previously. And contrary to my expectations, I found them growing very indifferently. The plants sent to the Andamans I have never seen, but I understand that they did not come to much."

Large numbers of plants have been freely distributed to private cultivators, but it may be concluded that it still remains to be demonstrated whether or not the medicinal properties are preserved in the Indian cultivated stock. These may improve as in the case of some of the Cinchonas, but on the other hand, they may decline, so that it must be concluded Ipecacuanha in India is even now but in its most early experimental stage.

Medicine

MEDICINE.
Root.
917

The treatment of this disease by large doses of Ipecacuanha (grs. xxx to grs. lx), of late years re-introduced, has been found most effectual. In diarrhoea, and in some forms of dyspepsia, especially when connected with functional derangement or torpidity of the liver, it acts beneficially. As an expectorant it is in common use in catarrhs, chronic bronchitis, asthma, phthisis, the early stages of hooping-cough, &c. In hæmorrhages, especially in uterine hæmorrhages and in menorrhagia, it has proved an effectual remedy. For removing crude and indigestible matter from the stomach, Ipecacuanha acts with certainty and safety as an emetic, without inducing nearly the same amount of subsequent depression that follows tartar emetic; it is especially adapted for childhood and for persons of a delicate constitution. As a counter-irritant (2 drs. of powdered Ipecacuanha incorporated with 2 drs. of olive oil and 4 drs. of lard, rubbed into the skin for a few minutes, once or twice daily), it has been advanced

CHEMISTRY.
918

the alkaloid, which, taken internally, is a potent emetic.

C. 918

341-
are
of a
to

Medicinal properties of Ipecacuanha

CEPHAELIS
Ipecacuanha.

CHEMISTRY.

"Emetine, discovered in 1817 by Pelletier and Magendie, is a bitter substance with distinct alkaline reaction, amorphous in the free state as well as in most of its salts, we have succeeded in preparing a crystallized hydrochlorate

"The root yields of the alkaloid less than 1 per cent, the numerous higher estimates that have been given relate to impure emetine, or have

$\text{C}_{20}\text{H}_{30}\text{N}_2\text{O}_5$,
found in 1877

1 bark of the

solution containing but
be

of
1 a
to
then
and
will
attach-
satu-
on of
may

as a cathartic

"If the wood, separated as exactly as possible from the bark is used

res
anc
anc

p

Special Opinions — § "Applied locally to bites of venomous insects and scorpions" (Surgeon-Major C. W. Calthrop, Morar) "With out door patients suffering from dysentery, it was found to be most unsuited and inconvenient

used with much benefit

pill, and given every three hours, when for a malarious origin, quinine one grain to each pill was added" (Honorary Surgeon Peter Anderson, Madras Presidency) "In 3 gr doses it is a most efficient calmative and sedative in delirium tremens" (Surgeon-Major W. Farquhar, Ootacamund)

CEPHALOSTACHYUM
capitatum.

Coccinia Indica.

CEPHALANDRA, Schrad.; *Gen. Pl.*, I., 827.

- 919 *Cephalandra indica*, Naud ; *Fl. Br. Ind.*, II., 621 ; *Wight*, III.,
 1. 105 ; **CUCURBITACEÆ.**

bung, tsa-tha-khwa, BURM ; *Kóváká*, SING

References.—*Ro* 128 ; *Dals* 63

Cat. Pl., I., 128 ;
Ro 128 ; *Dals* 63

1 ; *Lisboa*,

Habit

Medic

plant is
 metallic

MEDICINE.

Juice.

920

used by preparations prescribed by them in diabetes." "The expressed juice is directed to be taken in doses of one *tola* along with a pill, every morning." (*U C Dutt, Mat Med Hind*). The root, according to Moodeen Sheriff, is sold as a subst

in the bazars of Southern India and are useful as a colouring ager

Root.

921

the essential oil. "The root when which hardens into a reddish gum on drying, and is very astringent, but not bitter like the fruit" (*Dymock*). "The bark of the root, dried and reduced to powder, is said to act as a good cathartic, in a dose of 30 grains" (*Medical Topography of Dacca*, 58). "The LEAVES, mixed with *ghí*, are applied as a liniment to sores. The whole plant, bruised and

Leaves.

922

ered cummin seeds, is used) "The leaves he plant internally in

gonorrhoea" (*Balfour*). "In the Concan the green fruit is chewed to cure sores on the tongue" (*Dymock*).

FOOD.

Fruit.

923

Food—"The oblong fruit, about 2 to 2½ inches long, green when young, scarlet-red when ripe, fleshy, smooth, is eaten both raw and cooked. The ripe fruit is sweet" (*Lisboa*). The fruit is one of the commonest of native vegetables (*Dymock*). It is eaten fresh when ripe and cooked in curries when green (*Roxb*)

924

Cephalocroton indicum, *Beddome*, 261 ; **EUPHORBIACEÆ.**

A common tree in the moist forests of South India (altitude 1,500 to 4,000 feet) ; yields a timber useful for building purposes.

CEPHALOSTACHYUM, *Munro* ; *Gen Pl*, III., 1213.

(See Vol. I., B 69, No. 9.)

925

Cephalostachyum capitatum, *Munro* ; **GRAMINEÆ.**

VERA.—*Gobia, gopi*, NEPAL, *Payong*, LEPCHA ; *Silli, sullea*, KHASIA.

Reference—*Gamble, Man. Timb*, 429.

Habitat—Found in Sikkim and the Khásia Hills.

C. 925

Wax.	CERA alba.
<p>Food — This semi-scandent and often gregarious bamboo, on flower- the natives in times of to 30 feet long, strong. with internodes about 2½ feet, thin, yellow, used for bows and arrows by the Lepchas. It flowered in Sikkim in 1874 (Gamble)</p>	<p>FOOD. Grain. 926</p> <p>TIMBER. 927</p>
<p><i>Cephalostachyum latifolium</i>, Munro</p>	<p>928</p>
<p>Reference — Gamble, <i>Man Timb</i>, 429</p>	
<p>Habitat — A species with large leaves, found in Bhutan.</p>	
<p><i>C. pallidum</i>, Munro, Kurr, <i>For Fl Burm</i>, II, 563</p>	<p>929</p>
<p>Vern. — <i>Beti</i>, Ass.</p>	
<p>Reference — Gamble <i>Man Timb</i> 429</p>	
<p>Habitat. — A bamboo with shrubby stems. It grows in the Mishmi</p>	
<p>Hills and in Ava</p>	
<p><i>C. pergracile</i>, Munro, Brandis, <i>For Fl</i>, 567</p>	<p>930</p>
<p>Vern. — <i>Tin-wa, kengwa</i> Burm</p>	
<p>References — Kurr, <i>For Fl Burm</i>, II, 564, Gamble, <i>Man Timb</i>, 429</p>	
<p>Habitat — A bamboo common in upper mixed forests of Burma, often</p>	
<p>gregarious. It has stems often 40 to 50 feet long</p>	
CERA.	
<p><i>Cera alba</i> and <i>flava</i>.</p>	<p>931</p>
<p>Ve.</p>	
<p>References — <i>Pharm Ind</i>, 278, Moodeen Sheriff, <i>Supp Pharm Ind</i>, 97; <i>Ainslie Mat Ind I</i>, 470, <i>Bidie Cat Raw Prod</i>, Paris Exb. 64, Baden Powell, <i>Pb Prod</i>, 166; <i>Fleming, Med Pl and Drugs</i>, Asia tic Res, XI, 195</p>	
<p>Description. — The prepared Honeycomb. Occurs in masses, firm, breaking with a granular</p>	
<p>light. Occurs in circular</p>	
<p>not urticuous to the touch. It does not melt under 130° F. (<i>Pharm.</i> <i>Ind</i>)</p>	
<p>Medicine — Honey is emollient and slightly laxative, and is often</p>	<p>MEDICINE.</p>
	<p>932</p>
<p><i>Ind</i>) For further information see Bees, also Wax</p>	
<p>Special Opinions — § 'The oil is used as a liniment and is of great value</p>	
<p>in muscular and chronic rheumatism' (<i>Surgeon Major A S G Jaya-</i> <i>kar, Muskat, Arabia</i>)</p>	
	<p>C. 932</p>

CERATONIA

Siliqua

The Carob Tree.

Ceramic Manufactures, *see* Earthen-wareCerasus cornuta, *Wall*, *see* Prunus Padus, *Linn.*CERATONIA, *Linn*, *Gen Pl.*, I, 574

933

Ceratonía Siliqua, *Linn.*, *DC Prodr*, II, 486, LEGUMINOSÆ.THE LOCUST TREE, THE CAROB TREE, ST JOHN'S BEAN, OR BREAD OR LOCUST BEAN, ALGARODA of Spain, CARRUBIO, *It*, CARUBA, *Ger*Vern — *Kharnub*, *kharnáb nubti* (the pods), *Ps*; *Kharnub shámi* or *kharnub nubti*, *ARAB*References — *Roxb*, *Fl Ind*, Ed C B C, 361, *Brandis*, *For Fl*, 166; *Gamble Man Timb*, 133 145, *Dals & Gids*, *Bomb Fl Suppl*, 28; *Cult* 364; *nson*, *omb*, *asury*, *ndia*,

Habitat — A slow-growing, evergreen tree, indigenous in Spain and

CULTIVA-
TION.

934

Cultivation of the Carob.

CERATONIA
Siliqua.

In the Panjáb, considerable quantities of seed have been sown from as early as 1844, in the districts of Panipat, Gurgaon, Rohtak, and Delhi,

CULTIVA-
TION.
935

(Stewart, *Fb Pl.*, 63) Mr. Ricketts was of opinion that the seeds should be well soaked before planting, and the trees when planted out should not be too far from each other to ensure their fruiting.

In Madras, the experiments were made in various localities, but the general result was anything but satisfactory. The seeds did not germinate in some cases, and in others, the seedlings soon died off.

936

In Bombay and Sind—"During the last two years, District Forest Officers in the Bombay Presidency have been engaged in carrying out experiments with carob seed, but the results do not appear to have been very promising. In Sind the Conservator states that all the plants were protected by mats from the frost during the cold season, and adds that when once these plants have established themselves in the soil, they should be able to exist without artificial irrigation or protection; at present they are too small, and it would be premature to express an opinion as to their flourishing in Sind or not. The Superintendent of the Economic Garden at Haidarabad, Sind, also states that, though the plant will grow, the slowness of growth

937

mental
the Pri
died, a
at Poc
this tri

tree in the *Woda gardens* about 71 lb of fairly good fruit were obtained in May last year, and the crop would have been heavier if protected from

938

as the
whole t
Met
not use

1 condition.

Food.—The pods, full of sweet pulp, are a good source of food in the Mediterranean and have been imported into the Panjáb and other parts of India. They form an important constituent of the "husks" of the "John the Baptist."

ey are said by
ringent. The
pectoral, and
les to them as

MEDICINE.
Pods.
939

FOOD.
Pods.
940

CERBERA
Odollam

The Carob Tree.

In the *Treasury of Botany* occurs the following account of Carob pods as a food stuff "These pods contain a large quantity of agreeably-flavoured, mucilaginous, and saccharine matter, and are commonly employed in the south of Europe for feeding horses, mules, pigs, &c, and occasionally, in times of scarcity, for human food. During the last few years, considerable quantities of them have been imported into England and used for feeding cattle, but although they form an agreeable article

price, and were used by singers, who imagined that they softened and cleared the voice. By fermentation and distillation, they yield a spirit which retains the agreeable flavour of the pod." Professor Church in *Food-Grains of India* (p. 170) states that "The nutrient ratio is here about 1 : 8.5, and the nutrient value 68. As sugar, pectose, gum, &c, occupy the place of starch in these pods, the starch equivalent cannot be

TIMBER.

941
DOMESTIC.
Seeds.

942

Carpenter work (Odollam)

CERBERA, Linn, Gen Pl, II, 699

Cerbera manghas, Linn, see *Tabernaemontana dichotoma*, Roxb,

[APOCYNACEÆ

943

C. Odollam, Gaertn, Fl Br Ind, III, 638, Wight, Ic, t. 441

Syn—C. LACTARIA, Ham, TANGHINIA ODOLLAM, LACTARIA, and LAURI FOLIA, Don

Vern—Dabur, dhakur, BENG, Kada ma, kat arali, kadalalai, kadu, TAM, Odallam, MALA, Gen kaduru, SING, Ka lwah, BURM

References—Roxb, Fl Ind, Ed C B C 232, Brandis, For Fl, 322, Kura, For Fl Burm, II, 171, Gamble Man Timb 252, Thwaites, En Ceylon

FIBRE.
Bark.

944

OIL

Seeds.

945

MEDICINE.

Sap.

946

Leaves.

947

Habit

C 947

Cerbera : The Yeast Plant.

CEREVISIÆ
Fermentum.

MEDICINE.

Nut.
048
Fruit.
040
Bark.
950TIMBER.
951DOMESTIC.
Ordeal Nut.
952

gative.

ally used for firewood

Domestic Uses.—The poisonous JUICE of the fruits was formerly used in Madagascar as an ordeal in cases of suspected crime or apostasy (*Kew Cat.*, 96)

Cerbera Thevetia, *Linn*, see *Thevetia nerifolia*, *Juss.*

CEREALS.

953

the reader is information, such as the into Cereals or Pulses, such as buckwheat, amarantus, &c.

from the
cereals
RN, and
parately.

CEREVISIÆ FERMENTUM.

Cerevisiæ Fermentum.

954

YEAST PLANT OF TORULA CEREVISIÆ.

Reference.—*Pharm. Ind.*, 262

The history of yeast is replete with interest, even although many of the details of the action of the plant in the process of fermentation are

955

undant
ated to
action
agent

with the sugary liquid. This must be viewed as a closely allied phenomenon to the effect of sulphuric acid on starch, contact converting the latter into sugar, while the acid itself remains unchanged in quantity or

CEREVISIÆ
Fermentum.

The Yeast Plant

chemical nature. In the process of beer-brewing two manifestations of the same kind are met with. The grain from which the beverage is to be prepared is first moistened either with hot water or by being placed in a warm confined atmosphere. As the result, it sprouts or germinates. The chemistry of this action consists in the fact that in a warm moist atmosphere the simple contact of a substance known as *diastase* with the starch of the grain converts the latter into sugar. Diastase may be defined as a transformed condition of gluten produced within the seed during the first stage of germination, and no sooner is the diastase formed than it immediately commences to act upon the insoluble starch. This is a wise provision of nature. The embryo plant is imbedded in a mass of starch. The base of the embryo contains gluten, but both starch and gluten are insoluble, and cannot be transformed into the structure of the young plant.

new substance is rapidly absorbed, and for the first period of its existence the infant plant feeds upon the food stored up for it within the seed. It

when the diastase completes its action on the still insoluble starch. It has been found that for every 100 parts of starch, in good malt, 1lb of diastase is produced, but that quantity will suffice to convert the starch of 1,000lb

brewer filters the wort, for the boiling has not only killed the diastase, but has also acted upon all the other soluble matters.

956

nourishment these minute plants take has never been clearly established,

in some respects better than the beers that used formerly to come to this country in such large quantities. The yeast is killed by the process of heating to 60°. In the brewing of beer only about a quarter of the fermentable substance is converted into alcohol, the remainder giving the

or *Torula Cerevisiæ*CEREVISIÆ
Fermentum

sweet flavour to the beverage. The yeast lives and increases in the fermenting liquid, but appears to abstract nothing from it, and just as contact of diastase has changed starch into sugar, so contact of yeast with sugar produces alcohol.

It has already been said that there would appear to be other sub-

957

958

and distilled. The flowers are placed in earthen vessels and mixed up |

for future use, having discovered that if not washed out these vessels |

Sarami (lur), to make the beverage intoxicating. According to some |
authors, an alcoholic beverage is prepared from the juice of *Calotropis* |

FERREVISIÆ
Fermentum

The Yeast Plant.

959

shown that the substances indicated are after all only flavouring ingredients or at most auxiliaries to fermentation; but in that case the true fermentum is the sweet liquids, since, with the to, no other instance is known

960

wood 20 feet in length and 3 feet in thickness is hewn out into a large trough. This is placed in the centre of the village, constituting the communal brewery. A large quantity of water is poured when on the trough beer from rice. A log of

961

Afghanistan from raisins. But apparently wheat and barley are but rarely used for this purpose, the liquor from the former being called *Madulika* and from the latter *Kohala*.

962

In India the favourite beverages are prepared from the juices of trees, chiefly palms (*Vartuni*), or from sugar-cane (*Sidhu*). For this purpose the juice is extracted from the cocoanut, the date, the palmyra, *Caryota urens*, and the *nim* tree. Fermentation is generally set up in these beverages by means of fermentation seed. This consists of rice saturated in a former fermentation, the grains of rice retaining apparently the germs of the yeast plant. Yeast from the *tari* beverage is largely used

MEDICINE
963

used as a poultice. In India, where yeast is rarely procurable, the toddy (*tari*) poultice, in a great measure, answers the purpose. (*Pharm Ind.*; see also the fermentation seed of *Borassus*, B. 689)

C. 963

The Mangrove.

CEROPEGIA
Arnottiana.

CERIOPS, Arn.; Gen. Pl., I, 679.

Ceriops Candolleana, Arnott, Fl. Br. Ind., II, 436; Wight, Ic,
[t. 240; RHIZOPHOREÆ.

THE MANGROVE.

Vern.—Kirrari, kiri, chauri, SIND; Gorán, BENG; Mada, AND.

References.—Brandis, For Fl. 218, Kura, For Fl. Burm., I, 448, Bed-
domé, Fl. Sylva Anal., Pl., XIII, Fig 5, Gamble, Man. Timb., 176,
Thwaites, En. Ceylon Pl., 120, Aitchison, Cat. Pb. Pl., 50, Murray,
Fl. and Drugs, Sind, 190

Habitat.—A small, evergreen tree, met with on the muddy shores and

964

DYE.
965TAN.
966MEDICINE.
Plant.

967

Bark.

968

Shoots.

969

TIMBER.

970

Litter for

Cattle.

971

972

decoction of the BARK is used to stop hæmorrhage, and is applied to
malignant ulcers. On the African coast, a decoction of the SHOOTs is
used as a substitute for quinine.

Structure of the Wood.—Red, hard, weight, 63lb per cubic foot. Used
in Sind for the knees of boats and other similar purposes; in Lower
Bengal for houseposts and for firewood

Domestic Uses.—The bark is used as a litter for cattle.

C. Roxburghiana, Arnott; Fl. Br. Ind., II, 436.

Vern.—Garán or Ghorán, BENG; Kabaying, kyadaing, ka-pyaing, BURM.

References.—Kura, For Fl. Burm., I, 448; Gamble, Man. Timb., 176;
McCann, Dyes and Tans, Beng., 133, 158, 458.

Habitat.—A large shrub of the coast of Chittagong, down to Tenas-
serim (Kura)

TAN.

BARK.

973

DYE.

BARK.

974

cloths (McCann)

Structure of the Wood—Weight of the wood, 46lb per cubic foot.

TIMBER.

975

976

CERIUM.

This metal is used medicinally in India. Minerals supposed to contain
it have been collected in the Karnal district, in Madras, and in Nepal
(See Ball's Econ. Geology).

CEROPEGIA, Linn.; Gen. Pl., II., 779.

Ceropegia Arnottiana, Wight; Fl. Br. Ind., IV., 74; ASCLEPIADÆÆ.

Vern.—Uta-long, BURM.

977

CHÆTOCARPUS
castaneæcarpus

Ceropegia Iceland Moss

Reference — *Balfour Cyclop*

Habitat — Grows in Khásia Mountains, Burma, and Tenasserim

978

Ceropegia bulbosa, *Roxb*, var *esculenta*, *Fl Br Ind*, IV, 67,
[*Wight, Ic*, t 845]Vern — *Khappar kadu*, HIND, *Patalatum bari* BOMBReferences — *Roxb Fl Ind*, Ed C B C, 250; *Dals & Gibs Bomb Fl*
153; *Voigt Hort Sub Cal* 534; *Dymock Mat Med W Ind* 2nd
Ed, 526; *Isisboa, U Pl of Bomb*, 165; *Balfour, Cyclop*

Habitat — Met with in the Panjáb and in the Bombay Presidency

Food — TUBERS and LEAVES are used as pot-herbs in Multan and Sind
Shepherds are fond of eating the tubers, which they consider to be tonic
and digestive "Every part of this plant is eaten by the natives, either
raw or stewed in their curries The fresh roots taste like a raw tur-
nip" (*Roxburgh*)FOOD
Tubers

979

Leaves

980

Roots

981

982

C. tuberosa, *Roxb*, *Fl Br Ind* IV, 70Syn — *C. ACUMINATA*, *Dals & Gibs*, 1c not of *Roxb*Vern — *Khappar kadu*, BOMB, *Pátál tumbdi*, MAR *Commumadu*, TELReferences — *Roxb, Fl Ind*, Ed C B C, 251; *Dals & Gibs Bomb*
Fl 153; *Dymock Mat Med W Ind* 436; *Murray, Pl and Drugs*
Sind 162; *S. Arjun Bomb Drugs*, 85Habitat. — Met with in the Deccan Peninsula from the Konkan south-
wards

MEDICINE

Tubers

983

appl cable to both plants and perhaps to one or two other species such
as *C. juncea* and *C. acuminata*Cetaceum, see *Physeter macrocephalus*, *Linn*, MAMMALIÆ

984

Cervidæ, the famly of the deer of interest economically for their antlers
and their skins See 'Horns' and also 'Skins'

CETRARIA.

985

Cetraria islandica, *Achar*, LICHENES.

ICELAND MOSS

References — *Pharm Ind* 258; *Flück & Hanb Pharmacog* 737,
O'Shaughnessy Beng Dispens, 672

Medicine — Imported into India and sold in chemists' shops

MEDICINE

986

Cevadilla or *Sabadilla*, see *Asagraea officinalis*, *Lindl*, LILIACEÆCeylon Moss, see *Gracillaria (Pilocaria) lichenoides*, *Greville*, ALGÆCHÆTOCARPUS, *Thw Gen Pl*, III, 323

987

Chætocarpus castaneæcarpus, *Thw*; *DC Prodr*, XV 2, 1127,
[EUPHORBIACEÆ]Vern — *Bulkakra* BENG, *Palakuna, sadavaku*, TAM, *Hedbaka, hálak-
waka* SING

C 987

Chara and Nitella.

CHARA
involucrata.

References — *Kurz For Fl Burm*, II 429 *Gamble Ma: Tr 16*, 366,
Thwaites En Ceylon Pl 275, *Trimen System. Cat*, Ceylon Pl, 82

Habitat — A moderate-sized tree, found in the Khasia Hills, Eastern Bengal, Burma, the Andaman Islands, and Ceylon

Structure of the Wood — Light red, moderately hard, close-grained, weight 58 lb per cubic foot, used in Ceylon for building

TIMBER.
988CHAILLETIA, DC, *Gen. Pl*, I, 341.

Chaillietia gelonioides, Hook, *Fl Br Ind*, I, 570; CHAILLETIACEÆ

989

Syn — *MOACURRA GELONIOIDES* Roxb, *Fl Ind*, Ed C B C, 254

Vern. — *Moakurra*, SILHET, BENG, *Balu nakuta*, SING

References — *Kurz For Fl Burm*, I, 230 *Gamble Man Timb*, 80,
Bedd. Fl. Sylv 59, *Thwaites En Ceylon Pl*, 79, *Trimen System Cat*
Ceylon Pl, 17, *Dalziel & Gibs, Bomb Fl*, 52 *Lisboa U Pl Bomb* 47

Habitat — A small subdioecious tree, commonly met with in the hilly eastern parts of Bengal and Silhet, in the forests of Madras, and in the Western Peninsula on the Ghats from the Konkan southwards, it is also met with in the moister parts of Ceylon up to an elevation of 3,000 feet

Structure of the Wood — This is one of the timber trees specially mentioned by Dr Lisboa in his *Useful Plants of the Bombay Presidency*, but very little of a definite character can be learned regarding the value of the wood

TIMBER.
990

Chalcedony, see Carnelian

Chalk, see Carbonate of Lime

CHAMÆEROPS.

991

Chamæerops Ritchieana, Griff, *Gen. Pl.*, III, 924; see *Nannorhops Ritchieana*, PALMÆ

Chamois Leather, see Leather & Skins.

Chamomile or Camomile, see *Matricaria Chamomilla*, Linn; COM-

Chanáy Kéléngu, see *Tacca pinnatifida* (?) [POSITÆ

Chank shells, see Shells and also Pearl Fisheries

CHARA.

Chara involucrata, Roxb, *Fl Ind*, Ed C B C, 648

992

Vern. — *Jangli pati*, HIND, *Jhang*, BENG (These vernacular names are applicable to all Charas, indeed to most submerged plants)

Habitat — There are a large number of species both of *Chara* and *Nitella* found in tanks and pools of water near Calcutta during the cold and hot season

Domestic use — *Atkinson* (Roxb), is employed use this finished in the

DOMESTIC.
Clarify
sugar.
993

CHARCOAL.

Timbers used for Charcoal.

Charcoal, see Carbon.

994

CHARCOAL, Timbers used for—

Abies Smithiana.
Acacia arabica.
A. Catechu
A. modesta
Adhatoda Vasica (gunpowder)
Albizia procera.
A. stipulata.
Anacardium occidentale
Anogeissus latifolia.
Betula cylindrostachys
Boswellia serrata.
Butea frondosa (gunpowder)
Cajanus indicus (gunpowder)
Callicarpa arborea.
Calotropis gigantea.
Casearia glomerata.
Cassia Fistula.
Castanopsis tribaloides.
Colebrookia oppositifolia (gunpowder)
Corchorus capsularis (gunpowder)
Cornus macrophylla (gunpowder).
Cynometra polyandra
Daphne mucronata (gunpowder)
Dillenia indica.
D. pentagyna.
Echinocarpus dasycarpus
Ehretia Wallichiana.
Elæocarpus lanceæfolius.
Eucalyptus Globulus
Eugenia tetragona
Euphorbia antiquorum

Excæcaria Agallocha
Ficus cordifolia.
F. infectoria.
F. religiosa.
Hippophae rhamnoides
Juniperus excelsa.
Lagerstroemia parviflora.
Mangifera indica
Mimosa rubicaulis (gunpowder).
Phyllanthus Emblica.
Pieris ovalifolia.
Pinus excelsa.
P. longifolia
Premna latifolia.
Prosopis glandulosa.
P. spicigera
Quercus Ilex.
Q. incana.
Q. semecarpifolia.
Q. spicata.
Rhododendron arboreum.
Salix tetrasperma (gunpowder)
Semecarpus Anacardium.
Sesbania ægyptiaca (gunpowder)
Sponia orientalis (gunpowder)
S. politoria (gunpowder)
Stereospermum suaveolens
Tamarix articulata.
Terminalia myriocarpa
T. tomentosa
Xylosma longifolium.

995

tion of *Anogeissus* and *Boswellia*, are not specially mentioned by writers on the subject as being good for fuel. These trees may, however, be added to the above list. Dr Schlich, in his note, estimated that to produce 15 tons of pig iron a day, 372,604 maunds of charcoal would be annually required, or say 1,800,000 maunds of firewood.

Chaulmugra, see *Gynocardia odorata*, *R. Br.*, BIXINÆÆ.

Chavannesia esculenta, *A. DC.*, see *Urceola esculenta*, *Benth.*

Chavica Betle, *Miq.*, see *Piper Betle*, *Linn.*, PIPERACEÆ.

C. officinarum, *Miq.*, see *Piper officinarum*, *C. DC.*

C. Roxburghii, *Miq.*, see *Piper longum*, *Linn.*

Chay root, see *Oldenlandia umbellata*, *Linn.*; RUBIACEÆ.

C. 995

The Wall-flower: The White Goose-foot

CHENOPODIUM
album.

Cheep, see Shells

Cheeronjee (chironji or chiraufi) oil, see *Buchanania latifolia*, Roxb.;

Cheese, see Ghl.

[ANACARDIACEÆ.]

Cheilanthes tenuifolia, Sw.; FILICES.Vern.—*Nanha, dodhari*, SANTAL.

The Reverend A. Campbell writes that the Santals prescribe a preparation from the roots of this fern for sickness attributed to witchcraft or the evil eye.

996

CHEIRANTHUS, Linn.; *Gen. Pl.*, I., 68.*Cheiranthus Cheiri*, Linn; *Fl. Br. Ind.*, I, 132; CRUCIFERÆ.

997

THE WALL-FLOWER.

V.

References.—*Stewart, Pb Pl.*, 13; *O'Shaughnessy, Beng. Dispens.*, 1861

Habitat.—Cultivated in gardens in North India, but is not indigenous; known as "Viole gialle," or yellow violets.

pur
ene

OIL.
Flowers.
998

MEDICINE.
Flowers.

999

Petals.

1000

Seeds.

1001

son, M D, *Vignor*).
"aphrodisiac" (*Surgeon J. Ander-*

CHENOPODIUM, Linn.; *Gen. Pl.*, III, 31.

1002

A genus of annual or perennial herbs, belonging to the Natural Order
CHENOPODIACEÆ.

too d or
Sty ssed.

There are about 50 species of the genus, met with in the world. These are distributed in all climates. India possesses seven species, with perhaps numerous varieties and cultivated forms of most of these.

Chenopodium album, Linn; *Fl. Br. Ind.*, V., 3; CHENOPODIACEÆ.

1003

THE WHITE GOOSE-FOOT.

SYN.—*C. VIRIDE*, Linn; *Roxb. Fl. Ind.*, II, 38.

C. 1003

Mexican Tea The Jerusalem Oak.

CHENOPODIUM
Botrys.

Domestic Uses —Baden Powell says that this plant is used in the Panjab "to clean copper vessels preparatory for tinning them"

DOMESTIC.
1008

Chenopodium ambrosioides, Linn, *Fl Br Ind*, V, 4.

1009

THE SWEET-PIGWEEED, MEXICAN TEA

Syn —*C. VALPINUM*, Wall, *AMBRINA AMBROSIOIDES*

Vern —*Herba Santa Maria* in Brazil In Chili this is known as *Culen*

References —*Dals and Gids*, *Bomb Fl Suppl*, 73, *Bent and Trim*, *Med Pl*, 216

racemes

Medicine —This is said to afford an essential oil to which the tonic and antispasmodic properties of the plant are attributed. It is commonly reported that this plant is used as a substitute for the officinal *C. anthelminticum*, having in a milder degree the anthelmintic properties of that plant. It is employed in pectoral complaints and enjoys the European reputation as a useful remedy in nervous affections, particularly chorea. *Officinal preparation an infusion*

MEDICINE.
Oil
1010

various species not being distinguished

Food.—This plant affords the Mexican tea.

C. Blitum, Hook f, *Fl Br Ind*, V, 5

Syn —*BLITUM VIRGATUM*, Linn

Vern —*Sundar* (J), *kupala* (C), Ps

References —*Stewart*, *Pb Pl*, 177; *Von Mueller*, *Extra Tropical Plants*

Habitat —North Western India. Kashmir, altitude 8500 feet and Stewart found the plant wild in the Trans-Indus at altitudes

FOOD.
1011
1012

its fruits furnish a red dye "

Food —Stewart remarks that "the extremely insipid FRUIT is sometimes mistaken by Europeans for a kind of strawberry, and which it much resembles. In Ladak the LEAVES are eaten as a pot herb "

DYE.
1013
FOOD.
Fruit.
1014
Leaves.
1015
1016

C. Botrys, Linn, *Fl Br Ind*, V, 4

THE JERUSALEM OAK

Syn —*C. ILICIFOLIUM*, Griff *Natul*, IV, 337

References —*Dals & Gids*, *Bomb Fl Suppl*, 73

Habitat —Temperate Himalayas from Kashmir to Sikkim at altitudes from 4,000 to 10,000 feet. Tibet 11,000 to 14,000 feet. Stewart says it occurs at Peshawar, and Dalzell that it was originally introduced into Bombay but has now gone wild. A weed of fields

Medicine —Reported to be used as a substitute for *C. anthelminticum* and to possess the same properties as *C. ambrosioides*. According to *U S Dispensatory* it has been used in France with advantage in catarrh and humoral asthma. The officinal preparation is an oil.

MEDICINE.
Oil
1017

C. 1017

CHICKRASSIA
tabularis

The Quinoa : The Chittagong Wood.

1018

Chenopodium murale, Linn ; *Fl Br. Ind*, V, 4.Vern — *Bátá, káránd, kharatua*, PsReferences.—*Stewart, Pb Pl*, 178

FOOD.

1019

Habitat — General in many parts of India from the Panjab to the Gangetic Valley, the Deccan, and South India.

1020

Food — Used as a pot-herb in the Panjab

C Quinoa, an American species, has once or twice been tried in India, but apparently with little success (*See Church, Food Grains of India*, p 110)Cherry, see *Prunus Cerasus*, Linn., ROSACEÆ.Chestnut, Horse, see *Æsculus indica*, Colebr (A 567), and *Æ. Hippocastanum*, Linn (A. 573) ; SAPINDACEÆ.Chestnut, Sweet, see *Castanea vulgaris*, Lam , CUPULIFERÆChestnut, Water, see *Trapa bispinosa*, Roxb, and *T. nutans*, Linn .
ONAGRACEÆCHICKRASSIA, A Juss , *Gen Pl*, I, 339

1021

Chickrassia tabularis, Ahr Juss ; *Fl Br Ind*, I, 568, *Beddome, Fl Sylat.*, t 9, MELIACEÆ

THE CHITTAGONG WOOD

Syn.—*SWIETENIA CHICKRASSIA*, Roxb, *Fl Ind*, Ed C B C, 370, C

Habitat — A large tree, native of the hills of Eastern Bengal, South

GUM.

1022

DYE.
Flowers.

1023

MEDICINE
Bark

1024

TIMBER

1025

It is used for ever
"The wood is
extensively used

C. 1025

The Chittagong Wood Chlorophytum

CHLOROPHYTUM
breviscapum

'Chittagong wood,' being imported from that district, though it is abundant in the mountainous parts of the peninsula. It is close grained

but tough and close grained, and, from its general situation, it is hardly known to the carpenter. It grows in the warmer parts of Ceylon" (*Balfour, Cyclop*)

Chicory, see *Cichorium Intybus*, *Linn*, COMPOSITÆ

China Root, see *Smilax china*, *L*, LILIACEÆ

Choinanthus albidiflora, *Thw*, see *Linociera albidiflora*, *Thw*

C zeylanica, *Linn*, see *Linociera purpurea*, *Vahl*, OLEACEÆ

Chireta, see *Swertia Chirata*, *Ham*, GENTIANACEÆ

Chloride of Ammonium, see *Ammonium chloride*.

Chloride of sodium, see *Sodium chloride*

CHLORIS, *Sw*, *Gen Pl*, III, 1165

Chloris barbata, *Swartz*, *Duthie, Fodder Grasses*, 53, GRAMINEÆ

1026

Syn — *ANDROPOGON BARBATUS* *Linn*

Vern — *Gandi* " " " " " " " " " " " "

jharna Pa

PUR Bārdiy

SOUTH INDIA

References —

371 *Dals* &

Murray Pl *Orings*, 51 & 12 *Dialie*, *Cat Kaw Prod*, *Paris Exh*, 76

grows

hey do

FODDER.

1027

1028

CHLOROPHYTUM, *Ker*, *Gen Pl*, III, 788

Chlorophytum breviscapum, *Dalz* in *Kew Journ*, II, 142,

1029

[LILIACEÆ

Vern — *Bimphol* *Sing*

References — *Dals & Gibz*, *Bomb Fl*, 252 *Thwaites*, *En Ceylon Pl*, 339, *Baker*, *Linn Soc*, XV, 321, *Treasury of Botany*, II, 1280

Habitat. — Frequent in the Malwan District, Bombay, in rocky situations. C Heynel *Baker*, a newly allied species met with in the southern and central parts of Ceylon, at no great elevation

C 1029

CHLOROXYLON
Swietenia.

The Indian Satin-wood.

MEDICINE
Balb.
1030

ascending the Himalaya to 5,000 feet in altitude. *C. nepalensis* occurs in the eastern sub-tropical Himalayas, while *C. arundinaceum* occurs on the sub-tropical Himalaya and on Parasnath in Behar, altitude 4,000 feet

1031

CHLOROXYLON, DC ; Gen Pl, I, 340
Chloroxylon Swietenia, DC, Fl Br Ind, I, 569; Beld., Fl Syl-
vat, t 11, Wight, Ic, t 56; MELIACEÆ.

THE INDIAN SATIN-WOOD

Syn.—SWIETENIA CHLOROXYLON, Roxb Fl Ind, Ed. C B C, 370

Vern.—Dhaura, bhirra, girya, HIND, Behru, biluga, bhavru, bheyl, URIYA, Behra, girya, behru, bhiru, bhirra bhirra C. P, Sengel sa's, AOL, Banarhal, KARWAR, Bhira, GOND, Bhirra BAIGAS, Hulda, bills, hardi, bheria, BOMS, Halda, bheria, MAR., Mudada, burus, purush-mudada marum, purus-burus, tummay, mudada, tummai pota-

burnte, BING

References.—Brandis, For Fl, 74 Gambel, Man Timb, 77, Thaxates, En Ceylon Pl, 61 Dals & Gibs Bomb Fl, 39, Voigt, Hort Sub Cal, 137 Dymock, Mat Med W Ind, 2nd Ed, 177 Drury, U Pl, 131 Cooke, Gums and Gum resins, 25, 115 Atkinson Gums and Gum resins, 34 Atkinson, Him Dist, 814 Lusbo, U Pl, Bomb, 4, Bal four, Cyclop, Treasury of Botany Ann Cal 22.

Habitat.—A moderate-sized, deciduous tree, found in Central and South India, and Ceylon Common in the forests of the Konkan, Decan, and Coromandel, flower in March

Gum.—"Satin-wood gum was contributed by Dr Cleghorn to the Madras Exhibition of 1855 The specimen in the collection from Salem (1873) refer tears, very lucent, bro ble in water mahogany

surface of the solution

"Another sample in the reference collection is from Ceylon, paler in colour, and in definite, rounded, shining, amber-coloured tears" (Cooke, Gums and Gum-resins, 25).

Dye—"Yields a yellow dye" (C. P. Gaz, '03)

Oil.—The tree yields a wood-oil (Beddom)

Medicine.—"The astringent BARK is prescribed sometimes by Hindû

DYE
1033
OIL
1034
MEDICINE
Bark.
1035
Leaves
1036
TIMBER
1037

56lb per cubic foot.

C. 1037

Garden Chrysanthemums.

CHRYSANTHEMUM.

SATIN-WOOD

very str
as it me
the bro
a ton, c
furnitur
of 8 to
kotties
district

kotties part of the satin wood cut is exported to Madras, where it is used for furniture and general building purposes" (*Indian Forester*, X, 1 38)

Chocolate nut and bean, see *Theobroma Cacao*, Linn, STERCULIACEÆ

CHONEMORPHA, Don, *Gen Pl*, II, 720

Chonemorpha macrophylla, G Don, *Fl Br Ind*, III, 661,
[Wight, *lc*, 1 432, APOCYNACEÆ

Syn — *ECMITES MACROPHYLLA* Roxb, *Fl Ind*, Ed C B C, 245

Vern — *Garbadero*, HIND, *Yokchounrik*, LEPCHA, *Harki*, SYLHET

References — *Brandis*, *For Fl*, 323, *Kurs For Fl Burm II*, 187
Gamble Man Timb, 261 *Dala & Gibs*, *Bomb Fl*, 146, *Voigt, Hort*
Sub Cal, 523, *Balfour, Cyclop*

Habitat — A large climber with milky sap, met with in North and East

the leaves of
and the roots
The *Flora of*

GUM
1030
MEDICINE
1040

British India alludes to that plant as a doubtful species

Chowli, or Chauli, see *Vigna Catiang*, Endl, LEGUMINOSÆ.

CHROMIUM AND CHROMITE.

The metal Chromium

of
in
suc
in
is 1

information see *DAVIS* *Leon Geology*, 331. *Mallet, Mineralogy*, 53.
Balfour's Cycl, 717

CHRYSANTHEMUM, Linn, *Gen Pl*, II, 424

There are three wild species belonging to this genus met with in Western Thibet and one in upper Sikkim—all alpine in their character, never occurring below 9000 feet The Chrysanthemums of Indian pharmacy are the two garden species

1041

1042

C 1042

CHRYSANthemum indicum.

The Common Garden Chrysanthemum.

1043

Chrysanthemum coronarium, Linn ; Fl Br Ind, III, 314, Bot
CHRYsanthemum [Mag, t. 1521 ; COMPOSITE.
Syn — C. Pinnatifidum, DC Prodr, VI, 62, CHRYsanthemum
Ve
CYNACEÆ
Pb 99, 183,
It is one of the Medicinal plants of India

almost naturalised in India, and to such an extent that Roxburgh viewed them as "natives of Bengal"

Medicine.—"The FLOWERS are stated by Dalzell and Gibson to form

MEDICINE
Flowers.
1044
Root.
1045

(Pharm Ind)

Garlands.
1046

an agent for opening the mouths of wounds" (Murray, Plants and Drugs of Sind)

Sacred Uses —"The beautiful yellow fragrant flowers of this plant are made into garlands and offered at the shrines of Vishnu and Siva" (Balfour)

1047

C indicum, Linn ; Fl Br Ind, III, 314 ; Bot Mag, t 327, 2042,
THE COMMON GARDEN CHRYsanthemum OF INDIA [2556

Syn — PYRETHRUM INDICUM, DC Prodr, VI, 62, CHRYsanthemum
INDICUM Willd in Roxb Fl Ind, Ed, C B C, 604
Vern.—Gul dāudi, HIND, a name applied, according to Roxburgh, to all
the varieties, Gendi, bāgdur (genda is the Hindustani for Tagetes erecta),
PB, Kalsang, LADAK, Chevati akurkura, BOMB ; Shevati, MAR ;
Akkara carum, TAM, Chamunti, TEL

C. 1047

Chrysanthemum. Fodder Grasses.

CHRYSOPOGON
aciculatus

References — *Roxb, Fl Ind, Ed C B C, 601, Clarke, Composita Ind, 145, Dals & Gibs, Bomb Fl Supp 48, Stewart, Pb Pl, 124; S Arjun, Bomb Drugs, 192, Baden Powell, Pb Pr, 358, Birdwood, Bomb Prod, 50*

Habitat.—Commonly cultivated in Indian gardens, and is in fact only

MEDICINE.
Flowers
1048

gonorrhœa

Sacred Uses.—The flower-heads are sacred to Vishnu and Siva.

Garlands.
1049

CHRYSOPHYLLUM, *Linn, Gen Pl, II, 653*

Chrysophyllum Roxburghii, *G Don, Fl Br. Ind, III, 535; Bedd, Fl Sylv, t 236, MELIACEÆ*

1050

THE STAR APPLE

Syn — C ACUMINATUM, *Roxb, Fl Ind, Ed C B C, 201*

W. & A. D. P. B. C. D. L. A. C. E. I.

Thwaites En Ceylon Pl 174, Dals & Gibs, Bomb Fl 138 Voigt, Hort Sub Cal, 340; Lisboa, U Pl Bomb, 88, Balfour, Cyclop

Habitat — An evergreen tree of Bengal, Burma, the Western Ghâts, and Ceylon

Food — FRUIT edible Roxburgh says "The fruit ripens in October,

FOOD
Fruit
1051

TIMBER
1052

general use (*Bomb Gas, LV, pl 1, 00*)

CHRYSOPOGON, *Trin, Gen Pl, III, 1135.*

Chrysopogon aciculatus, *Trin, Duthie, Fodder Grass, 39, GRAMINEÆ*

1053

Habitat — A small, coarse grass, growing on barren, moist pasture

Fodder.—Cattle do not seem to like it. Its thin, straight culms, 1 to 2 feet high, flower, and the small spikelets of awned, barbed, fruits which follow, are troublesome to those who walk through the grass, as they stick

FODDER.
1054

T

C. 1054

CHRYSANTHEMUM
indicum.

The Common Garden Chrysanthemum

1043

Chrysanthemum coronarium, Linn ; Fl Br Ind, III, 314, Bot

CHRYSANTHEMUM [Mag, t 1521; COMPOSITÆ.

Syn — C. PERSICUM, DC Prodr, VI, 62. PERSICUM, DC Prodr, VI, 62.

Ve

I

CYNACEÆ

References — Dals & Gids, Bomb Fl Supp, 48, Aitchison, Cat Pb
Pl, 77, Pharm Ind, 127; Mooden Sheriff, Supp Pharm Ind, 99,
Dymock, Mat Med W Ind, 371, Murray, Pl and Drugs, Sind, 183,
S Arjun, Bomb Drugs, 79, Drury, U Pl, 133, Balfour, Cyclop

" of the Med Ind

almost naturalised in India, and to such an extent that Roxburgh viewed
them as "natives of Bengal"MEDICINE
Flowers.

Medicine — "The FLOWERS are stated by Dalzell and Gibson to form

1044

Root.

1045

(Pharm Ind)

Garlands.

1046

of Sina)

Sacred Uses — "The beautiful yellow fragrant flowers of this plant
are made into garlands and offered at the shrines of Vishnu and Siva"
(Balfour)

1047

C indicum, Linn ; Fl Br Ind, III, 314; Bot Mag, t 327, 2042,

THE COMMON GARDEN CHRYSANTHEMUM OF INDIA [2556

Syn — PYRETHRUM INDICUM, DC Prodr, VI, 62. CHRYSANTHEMUM
INDICUM Willd in Roxb, Fl Ind, Ed, C B C, 604V. Roxburgh, toall
Tagetes erecta),
Shevati, MAR ;

C. 1047

Chrysanthemum Fodder Grasses

CHRYSOPOGON
aciculatus

References — *Roxb Fl Ind, Ed C B C 601*, *Clarke Compositæ Ind, 145*, *Dalsh & Gibs, Bomb Fl Supp 48*, *Stewart Pb Pl 124, S Arjun Bomb Drugs, 192*, *Baden Powell, Pb Pr, 358*, *Birdwood Bomb Prod, 50*

Habitat — Commonly cultivated in Indian gardens, and is in fact only

MEDICINE
Flowers
1048

calculus and also to remove depression of spirits Drury says the "natives of the Deccan administer the plant, in conjunction with black pepper, in gonorrhœa"

Sacred Uses — The flower heads are sacred to Vishnu and Siva

Garlands
1049

CHRYSOPHYLLUM, Linn, Gen Pl, II, 653

Chrysophyllum Roxburghii, G Don, Fl Br Ind, III, 535,
Bedd, Fl Sylv, t 236, MELIACEÆ

1050

THE STAR APPLE

Syn — C ACUMINATUM Roxb, Fl Ind Ed C B C 201

Vern — *Petakara* BENG, *Pithogarkh* ASS, *Hali, hali-maru* KAN, *Tarsu, tarsiphala* BOMB, *Tarsu, Mar, Lawulé* SING, *Thankya, than kya*

δ, 242,
Voigt,

Habitat — An evergreen tree of Bengal, Burma, the Western Ghâts, and Ceylon

Food — FRUITEDIBLE Roxburgh says "The fruit ripens in October,

FOOD
Fruit
1051

CHRYSOPOGON, Trin, Gen Pl, III, 1135

Chrysopogon aciculatus, Trin, Dulhie, Fodder Grass 39, GRAMINEÆ

1053

Syn — ANDROPOGON ACICULATUS Linn (f Ret.) Roxb, Fl Ind, Ed

Habitat — A small, coarse grass, growing on barren, moist pasture

TIMBER
1052

Fodder — Cattle do not seem to like it. Its thin straight culms, 1 to 2 feet high flower, and the small spikelets of awned barbed, fruits which follow, are troublesome to those who walk through the grass, as they stick.

FODDER.
1054

CICER
arietinum.

Fodder Grasses The Common Gram

to the stockings and produce until removed a pricking and itching sensation As soon as the spikelets appear cattle refuse to eat the grass

1055 *Chrysopogon coeruleus*, *Nees*, *Duthie, Fodder Grasses*, p 39

Syn — *RHAPHIS CERULEA* *Nees*

Vern — *Dhaulian* Pb *Khar*, SALT RANGE *Dhaura* SIWALIK RANGE, *Ghweia*, KUMAON, *Tigri*, BUNDELKHAND, *Palla paggar gadi*, CHANDA, *Jhingra ka jhara*, *khidi*, BERAR

Habitat. — A common grass on the hilly tracts of Northern India, usually on stony or sandy soils

Fodder — On the Siwalik range it is extensively used as fodder

FODDER.

1056

1057

C. gryllus, *Trin*, *Duthie, Fodder Grasses*, 40

Syn — *C ROYLEANUM*, *Nees* *ANDROPOGON GRYLLUS*, *Linn*

Reference — *Aitchison, Cat Pb Pl*, 176

Habitat — The plains and hills of the Panjab and N-W Provinces

Fodder — *Mueller* says it is a useful fodder grass in Australia

FODDER.

1058

1059

C. montanus, *Trin*, *Duthie, Fodder Grasses*, p 40.

Syn. — *C PARVIFLORUS*, *Benth*, *ANDROPOGON MONTANUS*, *Roxb*

Vern — *Ballak* RAJ

Habitat — The hilly parts of Northern India (Mount Abu)

Fodder — In Rajputana it is said to be viewed as excellent fodder, and the grain is also sometimes collected and eaten by the natives

FODDER

1060

Cicca disticha, *Linn*, see *Phyllanthus distichus*, EUPHORBACEÆ

Cicendia hyssopifolia, *W & A*, see *Enicostema littorale*, *Blume*,
[GENTIANACEÆ]

1061

CICER, *Linn*, *Gen Pl*, I, 524

Cicer arietinum, *Linn*, *Fl Br Ind II*, 176, *Wight Ic*, t 20

[LEGUMINOSÆ]

THE COMMON GRAM OR CHICK PEA, *CECE It* *GARBANZOS*, *Sp*

Vern — *Chola bāt*, *but kalai* *BENG* *Chana chunna* *HIND* *But*,
SANTALI *Channa chola*, *PB* *Chola chana* *RAJPUTANA*, *Chana*

or Chick Pea

CICER
arietinumthe *εβίβη* of Dioscoridesto a not
in the poc
"gram"

where it

kurti (Do

33) In I'

Phaseolus "

'Bengal gram' these terms are however, unknown in other provinces,
where the word "gram" is exclusively given to the pea of *Cicer*

History—The

the Greeks in Homer's

Romans as *Cicer* and the

that it was early known

Europe It is supposed

to pt from the very earl est

HISTORY

time of the

lik

du

int

an

Ar

Eu

Gh

fro

ground where we do not know whether it springs from a stock originally
wild or from cultivated plants' (DC, *Orig Cult Pl*)

CULTIVATION

CULTIVA-
TION*N W Provinces*—The varieties grown in the North Western Pro
vinces are classed as follows

N W P

Large

1062

Small

1063

Cabuli

1064

heaviest clay to the lightest loam, but it is found to prefer the former
It does not require so fine tillage as wheat and barley do, nor much
irrigation and a deep rather than well pulverised seed bed is all that is
necessary The plants bushy like the

The cost of

follows —

Ploughing (four times)

Seed (Sole)

Sowing

Reaping

Threshing

Cleaning

Rs a p

3 0 0

2 0 0

0 14 0

1 9 0

2 0 0

0 6 0

TOTAL

9 13 0

Fert

3 0 0

GRAND TOTAL

12 13 0

CICER
arietinum.

The Common Gram

CULTIVA-
TION.

The approximate average outturn for unirrigated land in the several divisions varies from 5 to 8 maunds per acre in the case of gram, and from 6 to 9 maunds in the case of gram barley and gram-wheat. For irrigated land the outturn is estimated at 12 maunds for gram alone,

C. P
1065

est return was in Narsinghpur, where 873lb to the acre were obtained, and the lowest, 237lb, in Chanda. Taking the mean of all the returns in the eleven districts the yield may be expressed at 557lb. In the Chanda Settlement Report, it is stated that two kinds of gram are grown—the grey and the white. It is remarked that gram is not a popular crop in the Wardah District.

BOMBAY.

Bombay—There are 692,295 acres under this pulse, and in Sind 34,166 acres. The crop experiments made in the Bombay Presidency reveal

Large
1066
Small
1067Kills weeds
Improves soil

The following extracts from the Bombay Gazetteers will be found

Justification
of mixed
cropsWheat and
Gram

plained of by European merchants is the consequence of either of two things—*1st*, the wilful purchase of such admixture, for the natives of India regularly eat the two grains mixed, and to meet this demand the Indian

seems every reason to suppose that a certain amount of wilful—one might almost say criminal—admixture of gram takes place in wheat sold as pure wheat. Such admixture is mainly, if not entirely, effected by the dealer not by the cultivator.

C. 1067

or Chick Pea.

CICER
arietinum.CULTIVA-
TION.is the most
district. It
either water
to be used

Hols.

1068

Dal

1069

Puran-poll.

1070

Phutanaz.

1071

manured and irrigated lands In Belgaum gram is known as *kadli* in

p. 151).

In the ...
grown
soils P_r
wheat.

PANJAB

1072

The amount of seed to the acre varies from 32 to 48 lb. Rain in March
and April, so beneficial for wheat, and indeed abundant rain or prolonged
cloudy weather at any
causes the plant to s_i

Red

1073

Black.

1074

White.

1075

Cabull.

1076

C. 1076

or Chick Pea

CICER
arretinum

what has been said, it may be inferred that adulteration of gram with wheat grain is more an accident than a necessity of the habit of mixed cultivation

CULTIVA-
TION

GRAM AS AN ARTICLE OF CATTLE DIET — In an address delivered before Gram recom-

country has always a much larger percentage of pulses in it than in Europe. The animals thrive admirably on such a diet, and the opinion may be advanced that where muscular strength is required a diet that

of oats and Indian corn. To obtain the indispensably necessary amount of albuminoids from an English diet the animal has to eat a greatly

scribed by Principal McCall of Glasgow, in which the tongue becomes paralysed. We be said that our and that it has tried to the extent. The writer has

CICER
arietinum.

The Common Gram

These remarks regarding anthrax have however, been made in this place mainly to prevent undue alarm, until Professor Wallace's suggestions regarding a possible connection between it and gram-feeding have been proved correct.

CHEMISTRY
1084

CHEMICAL PROPERTIES OF GRAM

Professor Church, in his *Food-Grains of India*, gives an interesting account of this pulse, but is in error in too prominently restricting the name *gram* to the forms of *Phaseolus Mungo*. This is the case only in the Madras Presidency, throughout the rest of India the terms black and green gram are practically unknown, the word *gram* signifying the pulse *Cicer arietinum*, although the term horse gram is sometimes applied to the pea of *Dolichos biflorus*. In Madras it might fairly well bear that name, since it takes the place of *Cicer arietinum* as a food for horses. The Professor gives a valuable table as the result "of nine analyses of the unhusked peas and of four analyses of the peas from which the husk has been removed,"

"COMPOSITION OF THE CHICK-PEA.

IN 100 PARTS.

	Husked	With Husk	In 1 lb Husked	
			Oz	Grs.
Water	11.5	11.2	1	367
Albuminoids	21.7	19.5	3	207
Starch	59.0	53.8	9	192
Oil	4.2	4.6	0	294
Fibre	1.0	7.8	0	70
Ash	2.6*	3.1†	0	182

* 1.1 of Phosphoric Acid.

† 0.8 of Phosphoric Acid

"The nutrient ratio in the unhusked peas is 1 : 3.3; the nutrient value is 84."

The unhusked peas are therefore more nutritious than the husked, and it may be concluded that the process of steeping them in water before

a high reputation.

TRADE AND PRICES

Very little can be learned regarding the internal trade in gram. It is extensively eaten by the natives in every part of the country, and there must therefore exist a very considerable internal trade in the pulse. The grain could be most conveniently obtained from Bombay, Karachi, or Cal-

TRADE.
1085

or Chick Pea.

CICER
arietinum.

The foreign trade is at present not very extensive. The following were the exports during the past five years :—

TRADE.

	Cwt.	R
1882-83	312,953	8,28,647
1883-84	392,694	11,99,796
1884-85	314,965	9,28,848
1885-86	338,129	10,74,771
1886-87	306,979	9,84,646

The exports in 1870 were only 23,171 cwt, valued at Rs4,900; but it

various Indian pulses The majority of these gentlemen agreed in

other.

Prices.—In a recent number of the publication issued by the Department of Finance and Commerce under the title of *Prices and Wages in India*, Mr. O'Connor has published tables which afford perhaps the most trustworthy data for arriving at a knowledge of the price of gram; his figures represent seers (2½) to the rupee. Mr. O'Connor's results of average prices may be thus summarised :—

PRICES.
1086

	I 1873 to '76.	II 1877 to '80.	III 1881 to '84.	IV 1873 to '80.
Madras	23 63	17'77	32'05	20 7
Bombay and Sind	17 06	11'47	18 45	14 27
Bengal	20 58	15'31	21'77	17'94
North-Western Provinces and Oudh	26 61	18 36	24'53	22 48
Panjab	30 04	18 29	26 7	24'16
Central Provinces	31'02	18 1	27'25	24 56

It would, perhaps, be unsafe to carry these figures further; but the mean of Column IV, might give the reader an average approximation of the retail price of gram in India. But it must not be lost sight of that "gram" as presently exported means more than the pea of *Cicer arietinum*, and includes (as perhaps do the above figures) pulses that have a lower value than the true gram.

CICER
arietinum

The Common Gram

PRICES

seers to the rupee in wh ch of course a larger quant ty for the sum men
t oned would mean cheapness and a less quantity dearness —

DISTRICTS	August 15th	November 15th	February 15th	May 15th
Mandla	45	42	40	40
Damoh	39	27	29 8	40
Sambalpur	15	19 8	19 8	
Wardha	20	22	21	24

The d fference between the prices at which the cultivators sell the pro-
duce of their fields to the dealers at harvest time and at other periods

BENGAL
1088

it is accordingly
Director of Agri
at 24 seers to the

rupee after harvest and 20 seers at other seasons taking a h gh ex
change these quantities would represent 48 to 40lb for 1s 5d

BOMBAY
1089

Bombay—The quotation has been given in one of the *Crop Experi-
ments* of 60 seers to the rupee or, at the rate of exchange adopted in the
preceding estimates 120lb for 1s 5d It is probable however, that this
figure is much too low and that the average price in the Western Presi-
dency bears a closer approximation to that given for the Central Provinces
and Bengal

PANJAB
1090

Panjab—In the Lahore d strict according to the *Gazetteer*, gram is
stated to be sold at 100lb to the rupee (= 1s 5d) In the Mooltan dis-
trict, the average price for the past 20 years is given as 60lb and in the
Jhelam district for the past 44 years as from 68 to 110lb according to the

N W P
1091

am is consider

DYE
1092

s fact is known

..

MEDICINE
Seeds
1093

Gram
Vinegar
1094

or Chick Pea.

CICER
arietinum.

afterwards published in the Records of the Bombay Government (xvi) MEDICINE.

peculiar to the dew ... further on at p. 63, he observes that the natives

... drug
is sold
in some
given
lied as
The fresh juice of the leaves
administered with success in
The acid liquid is employed
in the treatment
and the patient
another way of

Chana-amta.

CICER
soongaricum

The Common Gram

MEDICINE
Chana-khar

ness' (Brigade Surgeon J H Thornton, B A, M B, Monghir) "The

(Native

'The

also in

cholera (Surgeon Major J J L Ratton, Salem) 'It is believed to have alterative properties' (Aligarh)

CHEMISTRY
1095

Chemical Composition—The seeds contain, according to Balfour, moisture 10.80 per cent, fatty matter 4.56 per cent, nitrogenous matter 19.32 per cent, mineral constituent (ash) 3.12 per cent, and starchy matter 62.20 per cent. Dr Warden however, gives the following composition "One hundred parts without husk contain water 11.39 nitrogenous matters 22.7, fat 3.76 starch 63.18, and mineral matter 2.60 (Parkes)" (Conf with Church's *Analysis of Pulse on a previous page*)

Food—Gram forms the chief food for horses. Amongst the poorer classes of natives parched gram (*chabena*) is much eaten. Masson informs us that in the Panjāb it is made into bread, which was a favourite article of food with the Sikh sirdārs. The natives also eat it boiled in the form of

FOOD
1096
Parched
Gram.
1097
Ragout
1098
Young plants
1099
FOODER
1100

ries instead of vinegar

The following account of gram given in the *Treasury of Botany* may be quoted here. In India the seeds form one of the pulses known under the name of 'Gram' and are greatly used as an article of food by the natives being ground into meal, and either eaten in puddings or made into cakes. They are also toasted or parched and in this state are commonly carried for food on long journeys. Rolled in sugar candy, these toasted peas form a rough sort of comfits, and gram flour made up with sesamum oil and sugar candy is an Indian sweetmeat."

Cicer Lens, Willd, see Ervum Lens, Linn

1101

C. soongaricum, Steph, Fl Br Ind, II, 176

Vern.—Tishu, jawāne banyars, sārri, serri, Pā

References—Stewart Pb Pl, 63, Murray, Drugs and Pl Sind 120, Church Food grains of India, p 131

Habitat—Met with in the Western Himālayas, temperate and alpine

FOOD
Seeds
1102Shoots
1103

gram is eaten by the people. The young shoots are prepared as a pickle by the Chinese, and a vinegar is made from the leaves. The latter are often covered by a viscid exudation, with a strong aromatic odour.

C. 1103

The Wild or Indian Endive.

CICHORIUM
Intybus

Aitchison states that in Lahaul shoots are used as a pot-herb, and that the peas are eaten there, as they are, both raw and cooked, in parts of Ladak" (Stewart, Pb Pl, 63, Hinderson, Mission to Yarkand)

CICHORIUM, Linn, Gen Pl, II, 506

Cichorium Endivia, Linn, Fl Br Ind, III, 391, COMPOSITÆ.

THE GARDEN ENDIVE

1104

Pl, 81, DC,
'Ed, Lisboa,
of Botany
a native of
it may, there
is no doubt of its having been used as an esculent food from a very
early period by the Egyptians, through whom the Greeks and Romans
probably became acquainted with it (Treasury of Botany) The Arabs call

1104

Medicine.—"Endive is much valued by the *hakims* as a resolvent and
ious complaints much as taraxa-
he four lesser cold seeds of old
East" (Dymock) The root is
refuge, given in 'munjus,' the
the seed is used in sherbets"

MEDICINE.

Seeds,

1105

Root

1106

Food—"Endive, radishes, and succory are mentioned by Ovid as
forming part of a garden salad, and Pliny states that endive in his time
was eaten both as a salad and pot herb. As such it has been used in

FOOD.

1107

C. Intybus, Linn, Fl Br Ind, III, 391, COMPOSITÆ

THE WILD OR INDIAN ENDIVE, CHICORY, OR SUCCORY.

Vern.—Kasni, HIND, PERS; Hindyba ARAB; Kashni-wira, TAM,
Kasni-wittulu, TEL, Hand gal, suchal, kasni, Pb Kasni, GUJ.

References—Brandis, For Fl, 77, Kurn, For Fl Burm, 77 Stewart,
Pb Pl, 124 Aitchison Pb Pl, 81, DC, Origin of Cult Pl, 56

1108

Habitat.—North-West India, Kumaon, distributed westward to the
Atlantic

§ "In the plains of the Panjáb it is cultivated by natives as a pot-herb
(sôg), and may be an escape, truly wild at 4,000 to 11,000 feet" (Sur-
geon-Major J. E. T. Aitchison, Simla)

C. 1108

CICHORIUM
Intybus

The Wild or Indian Endive.

HISTORY.

History —“The wild perennial chicory, which is cultivated as a salad, as a vegetable, as fodder, and for its roots, which are used to mix with coffee, grows throughout Europe, except in Lapland, in Morocco and Algeria, from Eastern Europe to Afghanistan and Beluchistán, in the Panjab and Kashmir, and from Russia to Lake Baikal in Siberia. The

CULTIVA-
TION
II09

fodder plant is simple enough. The seed is sown broadcast upon land that has been dug or deeply ploughed, from seven to twelve pounds per

rows. When the plants are about five inches in height, carefully hoe them and single out, leaving them about six inches apart, after the usual method in turnip culture,—that is, by boys following the hoers. Some recommend that the seed be sown in a bed, and when the plants are fit for transplanting—which will be when about five inches high—they are to be set out in rows nine inches apart, and at six-inch intervals from plant to plant in the rows. In either case, the land must be kept clean, and well

course of cropping pursued for a few years, and it may then be again sown or planted with chicory.

“In preparing the land for a root crop, deep ploughing is recom-

be carefully dug out and destroyed, when the time for taking up has arrived, because, if allowed to become mixed with the bulk, they will spoil the sample. T is been sown broadcast he crop being easily t nine to twelve inches quan- tity used M ants so as to leave spaces between them in the rows, each about six or eight

The of ing

is adopted’ (*Merton, Cyclop of Agri*, I, 457).

C. II09

Chicory and Coffee

CICHORIUM
Intybus.CULTIVA-
TION.

selling at 2 annas a seer. He mentions specimens of root and of seed as sent to the Lahore Exhibition from nearly every district

Great Britain imports annually close upon 200,000 cwts of the root. It is extensively grown in England, but the best roots are imported from

properties The seeds of the seed is used is bitter and used medicinally in the Panjab. It contains nitrate and sulphate of potash, of chicory

MEDICINE.
IIIO

(Assist in the liver
A strong
ous vomit-
"Much
t Surgeon

Pt.

vegetable"
me of Barbe

FOOD
IIII

roots once constituted half the food of the poorer classes, as they probably do at the present day. "Within the last few years, grocers mixing chicory

Chicory in
Coffee.

chicory by itself. In what the English Grocer requires to do is to sell pure "coffee" when he advertises - may be anything he pleases to ma

ground, Roasted chicory |

C. IIJI

CIMICIFUGA
foetida.

Chicory and Coffee: Black Snake Root.

FOOD.

contains a volatile empyreumatic oil, to which its aroma is due, and a bitter principle. It contains no caffeine. Infused in boiling water it yields a drink allied in flavour and colour to coffee. It is largely used in Bala. — In some parts of Germany the women are said to be regular

Warden, *Prof. of Chemistry*,

The following extract, relating to the fact of the chicory roots being a new source of alcohol, was published in the *Tropical Agriculturist* of 1st December 1882, page 495. also p 57 —

"According to *Erfindungen und Erfahrungen*, the celebrated coffee substitute, chicory, seems likely to become of importance as a source of alcohol. The root contains an average of 24 per cent of substances easily convertible into sugar, and the alcohol obtained by its saccharification, fermentation and distillation, is characterised by a pleasant aromatic taste and great purity" (*Chemist and Druggist*).

ADULTERATIONS.
III2

Adulterations. — "Roasted chicory is extensively adulterated. To colour

dead, dog-dish and baked livers of horses and bullocks (1), are substances which are said to have been used for adulterating chicory. A mixture of roasted pulse (peas usually) and Venetian red has been used under the name of *Hambro' powder* for the same purpose" (*Ure's Dict., Art and Manuf*). A recent examination of certain "coffee mixtures" revealed the fact that roasted cockroaches and iron rust were employed as adulterants. (*See Coffea arabica, para Adulterants*)

CIMICIFUGA, Linn; Gen Pl, I, 9.

III3

Cimicifuga foetida, Linn.; *Fl Br Ind.*, I, 30, RANUNCULACEÆ.

Vern — *Giunti*, PB

References. — *Stewart*, Pb Pl, 2, *Treasury of Botany*, *New Official Guide to the Museum*, 8

Habitat. — Found in the temperate Himálaya, from Bhután to Kashmir, altitude 7,000 to 12,000 feet.

MEDICINE.
Root.
III4

Medicine. — The root is said to be poisonous. In Siberia it is used to drive away bugs and fleas. Under the name of a nearly allied plant (*Actæa spicata*), the writer has already referred to this plant, and chiefly with the view of attracting attention to these useful but apparently neglected plants.

Garrod, in his *Materia Medica*, calls *Cimicifuga racemosa*, Linn., the

made known to Europe in 1690, and was scientifically introduced as a

Black Snake Root: Cinchona Bark.

CINCHONA.

cinal virtues. *C. racemosa* is chiefly prescribed in the form of tincture, and employed in rheu-
and chronic bronchial
been used to reduce
A section of the root

MEDICINE.

shaped sections, with a thick brittle
contains a resinous active principle
Macroton In its action this drug
and colchicum on the other. It is

most useful in acute rheumatism, and a powder of the root is perhaps the
best mode in which to give the drug, in doses of 20 to 30 grains (Royle's
Mat Med. ed by Harley)

Special Opinion — § "A poultice prepared of the fresh leaves is used
here, and said to be very useful in rheumatic affection of joints" (*Surgeon*
C. J. W. Merdous, Burrissal)

CINCHONA, Linn, Gen Pl, II, 32

III5

Cinchona, Linn, RUBIACEÆ.

CINCHONA BARK, PERUVIAN BARK, JESUIT'S BARK, COUNTESS'S
BARK, ECORCE DE QUINQUINA, Fr, CHINARINDE, Germ.

Reference — *U. S. Disp.*, 1825, 1827, 1828, 1829, 1830, 1831, 1832, 1833, 1834, 1835, 1836, 1837, 1838, 1839, 1840, 1841, 1842, 1843, 1844, 1845, 1846, 1847, 1848, 1849, 1850, 1851, 1852, 1853, 1854, 1855, 1856, 1857, 1858, 1859, 1860, 1861, 1862, 1863, 1864, 1865, 1866, 1867, 1868, 1869, 1870, 1871, 1872, 1873, 1874, 1875, 1876, 1877, 1878, 1879, 1880, 1881, 1882, 1883, 1884, 1885, 1886, 1887, 1888, 1889, 1890, 1891, 1892, 1893, 1894, 1895, 1896, 1897, 1898, 1899, 1900, 1901, 1902, 1903, 1904, 1905, 1906, 1907, 1908, 1909, 1910, 1911, 1912, 1913, 1914, 1915, 1916, 1917, 1918, 1919, 1920, 1921, 1922, 1923, 1924, 1925, 1926, 1927, 1928, 1929, 1930, 1931, 1932, 1933, 1934, 1935, 1936, 1937, 1938, 1939, 1940, 1941, 1942, 1943, 1944, 1945, 1946, 1947, 1948, 1949, 1950, 1951, 1952, 1953, 1954, 1955, 1956, 1957, 1958, 1959, 1960, 1961, 1962, 1963, 1964, 1965, 1966, 1967, 1968, 1969, 1970, 1971, 1972, 1973, 1974, 1975, 1976, 1977, 1978, 1979, 1980, 1981, 1982, 1983, 1984, 1985, 1986, 1987, 1988, 1989, 1990, 1991, 1992, 1993, 1994, 1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 2675, 2676, 2677, 2678, 2679, 2680, 2681, 2682, 2683, 2684, 2685, 2686, 2687, 2688, 2689, 2690, 2691, 2692, 2693, 2694, 2695, 2696, 2697, 2698, 2699, 2700, 2701, 2702, 2703, 2704, 2705, 2706, 2707, 2708, 2709, 2710, 2711, 2712, 2713, 2714, 2715, 2716, 2717, 2718, 2719, 2720, 2721, 2722, 2723, 2724, 2725, 2726, 2727, 2728, 2729, 2730, 2731, 2732, 2733, 2734, 2735, 2736, 2737, 2738, 2739, 2740, 2741, 2742, 2743, 2744, 2745, 2746, 2747, 2748, 2749, 2750, 2751, 2752, 2753, 2754, 2755, 2756, 2757, 2758, 2759, 2760, 2761, 2762, 2763, 2764, 2765, 2766, 2767, 2768, 2769, 2770, 2771, 2772, 2773, 2774, 2775, 2776, 2777, 2778, 2779, 2780, 2781, 2782, 2783, 2784, 2785, 2786, 2787, 2788, 2789, 2790, 2791, 2792, 2793, 2794, 2795, 2796, 2797, 2798, 2799, 2800, 2801, 2802, 2803, 2804, 2805, 2806, 2807, 2808, 2809, 2810, 2811, 2812, 2813, 2814, 2815, 2816, 2817, 2818, 2819, 2820, 2821, 2822, 2823, 2824, 2825, 2826, 2827, 2828, 2829, 2830, 2831, 2832, 2833, 2834, 2835, 2836, 2837, 2838, 2839, 2840, 2841, 2842, 2843, 2844, 2845, 2846, 2847, 2848, 2849, 2850, 2851, 2852, 2853, 2854, 2855, 2856, 2857, 2858, 2859, 2860, 2861, 2862, 2863, 2864, 2865, 2866, 2867, 2868, 2869, 2870, 2871, 2872, 2873, 2874, 2875, 2876, 2877, 2878, 2879, 2880, 2881, 2882, 2883, 2884, 2885, 2886, 2887, 2888, 2889, 2890, 2891, 2892, 2893, 2894, 2895, 2896, 2897, 2898, 2899, 2900, 2901, 2902, 2903, 2904, 2905, 2906, 2907, 2908, 2909, 2910, 2911, 2912, 2913, 2914, 2915, 2916, 2917, 2918, 2919, 2920, 2921, 2922, 2923, 2924, 2925, 2926, 2927, 2928, 2929, 2930, 2931, 2932, 2933, 2934, 2935, 2936, 2937, 2938, 2939, 2940, 2941, 2942, 2943, 2944, 2945, 2946, 2947, 2948, 2949, 2950, 2951, 2952, 2953, 2954, 2955, 2956, 2957, 2958, 2959, 2960, 2961, 2962, 2963, 2964, 2965, 2966, 2967, 2968, 2969, 2970, 2971, 2972, 2973, 2974, 2975, 2976, 2977, 2978, 2979, 2980, 2981, 2982, 2983, 2984, 2985, 2986, 2987, 2988, 2989, 2990, 2991, 2992, 2993, 2994, 2995, 2996, 2997, 2998, 2999, 3000, 3001, 3002, 3003, 3004, 3005, 3006, 3007, 3008, 3009, 3010, 3011, 3012, 3013, 3014, 3015, 3016, 3017, 3018, 3019, 3020, 3021, 3022, 3023, 3024, 3025, 3026, 3027, 3028, 3029, 3030, 3031, 3032, 3033, 3034, 3035, 3036, 3037, 3038, 3039, 3040, 3041, 3042, 3043, 3044, 3045, 3046, 3047, 3048, 3049, 3050, 3051, 3052, 3053, 3054, 3055, 3056, 3057, 3058, 3059, 3060, 3061, 3062, 3063, 3064, 3065, 3066, 3067, 3068, 3069, 3070, 3071, 3072, 3073, 3074, 3075, 3076, 3077, 3078, 3079, 3080, 3081, 3082, 3083, 3084, 3085, 3086, 3087, 3088, 3089, 3090, 3091, 3092, 3093, 3094, 3095, 3096, 3097, 3098, 3099, 3100, 3101, 3102, 3103, 3104, 3105, 3106, 3107, 3108, 3109, 3110, 3111, 3112, 3113, 3114, 3115, 3116, 3117, 3118, 3119, 3120, 3121, 3122, 3123, 3124, 3125, 3126, 3127, 3128, 3129, 3130, 3131, 3132, 3133, 3134, 3135, 3136, 3137, 3138, 3139, 3140, 3141, 3142, 3143, 3144, 3145, 3146, 3147, 3148, 3149, 3150, 3151, 3152, 3153, 3154, 3155, 3156, 3157, 3158, 3159, 3160, 3161, 3162, 3163, 3164, 3165, 3166, 3167, 3168, 3169, 3170, 3171, 3172, 3173, 3174, 3175, 3176, 3177, 3178, 3179, 3180, 3181, 3182, 3183, 3184, 3185, 3186, 3187, 3188, 3189, 3190, 3191, 3192, 3193, 3194, 3195, 3196, 3197, 3198, 3199, 3200, 3201, 3202, 3203, 3204, 3205, 3206, 3207, 3208, 3209, 3210, 3211, 3212, 3213, 3214, 3215, 3216, 3217, 3218, 3219, 3220, 3221, 3222, 3223, 3224, 3225, 3226, 3227, 3228, 3229, 3230, 3231, 3232, 3233, 3234, 3235, 3236, 3237, 3238, 3239, 3240, 3241, 3242, 3243, 3244, 3245, 3246, 3247, 3248, 3249, 3250, 3251, 3252, 3253, 3254, 3255, 3256, 3257, 3258, 3259, 3260, 3261, 3262, 3263, 3264, 3265, 3266, 3267, 3268, 3269, 3270, 3271, 3272, 3273, 3274, 3275, 3276, 3277, 3278, 3279, 3280, 3281, 3282, 3283, 3284, 3285, 3286, 3287, 3288, 3289, 3290, 3291, 3292, 3293, 3294, 3295, 3296, 3297, 3298, 3299, 3300, 3301, 3302, 3303, 3304, 3305, 3306, 3307, 3308, 3309, 3310, 3311, 3312, 3313, 3314, 3315, 3316, 3317, 3318, 3319, 3320, 3321, 3322, 3323, 3324, 3325, 3326, 3327, 3328, 3329, 3330, 3331, 3332, 3333, 3334, 3335, 3336, 3337, 3338, 3339, 3340, 3341, 3342, 3343, 3344, 3345, 3346, 3347, 3348, 3349, 3350, 3351, 3352, 3353, 3354, 3355, 3356, 3357, 3358, 3359, 3360, 3361, 3362, 3363, 3364, 3365, 3366, 3367, 3368, 3369, 3370, 3371, 3372, 3373, 3374, 3375, 3376, 3377, 3378, 3379, 3380, 3381, 3382, 3383, 3384, 3385, 3386, 3387, 3388, 3389, 3390, 3391, 3392, 3393, 3394, 3395, 3396, 3397, 3398, 3399, 3400, 3401, 3402, 3403, 3404, 3405, 3406, 3407, 3408, 3409, 3410, 3411, 3412, 3413, 3414, 3415, 3416, 3417, 3418, 3419, 3420, 3421, 3422, 3423, 3424, 3425, 3426, 3427, 3428, 3429, 3430, 3431, 3432, 3433, 3434, 3435, 3436, 3437, 3438, 3439, 3440, 3441, 3442, 3443, 3444, 3445, 3446, 3447, 3448, 3449, 3450, 3451, 3452, 3453, 3454, 3455, 3456, 3457, 3458, 3459, 3460, 3461, 3462, 3463, 3464, 3465, 3466, 3467, 3468, 3469, 3470, 3471, 3472, 3473, 3474, 3475, 3476, 3477, 3478, 3479, 3480, 3481, 3482, 3483, 3484, 3485, 3486, 3487, 3488, 3489, 3490, 3491, 3492, 3493, 3494, 3495, 3496, 3497, 3498, 3499, 3500, 3501, 3502, 3503, 3504, 3505, 3506, 3507, 3508, 3509, 3510, 3511, 3512, 3513, 3514, 3515, 3516, 3517, 3518, 3519, 3520, 3521, 3522, 3523, 3524, 3525, 3526, 3527, 3528, 3529, 3530, 3531, 3532, 3533, 3534, 3535, 3536, 3537, 3538, 3539, 3540, 3541, 3542, 3543, 3544, 3545, 3546, 3547, 3548, 3549, 3550, 3551, 3552, 3553, 3554, 3555, 3556, 3557, 3558, 3559, 3560, 3561, 3562, 3563, 3564, 3565, 3566, 3567, 3568, 3569, 3570, 3571, 3572, 3573, 3574, 3575, 3576, 3577, 3578, 3579, 3580, 3581, 3582, 3583, 3584, 3585, 3586, 3587, 3588, 3589, 3590, 3591, 3592, 3593, 3594, 3595, 3596, 3597, 3598, 3599, 3600, 3601, 3602, 3603, 3604, 3605, 3606, 3607, 3608, 3609, 3610, 3611, 3612, 3613, 3614, 3615, 3616, 3617, 3618, 3619, 3620, 3621, 3622, 3623, 3624, 3625, 3626, 3627, 3628, 3629, 3630, 3631, 3632, 3633, 3634, 3635, 3636, 3637, 3638, 3639, 3640, 3641, 3642, 3643, 3644, 3645, 3646, 3647, 3648, 3649, 3650, 3651, 3652, 3653, 3654, 3655, 3656, 3657, 3658, 3659, 3660, 3661, 3662, 3663, 3664, 3665, 3666, 3667, 3668, 3669, 3670, 3671, 3672, 3673, 3674, 3675, 3676, 3677, 3678, 3679, 3680, 3681, 3682, 3683, 3684, 3685, 3686, 3687, 3688, 3689, 3690, 3691, 3692, 3693, 3694, 3695, 3696, 3697, 3698, 3699, 3700, 3701, 3702, 3703, 3704, 3705, 3706, 3707, 3708, 3709, 3710, 3711, 3712, 3713, 3714, 3715, 3716, 3717, 3718, 3719, 3720, 3721, 3722, 3723, 3724, 3725, 3726, 3727, 3728, 3729, 3730, 3731, 3732, 3733, 3734, 3735, 3736, 3737, 3738, 3739, 3740, 3741, 3742, 3743, 3744, 3745, 3746, 3747, 3748, 3749, 3750, 3751, 3752, 3753, 3754, 3755, 3756, 3757, 3758, 3759, 3760, 3761,

CINCHONA.

Cinchona Bark.

compiling the present article, but has at the same time verified the historic and other facts by consulting the works enumerated above

Habitat.—Dr. King says: "The trees producing the medicinal barks are all natives of tropical South America, where they are found in the dense forests of the mountainous regions of the western parts of that continent at a height of from 2,500 to 9,000 feet above the level of the sea, and in an equable but comparatively cool climate. The Cinchona-producing region forms a crescentic zone which follows the contour of the coast line, but nowhere actually touches the coast, extending to 20° S latitude. a hundred miles in width, but than two thousand. During its of Venezuela, New Granada,"

the zone in 20° S were described by M. Weddell in his splendid monograph published at Paris in 1849"

HISTORY.

HISTORY OF THE INTRODUCTION OF THE DRUG INTO EUROPE.

"The introduction of the medicinal Cinchona bark to Europe was the result of a Spanish Viceroy of Peru, who, after recovering from an attack of fever contracted in the mountains, brought the bark to Europe on his return in the year 1639. Jesuit missionaries were also taken as active part in its introduction. Hence the name *jesuit's bark*, and the name of the tree *Quina*, from Jussieu, member of the Académie des Plantes at Paris, who discovered it near the mouth of the River Amazon. The first living Cinchonas ever

introduced this invaluable remedy to Europe (King).

HISTORY OF THE ALKALOIDS.—"The most important and at the same time peculiar constituents of Cinchona barks are the alkaloids

C. 1116

History of the Alkaloids.

CINCHONA.

enumerated in the following table:—

Alkaloid	Chemical composition	HISTORY OF THE ALKALOIDS.
Cinchonine	$C_{20}H_{21}N_3O$	
Cinchonidine (quinidine of many writers)	Same formula.	
Quinine	$C_{20}H_{21}N_3O_2$	
Quinidine (conquinine of Hesse)	Same formula	
Quinamine	$C_{20}H_{25}N_3O_4$	

so fe
febr
cert.
alka

the outward appearance of these being alike. With the separation of the new alkaloids, chemical tests for their recognition began to be inserted in

bark still continues to be rated by the European quinine-makers in proportion to the percentage of quinine it contains, the other alkaloids being

devoid of quinine, while those of the same species from a neighbouring

down to less than 1 per cent.

"Among"
are a great n
principles, of
altogether w

CINCHONA.

History of its Introduction into India.

HISTORY
OF THE
ALKALOIDS.

observed, was obtained by Broughton from a bark grown at Ootacamund. This bark afforded not less than 13½ per cent. of alkaloids, among which quinine was predominant.

"The few facts just mentioned show that it is impossible to state even

quinine

"As to *Crown* or *Loxa* bark, the *Cortex Cinchonæ palidæ* of pharmacy, its merits are, to say the least, very uncertain. On its first introduction in the seventeenth century, when it was taken from the trunks and large branches of full-grown trees, it was doubtless an excellent medicinal bark; but the same cannot be said of much of that now found in commerce, which is to a large extent collected from very young wood. Some of the *Crown* Bark produced in India is, however, of extraordinary excellency, as shown by the recent experiments of DeVry.

"As to red bark, the thick flat sort contains only three to four per cent. of alkaloids, but a large amount of colouring matter. The quill Red Bark of the Indian plantations is a much better drug, some of it yielding 5 to 10 per cent. of alkaloids, less than a third of which is quinine and a fourth cinchonidine, the remainder being cinchonine and sometimes also traces of quimidine (conquinine).

"The variation in the amount of alkaloids relates not merely to their total percentage, but also to the proportion which one bears to another. Quinine and cinchonine are of the most frequent occurrence; cinchonidine is less usual, while quimidine is still less frequently met with, and never in large amount. The experiments performed in India have

HISTORY OF THE INTRODUCTION OF CINCHONA INTO INDIA.

Dr. King writes "The practice of the bark collectors in the wild regions in which Cinchonas naturally grow involved the destruction of each tree felled for its bark, yet no measures were ever taken by the owners of either public or private forests to secure supplies for the future by conservancy or re-planting. Meanwhile the consumption of bark in Europe steadily increased, and, as a natural result, prices rose, and fears began to be entertained that the supply would ultimately fail. The British and Dutch Governments being, by reason of their tropical possessions, the

INTRODUC-
TION INTO
INDIA.

History of its Introduction into India.

CINCHONA.

the preservation of the natural forests, that great fears have been entertained that the supply might altogether cease, or be obtainable only at a price which would place it beyond the reach of the mass of the community."

HISTORY
OF THE
INTRODUC-
TION INTO
INDIA.

"Dr. Royle's recommendations, although approved of, were not at the time acted upon, but were allowed to remain in abeyance until 1859, when the increasing and constantly increasing destruction of Government forests seemed almost certain to result in the complete destruction of the Cinchona forests of India."

Gardner, recommended that an intelligent and qualified gardening collector should be deputed for a couple of years to the mountains of South America for the purpose of exploring the Cinchona forests, and of procuring seeds and barks."

the matter, as also did the late Dr. J. Anderson. The Medical Board of India, in 1859, recommended that an intelligent and qualified gardening collector should be deputed for a couple of years to the mountains of South America for the purpose of exploring the Cinchona forests, and of procuring seeds and barks."

to be found
bark forests.
Spruce and
the eastern
Markham
which he had

the inhabitants and flora of regions he traversed. Landing at Islay in March 1860, Mr. Markham, accompanied by Mr. Weir (a practical

CINCHONA.

History of its Introduction into India

HISTORY
OF THE
INTRODUC-
TION INTO
INDIA

great plain of western Brazil. Mr. Markham penetrated this valley

micrantha.

"Instead of sending these plants direct to India, Mr. Markham was compelled by his orders to take them to India *via* Panama, England, the Mediterranean and the Red Sea, and thus to expose them to transshipments and alterations of temperature which ultimately killed them all

"About the time Mr. Markham was exploring the yellow bark forests of Southern Peru, Mr. Pritchett was collecting seeds and plants of the

Andes, and he was thus enabled very speedily to form at Limon a nursery of young plants of *Cinchona succirubra*, which were ultimately conveyed safely to India by Mr R Cross. A quantity of seeds of this species was also collected and sent to India by post. Mr. Cross was subsequently commissioned to procure seeds of the pale barks in the forests near Loxa, and this commission he executed with great success. A third expedition to New Granada was made by the same collector with the object of securing seeds of the Cartagena bark, *Cinchona lancifolia* and *pita-yensis*. The seeds obtained by Mr. Cross were sent to Kew, where they

Kew, where some were retained and sown. A few of the plants brought from South America were also retained at Kew, so that a sort of reserve depôt was formed there in case of failure in India. For the successful introduction of *Cinchona* into India and other British possessions, Government are largely indebted for advice, as well as for more active

History of its Introduction into India.

CINCHONA.

two months later. In the month of December 1861, Dr Anderson delivered over to Mr. McIver at Ootacamund the plants he had brought from the Cinchona plantation which the Dutch had just succeeded in establishing in Java. Dr. Anderson had been sent by the Government of India to the Netherlands to obtain the plants, and by the courtesy of the Dutch authorities he secured 50 plants of Cinchona Calisaya, four of which were of the variety of Pahudiana. On the 4th March 1862, Dr. Anderson sent Mr. McIver 50 crown bark seeds from Loxa archona to India became thus an accom-

HISTORY
OF THE
INTRODUC-
TION INTO
INDIA

plished fact" (King).

Introduction into South India.—"The success of Cinchona succirubra and officinalis on the Nilgiris has been remarkable. Not only do the trees grow luxuriantly, but their bark is richer in alkaloids than much of the Cinchona bark imported from South America. The Government plantations there, according to the returns for 1884-85, contain 1,618,744 trees of sorts. The Nilgiri plantations were under the superintendence of Mr. McIver until his death, since which they have been under Mr. M. A. Lawson.

South India.

"Encouraged by its success on the Nilgiris, Cinchona cultivation was warmly taken up by European residents in the other high lands and hill ranges of the Madras Presidency. The coffee planters of Wynad put out a good many red bark trees on their estates, and these are found to grow well. In South Canara a small plantation was formed in 1869, at a place called Nagooli, above the Kooloor Ghât, and at an elevation of 2,500 feet above the sea, but the experiment there was pronounced by the Madras Government as unlikely to be productive of useful results, and was abandoned. On the Mahendra Mountain, in the Ganjam district, the Madras Government (the Forest Department, the Nulla Mully barks), and, as the experiment was un-
nivally, and She
Cinchona was taken up to a greater or less extent, both by private planters and the Government" (King).

most probably thrive best. For the hardier kinds Mr. Markham

CINCHONA

History of its Introduction into India

HISTORY
OF THE
INTRODUC-
TION INTO
INDIA

this ease is the result of the patience and intelligence which Mr McIver

tions Of these the following are the more important:—

- | | |
|-----------------------------------|---------------------------------------|
| (1) <i>C. officinalis.</i> | (8) <i>C. verde</i> (com form). |
| (2) <i>C. succirubra.</i> | (9) <i>C. zamba morada</i> (com form) |
| (3) <i>C. Calisaya.</i> | (10) <i>C. carthagena</i> (com form) |
| (4) <i>C. Ledgeriana.</i> | (11) <i>C. Pahudiana</i> |
| (5) <i>C. javanica.</i> | (12) <i>C. Humboldtiana.</i> |
| (6) <i>C. Santa Fe</i> (com form) | (13) <i>C. Pitayensis.</i> |
| (7) <i>C. morada</i> (com form) | (14) <i>C. micrantha.</i> |

He adds "Of these, the only kinds which are largely grown in the Govern-

Bengal

Sikkim plantation has been under the charge of Dr Anderson's successors, viz, Mr O B Clarke, during 1870 and 1871, and Dr George King, since the latter date Since 1866, the Sikkim plantations have

Calcutta from Ootacamund 193 plants of *succirubra* and of the species yielding grey bark Some of the Java plants died in Calcutta, and on the 19th January 1862 the total stock in the Botanical Gardens there from

him,
the
have

History of its Introduction into India

CINCHONA.

been largely increased, and at 31st March 1885 their contents were as follows —

	Red (Cinchona succubra)	Yellow (Cinchona Calisaya and Ledgeriana)	Yellow (Cinchona Calisaya verde and morada)	Hybrid (unnamed variety)	Other kinds	Total of all sorts
Mungpoo Division	2,132,000	801,118	134,300	345,100	25,593	3,438,111
Sitong „	1,100,000	70,000	15,000	40,000	...	1,225,000
Rungjung „		2,15,000	34,000			249,000
GRAND TOTAL OF ALL KINDS	3,232,000	1,076,118	183,300	385,100	25,591	4,912,111

HISTORY
OF THE
INTRODUC-
TION INTO
INDIA.

“A Cinchona plantation was begun by a private company in Sikkim almost simultaneously with that belonging to Government, and more recently a second such plantation has been opened out in Bhotan. Patches of Cinchona were also planted in several tea gardens in the

Khasia hills.

“*Into North-Western Provinces* — The cultivation also received a very patient trial for several years in the North-Western Provinces of India, and plantations were begun at various altitudes from 2,000 to 6,500 feet above the sea, but the plants all ultimately perished from frost. A similar result followed the spirited attempt of Colonel Nassau Lees to grow

N.-W.
Provinces.

Bombay.

Burma.

north of Toungoo, and about 54,000 plants are now alive. But the plantation does not thrive so well as could be wished, and it is desirable that the advice of an expert should be obtained as to the best course to be taken. It was hoped that Dr King would have visited Burma, but as yet he has been unable to do so. If the Government of Bengal can spare him, perhaps he will be able to come in May 1883. At Poonchoung the cultivation of Cinchona has done so poorly that orders have been given to abandon further outlay on the experiment there. About 300 lb of Cinchona bark were recently received from Thandoung, and

ultivation of Cinchona
otanist, Dr Thwaites,
It was subsequently

Ceylon.

CINCHONA
Calisaya.

The Yellow Bark of Commerce.

HISTORY
OF THE
INTRODUC-
TION INTO
INDIA.

taken up with great vigour by the very spirited planting community of that then most flourishing colony, and to such an extent was the cultivation carried, that in the year 1881 no less than three millions of pounds of dry Cinchona bark were exported from that island to England, and in subsequent years the exports have materially increased" (King) During the years 1885-86-87, Dr King informs the writer the annual exports from Ceylon touched 15 million pounds

THE SPECIES OF CINCHONA.

There are bet

will be necessary only to allude to the better known species and varieties which are cultivated in India

III7

Cinchona Calisaya, Weddell, RUBIACEÆ

THE CALISAYA BARK OF YELLOW BARK OF COMMERCE, a term also applied to the bark of C LEDGERIANA

Vern — Bārak, DEC, Shurappattai, TAM, Yradap-patta TEL

References — Kew Reports, 1877, pp 14, 28 1879, pp 12, 13 1880, pp 11, 25, 32, 1881, 25, 1882, pp 18, 19, 38; Trop Agriculturist, 1883, 706

most only second to C succirubra in point of importance in the Sikkim plantations. In a Resolution of the Bengal Government dated March 1883, it is stated that Mr Wood was of opinion that good quinine barks

tion was not acted upon for some time. Full effect has, however, been given to it of recent years, and succirubra has been supplanted by Calisaya to the extent of about a million trees. On the other hand, the attempt to cultivate this species in the Nilgiri hills has been practically abandoned. Calisaya was discovered by M Weddell in 1847, it is a native of Bolivia and South Peru. The supply of bark from natural

MEDICINE

Bark

III18

Powder

III19

Leaves

II20

C. II20

The Ledgeriana Bark of Commerce

CINCHONA
Ledgeriana

uncoated, consisting almost entirely of liber, is $\frac{1}{3}$ to $\frac{1}{2}$ inch thick Its MEDICINE.

Flax of the Pharmacopæia

Structure of the Wood — Reddish-grey, moderately hard, even-grained
Pores small, in short radial lines Medullary rays fine, closely packed

TIMBER.
II21

VARIETIES OF C CALISAYA

Numerous varieties and hybrids have been distinguished of this species, especially by Weddell. The best known are *var Josephiana* (named after Ledgeriana, but C zamba, are being experimentally

Josephiana.
II22

Zamba
II23

Morada
II24

Verde.
II25

Blanca.
II26

in 1873. "Our plantation " "The outward of know manner of harvesting, drying, and packing, but certain it is that their treatment is highly spoken of " "There are numerous varieties of C. Calisaya, but we possess one with which we have become acquainted,

Cinchona Ledgeriana (a cultivated form)

II27

Cinchonas, and consequently the amount of bark harvested in a given number of years is much smaller than that taken from other kinds. The bark also, when it is renewed, is less rich in quinine than the natural bark, so that the trees, instead of having their bark improved by the process of

C. II27

CINCHONA
officialis.

LOXA or Crown Bark of Commerce.

stripping, as is the case in the other kinds of Cinchona, decrease in value. These two circumstances make it doubtful if plantations of *C. Ledgeriana* will, in the long run, be much more profitable to the planter than those formed of the more hardy and reliable *C. Calisaya*.

was certain to prove more remunerative than that of any other species. It could be propagated at lower altitudes than the others (scarcely growing above 4,000 feet), and was, from this point alone, a more economical plant. In *Silva de C. Ledgeriana* Mr. Hooper remarks:—

"To-
veller,
near
= and

proved by far the most productive in quinine of all Cinchona barks. The tree is a mere form of *C. Calisaya*. Mr. Hooper, Quinologist to the Madras Government, in a recent report, remarks:—"In the Ledger bark it will be noticed that there is a steady rise of quinine up to the age of between five and six years, after which there is no apparent increase."

1128

Cinchona carthagenae (Commercial name)

This has been successfully introduced into the Nilgiri hills within the past few years, and Mr. Lawson alludes to it in his reports. In 1881-82 he says that up to date "the propagation of this valuable Cinchona was carried on with most satisfactory results." Again, in 1882-83, the plants "continue to make a very satisfactory growth."

1129

C. officialis, Hook.

LOXA or CROWN BARK; the Pale Bark of Commerce

Syn.—*C. CONDA-MINEA*, Humb.

References.—*Year Book of Pharm.*, 1873, 447, 1875, 161, 1878, 444

"The bark of *C. officialis* is obtained at high altitudes.

MEDICINE.
Loxa Bark.
1130
TIMBER.
1131

of *C. Calisaya*.

At the present day it is scarcely possible to obtain Loxa or Crown bark from South America; India, Ceylon, and Jamaica being the chief sources of the bark in commerce.

C. 1131

Red Bark of Commerce

CINCHONA
succirubra.

to 1 per cent.
5 per cent.
honidine and

CINCHONINE

Cinchona succirubra, *Pavon*

RED BARK

References — *Year Book of Pharm.*, 1873, 70—73, 447, 1874, 19—20, 150—154, 1875, 12, 159, *Ann. Report*, 1877, 28

Habitat — Cultivated on the Nilgiris and other hills of South India, at the plantations of Rangbi and Poomong in Sikkim, on the hills east of Toungoo, in Burma, and in parts of the Satpura Range in Central India

Mr. Lawson writes of South India, while speaking of *C. officinalis*: "The *C. succirubra*, on the other hand, has a bold sturdy stem, which in rich soil and sheltered situations, grows to the height of 50 feet or more. The leaves are a bright apple-green in colour, and a plantation made up of this species looks as light and bright, as that of the *C. officinalis* looks dark and gloomy."

The red *C. succirubra* and the *C. officinalis* have been compared

Medicine — This species thrives at a lower elevation than the others, but is comparatively poor in quinine, though rich in cinchonine and cinchonidine. It yields its best bark when eight years old. From it is chiefly derived the "Cinchona Febrifuge," which is now largely manufactured at the Government Plantation of Rangbi. Mr. W. Elborne remarks (*Pharm. Soc. Jour.*) "The experiments of Mr. J. E. Howard and others have proved that the bark of the root contains a larger proportion of alkaloids than that of the stem, and that the proportion of alkaloid diminishes upwards to the branches." Mr. David Howard has also shewn that the nature of the alkaloid varies according to the part of the tree from which the bark has been taken.

In the opinion of pharmacists the bark most suitable for medicinal use is the *Cinchona succirubra*. The cause of this preference, as pointed out by Mr. Holmes, are the following: (1) the red bark supply will probably be always equal to the demand on account of its growing on a much

MEDICINE.
Red Bark.
II33

CINCHONA
succirubra.**Hybrids of Cinchona****MEDICINE.**

ing matter. The brick-red colouring matter is not found in the growing plant but in the dried bark, and Mr. J. E. Howard considers that it is

TIMBER.**II34****HYBRIDS.****II35**

ations. They are now implicated with resin which appears to have also become oxidised so as to act the part of an acid, and is with difficulty separated. But the most remarkable feature is the altered condition of the alkaloids themselves. Quinine, which formed a considerable portion of the whole, is now diminished, while cinchonine and cinchonidine remain much the same. The quill red bark of Indian plantations is a much better drug, some of it yielding 5 to 10 per cent. of alkaloids, less than a third of which is quinine and a fourth cinchonidine, the remainder being cinchonine and sometimes traces of quinidine (*Elborne*)

Structure of the Wood—Yellow, moderately hard. Pores small in radial lines; medullary rays, closely packed, fine and very fine.

HYBRIDS OF CINCHONA.

Kuntze, after examining the living *Cinchonas* in the Indian planta-

manifest a greater tendency to variation and hybridization than do the plants referred to the genus *Cinchona*. Mr. J. Broughton, in a report

that this ready hybridism between the species of *Cinchona* affords an explanation of the occurrence

Angustifolia.**II36****Bonplandiana****II37****C. II37**

Chemical peculiarities of the Cinchonas

CINCHONA

guish it from the numerous self sown hybrids that are constantly appearing in the plantations. Of this form Mr C B Clarke wrote in 1871, that the gardener took it for *C pitayensis*. Mr McIver thought it was *C unita*.

variety
bark to

J L C N S

by hybridization or otherwise so as to produce a plant that will give the maximum of quinine or other alkaloid desired to be obtained.

CHEMICAL PECULIARITIES OF THE CINCHONA PLANTS

We may conclude this account of the forms of Cinchona grown in India by displaying their chemical peculiarities in the following table of comparative analysis taken from Mr Lawson's report —

CHEMICAL
PECULIARITIES
II38

The Analyses of the different kinds of barks grown on the Government estates given below have been made during the past year by Mr Hooper, the Government Quinologist

		Quinine	Cinchonine	Quinine	Cinchonine	Amorphous alkaloids	Total	Sulphur quinine
1	<i>C officinalis</i> natural	2.77	1.57	16	39	50	5.39	3.72
2	, mossed	3.40	1.50	20	45	62	6.17	4.57
3	, renewed	4.21	.85	22	65	70	6.63	5.66
4	<i>C. angustifolia</i> , natural	3.97	1.3	12	12	87	6.40	5.34
5	, mossed	5.60	1.41	33	04	97	8.35	7.53
6	, renewed	4.91	.89	38	19	114	7.51	6.60
7	<i>C. succirubra</i> natural	1.91	2.11		1.14	88	6.04	2.57
8	, mossed	1.69	2.03		1.68	98	6.38	2.27
9	, renewed	1.84	1.48		1.25	71	5.28	2.47
10	, branch	1.38	2.28		1.59	1.16	6.41	1.85
11	, root	1.24	.77	41	1.43	1.27	5.12	1.66
12	, renewed shavings	2.30	1.16		2.06	1.45	6.97	3.09
13	<i>C. robusta</i> natural	1.43	2.08		1.58	31	5.40	1.92
14	, mossed	1.92	3.16		.77	35	6.20	2.58
15	, renewed	4.40	2.54		51	1.05	9.10	5.92
16	, branch	1.64	2.71		1.17	50	6.02	2.20
17	<i>C. micrantha</i> natural				1.92	40	2.3	
18	, renewed	tr	2.45		1.12	1.0	4.59	
19	, branch				1.60	45	2.05	
20	<i>C. Calisaya</i> natural	1.71	2.32		2.13	29	5.95	1.62
21	, branch	.59	.73		1.93	48	3.73	.79

**CINCHONA
succirubra.****Hybrids of Cinchona.****MEDICINE.**

bark acquires its colour, the cinchotannic acid in which it abounds having become oxidised and changed into cinchona red, and under these conditions the alkaloids also appear to undergo some corresponding alterations. They are now implicated with resin which appears to have also become oxidised so as to act the part of an acid, and is with difficulty separated. But the most remarkable fact is the altered condition of the alkaloids. A portion of the whole remains much the same, but much better than a third of being cinchonin.

TIMBER.**II34****HYBRIDS.****II35**

Structure of the Wood—Yellow, moderately hard. Pores small in radial lines, medullary rays, closely packed, fine and very fine.

HYBRIDS OF CINCHONA.*argentea* Colla.**II36***Bonplandiana***II37**

allied to the form *Bonplandiana*. From the fact that it is reproduced by

Chemical peculiarities of the Cinchonas

CINCHONA

sively cultivated Dr King, in his report for 1874, says 'The analysis of the bark' of this hybrid or species 'shows it to contain much quinine. Since the discovery of this fact, every effort has been made to propagate this

by hybridization or otherwise, so as to produce a plant that will give the maximum of quinine or other alkaloid desired to be obtained

CHEMICAL PECULIARITIES OF THE CINCHONA PLANTS

We may conclude this account of the forms of Cinchona grown in India by displaying the chemical peculiarities in the following table of comparative analysis taken from Mr Lawson's report —

CHEMICAL
PECULIARITIES
II38

The Analyses of the different kinds of barks grown on the Government estates given below have been made during the past year by Mr Hooper, the Government Quinologist

		Quinine	Cinchonine	Quinine	Cinchonine	Amorphous alkaloids	Total	Sulph. quinine
1	C. officinalis natural	2.77	1.57	16	39	50	5.39	3.72
2	" mossed	3.40	1.50	20	45	62	6.17	4.57
3	" renewed	4.21	.85	22	65	70	6.63	5.66
4	C. angustifolia natural	3.97	1.32	12	12	87	6.40	5.34
5	" mossed	5.60	1.41	33	04	97	8.35	7.53
6	" renewed	4.91	.89	38	19	114	7.51	6.60
7	C. succirubra natural	1.91	2.11		114	83	6.04	2.57
8	" mossed	1.69	2.03		1.68	98	6.38	2.27
9	" renewed	1.84	1.48		1.25	71	5.28	2.47
10	" branch	1.38	2.8		1.59	116	6.41	1.85
11	" root	1.24	.77	41	1.43	1.27	5.12	1.66
12	" renewed shaved	2.30	1.16		2.06	1.43	6.97	3.09
13	C. robusta natural	1.43	2.08		1.58	31	5.40	1.92
14	" mossed	1.93	3.16		.77	35	6.20	2.58
15	" renewed	4.40	2.54		51	1.65	9.10	5.92
16	" branch	1.64	2.71		1.17	50	6.03	2.20
17	C. micrantha, natural				1.92	40	2.32	
18	" renewed	tr	2.45		1.12	1.07	4.59	
19	" branch				1.60	45	2.05	
20	C. Calisaya natural	1.21	2.32		2.13	29	5.95	1.67
21	" branch	.59	.73		1.93	48	3.73	.79

Climate, &c, suitable for Cinchona Cultivation

CINCHONA.

CULTIVATION.

the temperature, which shows a minimum temperature of 40° and Fahrenheit, the mean minima for mean maxima 71·7° and 72·28° 65·6° and 64·89°, respectively fairly suitable for succubra, but rather cold for Cansaya. A more congenial climate for both species is indicated by the figures obtained at a lower station (elevation above the sea 2,556 feet) which, for the years 1866 and 1867, are as follow —

Minimum temperature	40° and	41° Fahr.
Maximum	92 3°	94°
Mean minimum temperature	59 3°	60 94°
" maximum	80 6°	81 59°
" temperature	70 1°	71 26°

"In various parts of Ceylon a favourable climate for Cinchona is obtained, as will be seen from the following extract from a most reliable local publication —

and Cinchona without being injurious to human health. Dismissing the weather, in the shade of 73·2° Fahrenheit, resulting, as we 65·8° Fahrenheit Cinchonas were at first rather misunderstood, their preference for incessant rain and mist

weather affects the plants in flag. In Sikkim, succubra makes its most vigorous growth during the latter half of the rains, but both on the Nilgiris and Himálayas the trees continue to grow for two months after the rains cease.

"Observations which have been made show that (calculated on the returns of five years) there are at Ootacamund no fewer than 218 dry days in the year and at Neddiwattum about 240 dry days. The rainfall of the former locality (on an average of three years) is about 44 inches per

"As regards elevation above the sea, it is found that in the Nilgiris succubra succeeds best at altitudes of from 4,500 to 6,000 feet. An elevation of 7,000 feet is found to be too high, the growth being too slow to be profitable. Pale or crown barks thrive in a zone above this, and seem

CINCHONA.

Treatment of the Removed Bark.

COLLECTION.

stems that had been operated upon with a coating of moss or straw in process were very satisfactory so discovered that, provided to coat the partially decorticated stems, Director of the Dutch plantations in Java, suggested a modification of this process which consists in shaving off the superficial layers of bark from the whole surface of the stem, care being taken that at no point shall the young wood be laid bare. Mr. Moens found that the bark of trees thus treated gradually acquires its former thickness, and that the renewed bark is richer in alkaloids than the original bark. This process has been successful in

(King).
not resorted to
the bark under
if ants" (Roso-

In
Madras.
II43

detail (than in

the Govern-
ment plantations is that known under the name of stripping. The barker, with the sharpened point of an ordinary pruning knife, makes several cuts running down the stem parallel to each other, about an inch apart, and then with the blunt back of his knife, he raises every alternate narrow

away. If, on the other hand, the layer of cambium cells is crushed or scratched off by clumsy workmanship, no new bark will be formed. In order to facilitate this new formation of bark the stem is covered with moss or other material.

the latter case it is either up-rooted and a young plant put in its place, or it is cut down and one or more shoots are allowed to spring up from its stool.

TREATMENT

TREATMENT OF THE REMOVED BARK.

Bengal.
II44

In Bengal—"After removal from the trees, Cinchona bark has to be carefully dried, and on the best modes of doing this careful experiments have been made. From these it has been found that exposure to a high

Diseases of Cinchona Trees.

CINCHONA.

temperature, especially in a moist atmosphere, causes bark to become	TREATMENT OF BARK.
--	-----------------------

mouldy and to ferment, as is apt to happen if it be taken off during wet weather, deterioration more or less serious surely occurs. Dry bark, on the other hand, will keep unchanged for many months. Mr Broughton calculates that trunk bark loses from 70 to 74.8 per cent of weight in 76 per cent. The Sikkim experience sees 73 per cent, and twig bark 75

In Madras — After the bark is removed from the trees it is dried by the sun or by artificial heat. It is then packed in gunny bags, forming for sale, and Mr. Broughton exposing the idence of this, of the fact, so ears, however,

In
Madras.
II45

DISEASES OF THE CINCHONA TREES

"Cinchona trees are liable to a kind of canker, which often destroys the terminal and lateral branches, and not unfrequently kills the plants outright. This canker is most abundant in situations where the subsoil is

DISEASES.
II46

CINCHONA.

Diseases of Cinchona Trees

DISEASES

ly fatal, the other local and by no means fatal. The former disease is confined entirely to trees which have been originally planted in damp situations or in situations which have become damp subsequently by the oozing of drainage water in the way already explained. Disease first attacks the roots of such trees. Its existence becomes apparent by the discolorization of their leaves, which ultimately fall off. Gradual shrivelling of the

occasionally these appearances extend to the wood, but as a rule they do not. In size the patches vary, many are about the size of a shilling others are much larger. They are not numerous on one tree and are often confined to a single branch. When small no apparent affection of the general

prepared with a theory as to its cause. This disease is not confined like the last to certain spots, but is found on plants in all parts of the plantation.

A careful examination of all that has been written and of the evidence recorded before the Commission led to the conclusion that the diseases were by the Mr McIver professional cause the damp soil to the late Mr Scott in his a probable cause of the the atmosphere checking

Calicut that of the

It may be concluded that, with care in the selection of sites and the more perfect system of cultivation now pursued all danger from disease has been practically removed

Government Cinchona Febrifuge and Quinine

CINCHONA.

ANNUAL YIELD OF BARK.

In Bengal—The outturn of bark from the Government plantation was, in 1885-86, 339,201 lb, bringing the total yield of bark up to 3,256,927 lb. Almost the whole of the large amount has been used in the manufacture of quinine, which, during the year, has been (for the effect of the effect) paragraphs of

YIELD.

Bengal.
1147

Madras
1148

RESPECTIVE VALUE OF THE ALKALOIDS

"As has been already explained, the medicinal crystallizable alkaloids contained in the bark are quinine, cinchonidine, quinidine, and cinchonine together with an amorphous alkaloid. A fifth called aricine is occasionally found, but has never been used in medicine. M. Hesse has also recently announced the existence of another alkaloid occurring only in the succubra bark grown in Sikkim. This base has received the name of

VALUE OF ALKALOIDS

always been much esteemed and of late years (since it began to get scarce) has brought a price as high or even higher than that got for the barks richer in quinine" (*King*)

GOVERNMENT CINCHONA FEBRIFUGE AND QUININE

FEBRIFUGE
1149

"It had for many years been suspected that the other alkaloids in which red bark is so rich are nearly, if not quite, as efficacious febrifuges as quinine. The settlement of this point naturally demanded attention at an early stage of the Cinchona experiment. In order to settle it by actual trial, Commissions of medical officers of Government were appointed, and the result of an extended series of trials instituted by them may be given in the following extracts from their reports—

"In regard to the relative effects of the three new alkaloids, and with them chemically pure sulphate of quinine, the evidence derived from their use shows that with the exception of sulphate of cinchonine, as

CINCHONA.

Government Cinchona Febrifuge and Quinine.

FEBRIFUGE.

power, and in equal circumstances their use produced almost the same physiological results.

"The results confirm the report on the subject of the Commission, and doubt, that of quinine, and at sulphate of cinchonine, though considerably inferior to the other alkaloids, is, notwithstanding, a valuable remedial agent in fever.

"There is no longer room to doubt that the alkaloids are capable of being generally used with the best effects in India. They have been compared with quinine, a drug which possesses, more than any other that can be named, the confidence of medical practitioners here; and have been

others, and no more than "half" of the ... any

; and
other
d by

the quinine-maker as good American yellow. The establishment of the therapeutic excellence of these alkaloids largely increased the value of the red bark plantations in India, and made much easier of solution the problem of supplying its fever-stricken population with a cheap and effectual febrifuge. And for the solution of this problem the Government very speedily took active steps, by appointing Mr. J. Broughton, a skilled chemist educated in England, as Quinologist to the Nilgiri plantations. Mr. Broughton, after making some valuable observations on the chemistry of living Cinchonas and initiating a process for extracting the whole of the alkaloids from succubra bark, retired from the service of Government about 1877. The manufacture of Mr. Broughton's amorphous quinine was, however, discontinued on the departure of Mr. Broughton, and since then the whole of the bark produced on the Nilgiri plantations has been disposed of by sale. In 1873, Mr. C. H. Wood was appointed Quinologist to the Government plantation in Sikkim, and by him a process of manufacture was indicated by which the mixed alkaloids of red bark are extracted in the form of an amorphous white powder. This powder is called CINCHONA FEBRIFUGE, and up to the 31st March 1885, 70,491lb of this drug had been manufactured at the Sikkim plantation. This drug is disposed of only in India, and is

Government Cinchona Febrifuge and Quinine.

CINCHONA.

(but not powdered) and is put into wooden casks, where it is macerated in the cold with very dilute hydrochloric acid. The liquor is then run off into wooden vessels and mixed with an excess of a strong solution of caustic soda, a precipitate forms, which is collected on calico filters, and well wash-

FEBRIFUGE.

hours the liquor is carefully filtered. The filtrate is mixed with caustic soda, and the resulting precipitate collected on calico and washed with a small quantity of water, dried and powdered it is then ready for issue, and is sent out under the name of CINCHONA FEBRIFUGE."

QUININE.
1150

quinine in yellow bark can be extracted in a form undistinguishable, either chemically or physically, from the best brands of European manufacture. This can be done so cheaply that, as long as the supply of bark is kept up, quinine need never cost Government much above twenty-five rupees per pound. It is true that, at the present moment, quinine is obtainable in the open market at rates not very different from this; but that is due to entirely exceptional causes. For some time back the Ceylon planters have been up-rooting their Cinchona trees, both to save them from disease, and to make way for tea-planting which appears now to be becoming the principal industry of that Colony; and Cinchona

Method of extraction of the alkaloids from Cinchona bark by cold oil as used at the Government Cinchona Factory in Sikkim.

"In order that the oil may speedily and effectually extract the Cin-

is driven at the speed of about thirty revolutions to the minute. Any

CINCHONA.

Government Cinchona Febrifuge and Quinine

QUININE.

(about 5 parts) may be used in addition to the 8 parts of caustic soda; or caustic soda may be altogether omitted, and 15 parts of slaked lime may be used instead of it. The caustic soda is dissolved in the water and mixed with the bark. Then the oil is added, and the whole is kept

agitated, and is there thoroughly intermixed with acidulated water for

second need not be added. The quantity of

allowed to cool, and as it cools the crystals form out. These crystals are

crystalline mass obtained by filtration is then placed in small lumps on sheets of white blotting paper, and on a plate of glass or of paper.

is drained off, the precipitate is washed with a little pure water, and powdered. The powder is Cinchona Febrifuge ready for use."

TRADE.
1151

TRADE IN CINCHONA

PRESENT CONDITION OF THE BARK TRADE.—Dr. King has kindly furnished the following paragraph on this subject.—"The present condition

C. 1151

Foreign Trade in Cinchona.

CINCHONA.

of the Cinchona bark trade is one of depression. This is by no means due to any diminution of the demand for the Cinchona alkaloids, but in a great measure to the fact that an entirely new source of quinine has of late been discovered in the northern parts of South America. This

TRADE.

years been poured into the London market in enormous quantities under the designation of *Cuprea* bark. The depression is also greatly due to the enormous exports from Ceylon where cinchona is everywhere being up-rooted to make way for Tea. The effect of these flushings has been temporarily to swamp the market, the *Cuprea* crushing out the more costly Cinchona barks. The Cinchona planter, however, has only (if he can afford it) to play a waiting game for, if the importation of *Cuprea* bark goes on much longer at the present rate, *Remija* trees will soon become scarce in all easily accessible spots, and the exports from Ceylon must soon diminish. With the extension of civilization, and with the increase of wealth in tropical countries, the consumption of quinine must steadily increase, at any rate, as long as malarious fevers continue to exist in these countries.

1152

Remija plants have only recently been introduced into India. Plants are being grown in the Sikkim plantations, and Mr Lawson alludes to those in the Nilgiri plantations as too young to advance any opinions regarding the success of this new undertaking. It seems probable, however, that it may be found possible to cultivate the *Cuprea*-bark plant in regions where labour may be less expensive than is the case with the Cinchona plantations. *Remija pardiana* and *R. pedunculata* yield the *Cuprea* bark of commerce.

In the official correspondence regarding Cinchona, various opinions

plantations are not too numerous for profit." It must, however, be admitted
abandon
can see
more to
meeting,
bark h

INDIAN FOREIGN TRADE IN CINCHONA AND QUININE

The market in the London
market : e the bark as-
sumed a view of Trade
for 1875 value of the
imports of quinine in 1875-76 was Rs. 1,91,619, but it would seem that the
removal of the import duty in August 1875 has stimulated the imports
which, in the nine months of the current year, are valued at Rs. 2,28,978
It is manifest that as yet, even with the aid given by Government in the

C. 1152

CINNABAR.

Foreign Trade in Cinchona Cinnabar.

TRADE IN
CINCHONA

shape of imported quinine and the alkaloids of Cinchona produced in India at the cost of the State, this valuable febrifuge can reach only a fraction of the population."

are greater than at any previous year. The exports of Indian Cinchona bark have steadily increased for years past. In 1882-83 they amounted to 641,608 lb valued at R7,90,861, and last year 1,286,900 lb valued at R14,56,381. Thus, both in quantity and value the exports are double what they were five years ago. These facts would seem to almost point

of
bled,
2-83
the
the
Cincha
dia,
time

hoped, and indeed it has been somewhat a disappointment to those who invested in the business with expectations of large fortunes in the no distant future. The fall in prices and the competition of other countries have restricted the trade, but though its dimensions are still relatively small, the trade has been increasing."

In addition to the imports of quinine as a commercial article, reference

plantations the immense benefit conferred on the people of India by the Government effort to provide the only trustworthy specific against the

results of the Nilgiri plantations since their commencement shows a net surplus of profit of R551,743 (£55,174) "

CINNABAR.

1153

See Mercury.

C. 1153

Shun-
rofes-
pro-

Camphor Wood

CINNAMOMUM
inersCINNAMOMUM, *Blume, Gen Pl, III, 155*

Cinnamomum Camphora, *Nees, Fl Br Ind, V, 134, Wight, Ic, t 1818, LAURINEÆ*

1154

JAPAN CAMPHOR of Commerce is obtained from this tree

Syn — CAMPHORA OFFICINARUM, *Nees, LAURUS CAMPHORIFERA, Kamp ; Roxb, Fl Ind, Ed C B C, 340*

Habitat — A tall tree, with smooth, shining leaves, a native of China, Japan, and Malay Islands, introduced into the Botanic Garden at Calcutta in 1802 This is one of the sources of camphor For further information see Camphor

C. glanduliferum, Meisn ; Fl Br Ind, V, 135.

1155

THE NEPAL CAMPHOR WOOD, THE NEPAL SASSAFRAS

Syn — LAURUS GLANDULIFERA, *Wall, in Act Ser, Med and Phys, Cal, 1, 45*

Vern — *Malligiri marigiri, NEPAL, Rohu, LEPCHA, Gunserai, MECHI, Ass, Gundroi, CACHAR*

References — *Brandis, For Fl, 376, Gamble, Man Timb, 306, Voigt, Hort Sib Cal, 308, Pharm Ind, 196*

Habitat — A large tree of South Himālaya from Kumaon eastwards to Assam, the Khasia Hills, and Sylhet

Medicine — In the
tended as worthy of

MEDICINE
1156

TIMBER
1157

C. iners, Reinw, Fl Br Ind, V, 130, Wight, Ic, t 130, 122, 135

1158

cinnamomic odour and taste, and by careful drying and preparation appears capable of affording *cissia lignea* of good quality

Dr. A. Ross
Bark
1159

C. 1159

CINNAMOMUM
Parthenoxylon

Martaban Camphor Wood.

MEDICINE.

Branches.

1160

Seeds

1161

FOOD.

Bark.

1162

Leaves.

1163

TIMBER

1164

1165

states that this tree is very abundant in the Balaghat jungles of North Kanara, and that it was from this locality that the *cassia bark*, once so largely exported from that district, was obtained. The smaller BRANCHES when carefully prepared, he pronounces to be nearly equal to that of *C zeylanicum*. At his recommendation, Dr. Ross states, the Bombay Government now farms out these trees, and by this means a very considerable addition has been made to the revenue. It may be used as a substitute for cinnamon, to which, adds Dr Ross, it can hardly be reckoned inferior" (*Pharm Ind.*) "The SEEDS, bruised and mixed with honey, are used as a cathartic, and the LEAVES, and com-

(Lisboa) See

Structure of the Wood.—Billets of this tree are often sold, together with other kinds of firewood, by the wood cutters

Cinnamomum obtusifolium, Nees, *Fl Br Ind.*, V., 128; Wight, *Ic, t. 139*

Syn.—LAURUS OBTUSIFOLIA, Roxb, *Fl Ind*, Ed C B C, 339, L. CASSIA, in *Herb Ham.*

Vern.—*Tespat*, *ramtespat*, *kinton* BENG, *Bara singoli*, NEPAL, *Nupsor*, *LEPCHA*, *Patichanda*, ASS, *Dupatti*, *MECHI*, *Krowai*, *MAGN*, *Lu leng-kyaw*, BURM

References—Brandis, *For Fl*, 375, Kurz, *For Fl Burm*, II, 287; Gamble, *Man Timb*, 305, Voigt, *Hort Sub Cal* 307, *Flick & Haub*, *Pharmacog*, 528, Balfour, *Cyclop*, Simmonds, *Trop Agri*, 490; *Kew Cat*, 110

Habitat.—An evergreen tree, with grey aromatic bark, quarter inch thick, native of the outer North-East Himalaya, ascending to 7,000 feet, and of Eastern Bengal, Burma, and the Andaman Islands.

Fibre.—The "Muga" silkworm (*Antheraea assamæ*) sometimes feeds on its leaves

Medicine—Dr Kurz says the aroma of the BARK is variable, and the bark of the root of the Martaban plant is as aromatic as the best Ceylon cinnamon Dr Gimlette says the bark is "collected in Dunabaisia, a valley adjacent to that of Nepal proper, it is used in dyspepsia and liver diseases"

Food—LEAVES are aromatic, used in curry. In Assam the dried leaves are used as a spice

Structure of the Wood.—Reddish grey, moderately hard, shining, mottled on a vertical section by the medullary rays, the pores containing a gummy substance which exudes copiously on the wood being wetted Weight, 41lb per cubic foot Balfour says that the wood is useful for various purposes

C. Parthenoxylon, Meissn, *Fl Br Ind*, V., 135, Wight, *Ic, t. 1832.*

THE MARTABAN CAMPHOR WOOD

Syn.—*Laurus camphora*, Roxb, *Fl Ind*, Ed C B C, 339, L. CASSIA, in *Herb Ham.*

Vern.—*Tespat*, *ramtespat*, *kinton* BENG, *Bara singoli*, NEPAL, *Nupsor*, *LEPCHA*, *Patichanda*, ASS, *Dupatti*, *MECHI*, *Krowai*, *MAGN*, *Lu leng-kyaw*, BURM

References—Brandis, *For Fl*, 375, Kurz, *For Fl Burm*, II, 287; Gamble, *Man Timb*, 305, Voigt, *Hort Sub Cal* 307, *Flick & Haub*, *Pharmacog*, 528, Balfour, *Cyclop*, Simmonds, *Trop Agri*, 490; *Kew Cat*, 110

Habitat.—An evergreen tree, with grey aromatic bark, quarter inch thick, native of the outer North-East Himalaya, ascending to 7,000 feet, and of Eastern Bengal, Burma, and the Andaman Islands.

Fibre.—The "Muga" silkworm (*Antheraea assamæ*) sometimes feeds on its leaves

Medicine—Dr Kurz says the aroma of the BARK is variable, and the bark of the root of the Martaban plant is as aromatic as the best Ceylon cinnamon Dr Gimlette says the bark is "collected in Dunabaisia, a valley adjacent to that of Nepal proper, it is used in dyspepsia and liver diseases"

Food—LEAVES are aromatic, used in curry. In Assam the dried leaves are used as a spice

Structure of the Wood.—Reddish grey, moderately hard, shining, mottled on a vertical section by the medullary rays, the pores containing a gummy substance which exudes copiously on the wood being wetted Weight, 41lb per cubic foot Balfour says that the wood is useful for various purposes

C. 1172

MEDICINE.

Fruit

1171

oil.

1172

The Cassia Lignea

CINNAMOMUM
Tamala.

Cinnamomum pauciflorum , <i>Nees, Fl Br Ind, V, 129</i> <i>Syn</i> — <i>LAURUS RECURVATA</i> , <i>Roxb, Fl Ind, Ed C B C, 338</i> <i>Vern</i> — <i>Dinglatterdop</i> , KHASIA <i>References</i> — <i>Gamble, Man Timb 305, Fluck & Hanb, Pharmacog, 528, Simmonds, Trop Agri, 490</i> <i>Habitat</i> — Met with in the Assam Valley, Khasia Hills, and Sylhet	II73 Bark II74 Leaves. II75 TIMBER II76
per cubic foot	
C. sp. <i>Vern</i> — <i>Hmanthin</i> , BURM <i>Reference</i> — <i>Gamble Man Timb, 307</i> <i>Habitat</i> — Met with in South Tenasserim <i>Structure of the Wood</i> — White, with a pink tinge, shining, moderately hard, highly scented Weight 36 to 43lb per cubic foot It is plentiful at Tavoy and Mergui, where it is used for building.	II77 TIMBER. II78
C. sp , perhaps C Parthenoxylon , <i>Meissn (Kurz, II, 289)</i> , or Aperula Neesiana , <i>Bl (Brandis, 383)</i> <i>Vern</i> — <i>Ka away</i> , BURM <i>Reference</i> — <i>Gamble, Man Timb, 307</i> <i>Habitat</i> — Met with in South Tenasserim <i>Structure of the Wood</i> — Orange-brown, scented, moderately hard, oily to the touch It resembles the wood of <i>C. glandaliferum</i> in structure Weight 43 to 46lb per cubic foot, durable, used for house-building and shingles	II79 TIMBER II80
C. sp. (This is probably C. iners , <i>Reinw</i> , which see.) <i>Vern</i> — <i>Sinkosi</i> , BURM <i>Reference</i> — <i>Gamble, Man Timb, 307</i> <i>Habitat</i> — Met with in South Tenasserim; found by the late Mr Lee in Mergui, but rather scarce <i>Structure of the Wood</i> — Red, soft strongly scented	II81 TIMBER II82
C. Tamala , <i>Fr Nees, Fl Br Ind, V, 128</i> THE CASSIA LIGNEA OR CASSIA CINNAMOY. <i>Syn</i> — <i>LAURUS RECURVATA</i> , <i>Roxb, Fl Ind, Ed C B C, 338</i> <i>Vern</i> — <i>Dinglatterdop</i> , KHASIA <i>References</i> — <i>Gamble, Man Timb 305, Fluck & Hanb, Pharmacog, 528, Simmonds, Trop Agri, 490</i> <i>Habitat</i> — Met with in the Assam Valley, Khasia Hills, and Sylhet	II83

CINNAMOMUM
Tamala.

The Cassia Lignea.

Habitat.—A moderate-sized evergreen tree on the Hīmalaya, sparingly

DYE
Leaves
1184

Bengal the leaves and bark of *C. obtusifolium*, *Nees*, more commonly bear these names. In fact, the leaves of any species of the genus would be at once called *Teypat* by a native, but for economic purposes *C. Tamala* is superior to any of the other Indian species. The bark of this plant is the *Cassia Lignea* of Indian commerce. The *Cassia Cinnamon* of Europe is obtained from China, the source of which is still obscure. It is chiefly, however, attributed to *C. Cassia*, *Bl*, which it seems may be proved but a form of *C. Tamala*, *Nees* (Gamble reduces it to be a synonym). The true cinnamon is, however, *C. zeylanicum*, *Breyn*. The roots of *C. zeylanicum*, as also, sparingly, of *C. Tamala* and *C. obtusifolium*, yield camphor, but the true camphor plant of commerce is *C. Camphora*, *Nees*, a native of Japan.

OIL
1185

Oil.—The outer bark of the plant yields on distillation an essential oil has a to it what

Cinnamon or *Cassia Lignea* of Indian commerce is obtained from this plant. It is coarser and sold in larger pieces than the true cinnamon or bark of *C. zeylanicum*, for which it is often used as an adulterant. *Kurz* says the bark of the root is quite as good as the true cinnamon bark. In Manipur the writer found the natives on the eastern frontier regularly in

but on this point *Flückiger* and *Hanbury*, in their *Pharmacographia*, say: "Although it is customary to refer it (*Cassia* bark) without hesitation to a tree named *Cinnamomum Cassia*, we find no warrant for such reference, no competent observer has visited and described the *Cassia*—"

CASSIA
BUDS.
1186

C. 1186

The Cassia Lignea

CINNAMOMUM
Tamala

of the bark. It appears from a very old writing that the cassia buds were employed in preparing the spiced wine called *Hippocras* (*Pharmacop. Treasury of Botany*). Dr. Dymock alludes to "*Kāla nāgkesar* (known in Europe as Cassia buds)" as the immature fruits of Cinnamon — *C. alba*, and *C. cassia* buds, "exported into Bombay from

CASSIA
BUDS.in
by

... of Cassia Bark. "Cassia Bark" is a name...

CASSIA
BARK
II87

straight, even and regular, and are of a darker brown, and though some of the bark is extremely thin, other pieces are much stouter than fine cinnamon,—in fact it is much less uniform. The outer coat has been removed with less care than that of Ceylon cinnamon, and pieces can easily be found with the corky layer untouched by the knife.

"Cassia bark breaks with a short fracture. The thicker bark cut transversely shows a faint white line in the centre running parallel with the surface. Good cassia in taste resembles cinnamon, than which it is not less sweet and aromatic, though it is often described as less fine and delicate in flavour.

"An excellent medicine for the treatment of the following diseases:—

flavour

"The less esteemed kinds of cassia bark which of late years has been

MEDICINE
Bark
II88
Leaves,
II89

CINNAMOMUM
Tamala.

The Cassia Lignea

MEDICINE.

is a medicinal substance for the treatment of
 spleen, affections of the nerves or heart, pains in the womb, also in retention of urine and catamenia, and bites of serpents and poisoning by opium. "An infusion of the bark and leaves is used as a medicine in dispensary in place of true cinnamon."

CHEMISTRY.
1190

It is a substance with much
 coughs, properties
 to an essential oil, which, in a chemical point of view, agrees with that of cinnamon. It is different from the essential oil of cinnamon in the volatility of a column of water in respect to its specific gravity.

It is a colourless, inodorous substance, crystallizing in shining, brittle prisms. We have never met with it.

"If thin sections of cassia bark are moistened with a dilute solution of perchloride of iron, the contents of the parenchymatous part of the whole tissue assume a dingy brown colour, in the outer layers the starch granules even are coloured. Tannic matter is consequently one of the constituents."

salt. The colour of the substance present of cassia or cinnamon is bright blue coloration and becomes permanent only after much of the test has been done. We have not ascertained the nature of the substance that thus modifies the action of iodine; it can hardly be tannic matter, as we have found the reaction to be the same when we used bark that had been previously repeatedly treated with spirit of wine and then several times with boiling ether.

"The mucilage contained in the gum-cells of the thinner quills of cassia is easily dissolved by cold water, and may be precipitated by alcohol in the form of a precipitate."

FOOD
Bark
1191
Leaves
1192

It is much employed to adulterate true cinnamon.

C. 1192

The Cassia Lignea.

CINNAMONUM
Tamala.

Structure of the Wood—Reddish grey, splits and warps, moderately hard, close grained, slightly scented, not used Weight 39 lb per cubic foot

TIMBER.
1193

1194

thing can be gathered as to the likelihood of its being grown to a profit in Bengal as a source of Cassia bark "

FOREIGN TRADE OF CASSIA LIGNEA

TRADE.
1195

Year	IMPORTS		EXPORTS AND RE EXPORTS	
	Quantity	Value	Quantity	Value.
	cwt	R	cwt	Rs
1880-81	19,660	4 68,576	4,437	1,18,249
1881-82	9 705	1,00,891	3 863	94,408
1882-83	13 240	2 61,543	2,211	45 921
1883-84	19 917	3 84 491	5,365	1,05,310
1884-85	14,769	2,48,344	4 692	81,394

Imports for 1884-85

Presidency to which imported	Quantity	Value	Country from which imported	Quantity	Value
	cwt	R		cwt	R
Bombay	12 308	2,01 944	Aden		
Bengal	2,226	41,460	China—Hong-kong	13 537	2,24 803
Madras	235	4 940	Straits	1,212	23 536
TOTAL	14,769	2,48,344	TOTAL	14 769	2,48,344

Re exports for 1884-85

Presidency from which exported	Quantity	Value	Country to which exported	Quantity	Value
	cwt	R		cwt	R
Bombay	4 675	81,114	Persia	2,735	49 956
Bengal	13	225	Arabia	930	17,051
Sindh	4	55	Turkey in Asia	715	11 956
			Other Countries	212	3 561
TOTAL	4 692	81,394	TOTAL	4 692	81,394

Dr Dymock (*Mat. Med. W. Ind.*, 2nd Ed., 667) alludes to Cassia

of the truly Indian bark is exported

CINNAMOMUM
zeylanicum

True Cinnamon.

1196

Cinnamomum zeylanicum, Breyn.; Fl. Br. Ind., V., 131; Wight,
TRUE CINNAMON. [Ic, t. 123.

S. D.

..

..

Murray, Drugs and Pl Sind, 110; Bidse, Cat Nam Fou, 1413 LAV;
15, Waring, Basar Med., 43; S Arjun, Bomb. Drugs, 116; Baden

CAMPBOR.

1197

DYE

1198

p. from the root bark

from Nepal and from the North-
together with myrobalan, chiefly
"Ruck Dyes and Tans of
ld be referred to

OIL

1199

of cinnamon, an
is at present unknown

is at present unknown

C. 1199

True Cinnamon.

CINNAMOMUM
zeylanicum

here and there scars or holes at the points of insertion of leaves or twigs. The inner surface of the bark is of a darker hue. The bark is brittle and splintery, with a fragrant odour peculiar to itself and the allied barks of the same genus. Its taste is saccharine, pungent, and aromatic" (*Pharmacographia*, p. 525)

MEDICINE.
Bark
I200

Oil.
I201

carbon having the formula $C_{20}H_{16}$. It also contains a small quantity of benzoic acid. In medicinal properties and uses it resembles closely the oil of cloves (*Pharm Ind*). "Cinnamon is largely used in compound prescriptions. A combination of cinnamon, cardamoms, and *tejapatra* leaves, passes by the name of *trijataka*, these three aromatics being often

Special Opinions — "Powdered cinnamon in 20-grain doses is a reputed medicine in dysentery" (*Assistant Surgeon T. N. Ghose, Meerut*). "Appears to be useful in certain forms of amenorrhœa when chewed or as Oil Cinnamon" (*Surgeon-Major G. Y. Hunter, Karachi*). "The bark ground up with water into a paste is applied to the temples in neuralgia and severe headache" (*K. N. A., Dacca*). "Warm stomach cordial, carminative and astringent, useful in flatulence and diarrhœa. Cinnamon oil applied locally in very small quantity gives great relief in neuralgia headache" (*Surgeon C. M. P. Hall, Madras*).

England. It was prepared by Valerius Cordus, who stated, somewhat before 1544, that the oils of cinnamon and cloves belong to the small number of essential oils which are heavier than water, '*fundum petunt*.' About 1571 the essential oils of cinnamon, mace, cloves, pepper, nutmegs, and several others, were also distilled by Guintherus of Andernach, and again, about the year 1589, by Porta.

"In the latter part of the last century it used to be brought to Europe by the Dutch. During the five years from 1775 to 1779 inclusive, the average quantity annually disposed of at the sales of the Dutch East India Company was 176 ounces. The wholesale price in London between

with a variable proportion of hydrocarbons. At a low temperature it

C. I202

CINNAMOMUM
zeylanicum

True Cinnamon

CHEMISTRY

not examined
ed with res n

and tannic
Wittstein to
decoction of
Cinnamon
afforded to Schatzlar (1862) 5 per cent of ash consisting chiefly of the

recognition but if it should have been reduced to powder, the case is
widely different We have found the following tests of some service
when the spice to be examined is in powder Make a decoction of
powdered cinnamon of known weight of the suspected powder
each with one or two drops of
mon is but little affected but in that of cassia a deep blue black is
tara,
ell as
ed by
with

infec-
tionary, also in curry, and enters into the preparation known as *pan*

FOOD
Bark
I203
TRADE
I204

FOREIGN TRADE OF CINNAMON

Year	IMPORTS		EXPORTS AND RE EXPORTS	
	Quantity	Value	Quantity	Value
	lb	Rs	lb	Rs
1879-80	1 785	484	202	24
1880-81	7 707	3 511	19 432	4 833
1881-82	2 244	512	67 466	14 436
1882-83	18 731	3 641	27 768	11 068
1883-84	13 687	2 640	35 181	9 330

Detail of Imports 1883-84

Province into which imported	Quantity	Value	Country whence imported	Quantity	Value
	lb	Rs		lb	Rs
Bengal	9 6	437	Straits Settlements Other Countries	11 924	2 034
Madras	12 547	2 143		1 763	606
British Burma	224	60			
TOTAL	13 687	2 640	TOTAL	13 687	2 640

CINNAMOMUM
zeylanicum

True Cinnamon.

CHEMISTRY.

not examined
ed with resin

and tannic
Wittstein to
decoction of
Cinnamon

afforded to Schatzlar (1862) 5 per cent. of ash consisting chiefly of the
og, 526).
lia remark that "Cassia
very commonly substi-
re is no difficulty in its
1 to powder, the case is
widely different we have found the following tests of some service
when the spice to be examined is in powder Make a decoction of
powdered cinnamon of known genuineness, and one of similar strength
of the suspected powder When cool and strained, test a fluid ounce of
each with one or two drops of tincture of iodine A decoction of cinna-
mon is but little affected, but in that of cassia a deep blue black tint is

FOOD
Bark
I203
TRADE
I204

tionery, also in curry, and enters into the preparation known as *pan*.

FOREIGN TRADE OF CINNAMON

Year	IMPORTS		EXPORTS AND RE EXPORTS	
	Quantity	Value	Quantity	Value
	lb	R	lb	R
1879-80	1 785	484	202	14
1880-81	7,707	3 511	19 432	4 833
1881-82	2,244	512	67,466	14 436
1882-83	18,731	3,641	27,768	11,068
1883-84	13,687	2 640	35 181	9 330

Detail of Imports, 1883-84

Province into which imported	Quantity	Value	Country whence imported	Quantity	Value
	lb	R		lb	R
Bengal	916	437	Straits Settlements Other Countries	11 924	2 034
Madras	12 547	2,143		1,763	606
British Burma	224	60			
TOTAL	13 687	2,640	TOTAL	13,687	2 640

CITRULLUS
Colocynthis

False Pareira Brava; Colocynth

MEDICINE.

Root
I207
Bark
I208
Leaves
I209

wedge-shaped rays, which are often irregular, scattered, and indistinct. The axis is not often eccentric. In *Cissampelos Pareira* the root and stem are nearly alike in structure, and in transverse section there are concentric rings" (*Year-Book of Pharm*, 1873, 30)

Medicine.—The dried root and bark are used as mild tonics and diuretics in advanced stages of acute and chronic cystitis and catarrhal affections of the bladder, also exercises apparently an astringent and sedative action on the mucous membranes of the genito-urinary organs. They are generally administered in the form of decoction and extract. The leaves are applied to abscess. Ainslie writes; "The leaves of this plant are considered by the *vytians* as of a peculiarly cooling quality, but the root is the part the most esteemed, it has an agreeable, bitterish taste, and is considered as a valuable stomachic. It is frequently prescribed in the latter stages of bowel complaints, in conjunction with aromatics. *Cissampelos Pareira* has been very highly extolled by several writers for its medical virtues, particularly by Sloane, Marcgraaf, Barham, and Wright. The first speaks of the efficacy of the leaves as a vulnerary for

Special Opinions.—? Used locally in cases of unhealthy sores and

CHEMISTRY.
I210

a yellow bitter principle, a brown colouring matter, starch, an azotised substance, and various salts of ammonia and lime" (O'Shaughnessy).

to Fluckiger,
Warden, Pro-

Cissus carnosa, Lam, see *Vitis carnosa*, Wall, AMPELIDÆ.

C. discolor, Blume, see *V. discolor*, Dals.

C. edulis, Dals., see *V. quadrangularis*, Wall.

C. pedata, Lamk, see *V. pedata*, Vahl.

CITRULLUS, Schrad.; Gen. Pl., I, 826.

I211

Citrullus Colocynthis, Schrad; Fl. Br. Ind, II., 620; Wight, Ic,
1 498; CUCURBITACÆ.

COLOCYNTH, Eng.

C. I211

The Water melon

CITRULLUS
vulgaris

It, bitter taste, and contain 17 per cent of fat
t, if rubbed, emit a very unpleasant smell "

armacy (1873) gives the following account

FOOD
Fruit
1217

marogelie, ...
limba (Arabia) ...
Tung-ga ...
indrak G ...
Duk, The ...
Paycumati, ...
utite, TAN ...
veri pách-cha ...
kayt, KAY ...
varuni méliká ...
tolkh, khar-busaka ...
khia si, khia si ...

References — Thwaites ...
101, Stewart Ph Pl ...
Ind, 94, Modern ...
Pharmacog, 215, ...
Mat Med Hind ...
Mat Ind, 1 84, ...
and Drugs Sind, 39 ...
Year Book Pharm, 1891 ...
U Pl 139, Baden Powell ...

"st freed from pulp by roasting
sacks, and then deprived of
by grinding and winnowing.
old only taste, but several, if
The kernels contain about
buminous substances, be-
e be regarded as a suffi-

I with cold water, dried
other ways as food "
as food for horses
cold winter nights
ously been pierced
aters, until all the

Kernels
1218

Habitat — An annual found wild in
tral, and South India. It is the wild gourd
The plant cannot be said to be systematic
India, the fruits are collected from plants
desert tracts of North-West India (Duthie and ...)
Oil — Yields, according to Ainslie, a clear,
the southern provinces for burning in lamps
used to dye the hair

h The people
ch they use for
nts the camels
icog, 297)

DOMESTIC.
Tooth-
brushes.
1219
Tar
1220
1221

ascites, enlargements of the abdominal viscera, urinary ...
&c An oil prepared from the seeds of Indian Colocynthis ...
ening grey hairs. A poultice of the root is said to be useful ...
of the breasts." (U C Dutt, Mat Med Hind) According to ...
madan writers, Colocynthis is a drastic purgative, removing ...
parts of the system. They recommend the fruit, leaves and ...
used in costiveness, dropsy, jaundice, colic, worms, elephantiasis ...
acts as an irritant on the uterus, and its fumigation brings on the ...
flow. The author of the *Makhsan* describes a curious mode of ...
tration. "A small hole is made at one end of the fruit and ...
are introduced, the hole is then closed, the fruit enveloped in

B C, 700
samanka,
ana, P 8
Turbus,
Pittha,
kachrehn,

Pl, 95,
2, U. C
Ainslie,
Pb Pr,
1, 701,
this &
3, Dic,
2

espe-
well
uch
the
ten
son
in
er-

CITRULLUS
Colocynthis

Colocynth.

Colocynth is rarely employed alone, it is generally given in combination with other purgatives and carminatives. It commonly causes griping when used alone, in excessive doses it produces inflammation of the intestines and even death. The principal efficient forms for the use of this drug are the compound extract of Colocynth, compound Colocynth pill, and Colocynth and henbane pill. (*Bentley and Trim, Med. Pl., 114*) From the pulp a watery extract is prepared, which is much

compound extract of Colocynth the supply of the medical stores in Panjab the fruit is extensively employed as a purgative for horses. The pulp of the fresh fruit mixed with warm water, or the dried pulp with *aywain*, is reckoned a special remedy in cholera. The dried root reduced to powder is given as a purgative (*Bellew*). Stocks says the root and the juice are both used medicinally in Sind. In a report of the drugs shown at the late Colonial and Indian Exhibition from Baroda, the properties of the fruit and root are given in very nearly the same terms as above, so that the knowledge of this drug seems very extensively diffused over India.

Special Opinions — “Used in dropsy and amenorrhœa” (*Native*

CHEMISTRY
1216

principle remains partly in the aqueous liquid, partly in the resin, from which the *Colocynthin* is to be extracted by boiling water. The whole solution was then concentrated and mixed with carbonate of potassium, when a thickish viscid liquid separated. Hubschmann dried it and redissolved it in a mixture of 1 part of strong alcohol and 8 parts of ether. After treatment with charcoal, the solvents were distilled and the remaining bitter principle removed by means of water. This on evaporating afforded 2 per cent. of the pulp of a yellow, extremely bitter powder, readily soluble in water or alcohol, not in pure ether. Colocynthin is precipitated from its aqueous solution by carbonate of potassium. Colocynthin was further extracted by Lebourdais (1848) by evaporating the aqueous infusion of the fruit with charcoal, and exhausting the dried powder with boiling alcohol.

“Again, another method was followed by Walz (1858). He treated alcoholic extract of colocynth with water, and mixed the solution, firstly,

purgative, it is decomposed, according to Walz, by boiling with hydrochloric acid, and then yields colocynthein, $C_{44}H_{64}O_{12}$, and grape sugar. The extract of colocynth is dried with

The Water melon

CITRULLUS
vulgaris

even when crushed, but a faint, bitter taste, and contain 17 per cent of fat oil.

"The fresh leaves of the plant, if rubbed, emit a very unpleasant smell" (*Pharmacog*, p 296)

Food—The *Year Book of Pharmacy* (1873) gives the following account of the fruit as a food substance—

"The FRUIT, which is about as large as an orange, contains an extremely bitter and drastic pulp, from which colocynth is obtained. This pulp is said to be eaten by buffaloes and ostriches, but is quite unfit for human food. The seed kernels, however, which contain but a very small quantity of bitter principle, are used as food by some of the natives of the African desert and their A sin

The kernels contain about albuminous substances, therefore be regarded as a suffi-

"The KERNELS are heated to boiling, then washed with cold water, dried and powdered and eaten with dried dates, or used in other ways as food" (*Bentley and Trimen*). "The fruits are often used as food for horses in Sind, cut in pieces, boiled, and exposed to the cold winter nights. They are made into preserves with sugar, having previously been pierced all over with knives, and then boiled in six or seven waters, until all the

FOOD
Fruit
I217

Kernels.
I218

DOMESTIC
Tooth-
brushes.
I219
Tar
I220
I221

Citrullus vulgaris, Schrad, *Fl Br Ind*, II, 621

THE WATER-MELON.

Syn—*CUCURBITA CITRULLUS*, Linn, *Roxb*, *Fl Ind*, Ed C B C, 700

Vern—*Tarbua*, *tarbus*, *turmua*, *karbu*, *halinda*, *hindwana*, *samanka*,

kers, paye, *ina yaithi* BURN

References—*Dal*, & *Gibb*, *Bomb*, *Fl*, 103, *Stewart*, *Pb*, *Pl*, 95.

Habitat—Cultivated in gardens

CITRULLUS
vulgaris

The Water melon.

HISTORY

History.—The water melon is native of Southern Italy, while
Seringi
wards
saw dis
of wild animals eagerly devoured the
the ancient Egyptians, as appears from
only received the plant in the tenth c
Orig Cult Pl, 263)

Oil.—The seeds yield a clear, bland, pale coloured, limpid oil, used for
burning in lamps, and probably also as an edible oil (*Cooke*)

Me

says t

use

and ar

remar

and a

ministered it with good results

Special Opinion.—§ "Cooling as well as a diuretic" (*Assistant Sur-
geon Anund Chunder Mukerji, Noakhally*)

Food.—The FRUIT is large, ovoid, green, and smooth, the flesh is whit-
ish yellow or red. The SEEDS are compressed and variable in shape and
colour, they are sometimes dried and the kernels eaten. Stewart says
they are eaten parched with other grain. In the North-West Provinces

such numbers as to form for some months in the year no small part of the
food of the scanty population. The seeds of these and of other cucurbita-
ceous plants cultivated in gardens are ground during times of scarcity into
a kind of flour" (*Raj Gas* 31). The water-melons of the North-Western
Provinces are famed all over India and are used as refrigerants, and as a
sherbet ingredient.

Var *fistulosus*, *Stocks*, *Duthie & Fuller*, *Field and Garden Crops*,
N-W P, II, 46, Plate XLVII

In the *Flora of British India* *C. fistulosus* has been given as a syn-
onym to *C. vulgaris*, *Schrad*, but it seems desirable to retain it as a
variety.

Vern.—*Tandús, tendu, tind albinda, tensi*, *N W. P.*, *Tinda, albinda*,
dilpasand, Ps, *Meh, trindus, dilpasand, tinda, albinda*, *SIND*

References.—*Stewart, Pb Pl*, 96. *Balfour, Cyclop.*

Habitat.—In the North-West Provinces this fruit is sown some
little time before the rains, the fruit ripening during the rains. "Culti-
vated in Sind from April to September, generally in the same plot of
ground with common melons, gourds, and cucumbers. In the North-
s before
r vege-

MEDICINE

1228

FOOD

1229

Pickle

1230

C 1230

The Genus Citrus.

CITRUS.

black
lt, and
anner.
ls, and
cks, 121

Hooker's Journal of Botany, quoted by Brandis and Thwaites,

CITRUS, Linn, Gen. Pl., 1., 305.

1231

Rut

This genus comprises 5 tropical Asiatic species and 2 Australian.
The different varieties of the Orange, the Lemon, the Lime, and the Citron have been critically examined by a large number of patient and careful observers, but, it must be admitted, with but indifferent results. Brandis, after presenting a concise and pregnant account of the Indian

results regarding the spread and changes of arborescent species under cultivation." Since these words were penned, it is feared we have not advanced very far towards a solution of the problems which hinge upon the nativity of the orange and the lemon. Shortly after the appearance of Dr. Brandis' *Forest Flora*, Dr. Rice of New York published in *New Remedies* a most interesting account of the genus Citrus, but without

Risso, as a synonym under *C. nobilis*, *Lour.* (the Mandarin)—a species

mamillate in the oranges. Species characterised by the degree of development of a certain feature must naturally under cultivation become hopelessly intermixed, hybridisation rendering it almost impossible to distinguish the forms. This is true in its fullest extent with the mem-

C. ... long with
the writer
means so
the limes
aurantium,
C. Auran-
er of the
of *C. Medea*. Whether *K* is the correct name or the name of

C. Aurantium by their very much smaller flowers. It is usual to regard
larins,
culi-
navia
the *Khasia* hills but
of good Mandarins as
The true Mandarin,
but it would be in-
teresting to have the question of its relation to the sweet lime more clearly
established. According to *Kurz*, these two cultivated plants are one and
the same species, *C. nobilis*, being much cultivated all over Burma. This
conclusion may not, however, be regarded as satisfactory, from the fact
that the Mandarin and the sweet lime are both extremely common in the

subsequently), may be found useful:—

- *Young shoots and leaves perfectly glabrous; transverse vesicles of the pulp concrete.*

† A shrub, young shoots purple; petiole more or less naked, petals generally tinged with red; flowers

CITRUS
Aurantium.

The Sweet Orange.

Vern — *Nárangí, sanglara, nárenj, naríngi, náránge, sunthura, amrit-phal, kumla nebu*, HIND.; *Kamlá nembu, nárunji, nárengá*, BENG;

Re

Pharmacop., 144, U. S. Dispens., 151a Eu., 261; *Pent. & 1 rim*, Med Pl., 51; U. C. Dutt, *Mat. Med. Hind.*, 127; Dymock, *Mat. Med. W. Ind.*, 107, *Ainslie, Mat. Ind.*, I, 281; O'Shaughnessy, *Beng. Dis.*

1025-7, *Presse, Perfumery*, 159, *Dufour, Cylind.*, 216, *Smith, Dic.*, 300;

Habitat.—Cultivated in most parts of India, but specially so in the

oranges, but there are large tracts where none or inferior kinds only are produced. In India the fruit generally ripens between December and March, according to the climate of the locality. A variety which flowers twice a year (February and July), and yields two crops—the first

HISTORY.

ning of the Christian era." It was, according to some authors, taken to Europe by the Portuguese about 1548, the first tree having stood for some time at Lisbon. From this point, the cultivation of the sweet orange spread to Rome and along the Mediterranean. DeCandolle, however, is of opinion that the sweet orange may have reached Europe before the

The Sweet Orange.

CITRUS
Aurantium

HISTORY.

sweet and the bitter orange were unknown to the Romans. Whether or not the Portuguese deserve the credit of introducing the orange to

that the orange is a native of China, the names given to the various forms are represented by a particular character which occurs in the most ancient Chinese writings, whereas the names given to the pumelo and the lime are of a much more modern character.

Dr. Bonavia has given the subject of the Indian Oranges, Limes, and Lemons more careful consideration than any other Indian authority. In

derived from the Sanskrit. It is, according to the best authors, a Persian corruption, and it can hardly be doubted that Santara is derived from Cintra—a town famous for its fruits. Yule-Burnell say "As early as the beginning of the fourteenth century we find Abulfeda extolling the fruit of Cintra. His words, as rendered by M. Re naud, run "Au nombre des dependances de Lisbonne est la ville de Schintara, à Schintara on recueille des pommes admirables pour la grosseur et la gout" That these *pommes* were the famous Cintra oranges can hardly be doubted. Baber (*Memoirs of Zehir-ed din Muhammed Baber, Emperor of*

me of Sangtarah, which is, for a species of the fruit nge in Portugal would ac e name Cintra, but for the of Portugals has adhered as might be quoted in sup- rother fruit in (Tarang), but the skin of , page 328) Kirkpatrick, in Nepaul Santola orange as ys, "I take to be a corrup-

The Sweet Orange.

CITRUS
Aurantium.

or economic point of view, it is of little consequence whether, a sweet orange be referable to *C. Aurantium* or *C. nobilis*; we may therefore

CULTIVA-
TIONTHE RACES
OF SWEET
ORANGES.

Race 1st,
Santara.

1234

ber, December, and January.

Vern — T

form

BENG,

kompho

tengo,

latter in N-W P for the same orange), Santara, C P. (near Wardha)

two crops are obtained, one ripens in spring known as the miragahar,

Mr. Morris (in his Godavery District, Madras Presidency) says: "a

name with it. The plant could scarcely have been indigenous to both

ora

and

The ora is a thick tree, met with in the Godavery District, Mr

Race 2nd,
Keonia
1235

darker colour, thinner, and adhesive (e.g. jacket not loose). This is the orange that comes latest into the Calcutta market. It is plucked about

CITRUS
Aurantium.

The Sweet Orange.

RACES OF
SWEET
ORANGES

Before proceeding to discuss the third class of sweet oranges referred to by Dr. Bonavia it may be as well to refer to another author. Mr. Atkinson says of Kumaon: "The sweet orange is the form most usually cultivated, and there are several local varieties, some named after the localities in which they are produced, and others according to specific

names derived from a common source, and that the oranges they represent should be isolated from those designated *Santara* or some derivative from

doubts may be entertained as to whether it is a coincidence not in connection with any other names so much alike as remote parts of India

and be used in the same manner as the oranges of the same name could be used in the same manner as the oranges of the same name may be found in the same manner as the oranges of the same name That writer says: "The orange is cultivated in the same manner as the oranges of the same name and with great success in the same manner as the oranges of the same name late, acumin

possible to avoid the conviction that too strong opinions have, by all

mentioned regarding the vernacular names as given to the various forms of the Indian sweet oranges of cultivation (and even to the supposed wild oranges of Nepal), is sufficient to justify the conclusion that the whole subject is still involved in the utmost obscurity. A scientific exploration

Rare 3rd,
Malta.
1236

The oranges of Burma, for example, may have been derived from the indigenous plant, a species to which the name of the same name is given. Be that as it may, a

of the blood red forms, India might obtain a supply of oranges in

The Sweet Orange

CITRUS
Aurantium.

the hot season, the time when these fruits would be most acceptable. Speaking of the Gujranwala oranges Dr. Bonavia says Colonel Clarke introduced these from Malta in 1852-56 Dr. Bonavia himself in 1863, and Mr C Nickels 2. Prior to the Mutiny blood it there must have been earlier From these centres, however, the cultivation of the red oranges has been greatly extended, so that they are now met with in most districts in Upper India. At Poona a blood orange is grown under the name of the *Mussembi*, a name given

RACES OF
SWEET
ORANGES

1237

opinion, simply perfect I thought them equal to that of the blood oranges of Malta." "Mr. Steel states that the soil on which they grow is But the real secret, he thinks, is

there is a suitable soil and climate
is also skill to turn these materials to

oranges, and therefore would not compete with the *Sattara* oranges, which flood the Calcutta and Bombay markets from Silhet and Nagpur." Mr. Steel reports that in February they are "barely ripe, and would remain on the trees till the middle of March Last year, some by careful packing were kept in good condition till July"

Race 4th.
Mandarin.
1238

the true Mandarin, while found in Ceylon, does not exist in India Mr. C B Clarke, on the other hand, says the cultivation of this form is rapidly extending in the Khasia hills Dr. Bonavia recommends its introduction in "the highlands of Bengal," "where it would be out of the influence of the hot winds," which have killed or rendered useless all the plants grown in Upper India.

in
the
Nagpur

1.—ORANGES OF SILHET AND THE KHASIA HILLS — A most instructive paper appeared on this subject in the Journal of the Agri-Horticultural Society of India, from the pen of Mr. C. Brownlow (*Vol I, Part IV, New*

Silhet.
1239

C. 1239

CITRUS
Aurantium.

The Sweet Oranges of Silhet.

ORANGE PRO-
DUCTION IN
INDIA.

Series, 1869, p. 372. Mr. Brownlow gives the fullest particulars regarding the "Orange groves of Shalla," his paper being a model after which all

tion, collection, and transport are next duly disposed of. Indeed, so admirably has Mr. Brownlow fulfilled his task that any abridgment of his paper must mar its usefulness. The limited space at the writer's disposal precludes the reproduction of the entire paper, and the reader who may be specially interested in this subject is therefore referred to the original;

Soil.

back again, and that the floods inundate the land in spring tides, thus ruining the soil and preserving its fertility. "The land is flat, having a slight slope away from the river, there are a few points that rise above the gen-

received equal to dry 100

Soil dried at 212°F.

Alumina	6.09
Peroxide of iron	4.93
Lime	1.19
Magnesia	1.13
Alkalies (by difference and loss)	1.80
Silica solution	1.15
	12.29
	3.49
	5.66
	73.56
	100

The Sweet Oranges of Silhet.

CITRUS
Aurantium.

"It will be observed that this is a very siliceous soil, proceeding from the decomposition of siliceous rocks alone. It contains no carbonate of limes and is a very open and porous soil."

CULTIVATION—The seed is sown in January and February, thickly in troughs or boxes in about 6 inches of soil. These seed-boxes are raised at intervals to catch the rain and are often protected by nets.

ORANGE PRO-
DUCTION IN
INDIACultivation.
1240

root They are transplanted into a nursery in the grove; here they remain until retransplanted to their destined places in the grove. The system seems defective and the nursery is only once a year weeded, viz., in October. Grafting is quite unknown, and no care seems to be spent on the selection of the seed.

COLLECTION AND PRUNING—Each collector has a ladder, about 20 feet long, made of light bamboo. A coarse net bag, held open at the mouth by a cane ring, depends on his back by a strap passed over the right shoulder and chest. Into this he throws the oranges and before descending he removes the withered leaves and dead branches, or cuts out boughs injured by the loranthus parasite that does such damage to the plants. "The orange trees receive no other handling than the above, they are never systematically pruned or thinned, and are allowed to retain just what fruit they set, and yet the crop turns out wanting neither in size, flavour, nor abundance. Contrast with this the elaborate summer and winter pruning of the French gardens and the systematic cultivation and

Collection and
Pruning.
1241

the dogs have come by habit to relish this food.

TRANSPORT TO THE PLAINS—The oranges so collected are taken

Transport
1242

quarry are sold by contracting for rice, fish, &c., to the Muhammadan boatmen at R6 a *son*, being R4 less than the oranges at the Shalla groves, and yet this includes the cost of cultivation, labour of plucking, and carriage to the river.

TRADE IN SILHET ORANGES.

Mr G. Stevenson, Deputy Commissioner, Silhet, has furnished the following tabular statement.—

TRADE
1243

						BOAT TRAFFIC.	
						Quantity in maunds	Value in Rs.
1880-81	1,20,398	2,40,796
1881-82	1,46,392	not kn wn
1882-83	1,02,631	1,28,283
1883-84	1,14,969	2,27,662
1884-85	1,20,884	2,47,352

CITRUS
Aurantium.

The Sweet Oranges of Nagpur.

TRADE

Dr Bonavia, comm

about 1,21,095 maunds

of rupees, in favourable

to be equal to about 8

Bonavia further adds

small Taking 8,05,36c

low, the figures would be 2,41,60,800, or about 210 oranges to the maund"

Nagpur
1244

II —ORANGES OF NAGPUR IN THE CENTRAL PROVINCES —We have already given several passages that refer to the so-called wild oranges both of Nepal and the Central Provinces. It will only be necessary further to give here a brief account from the pen of Mr. J B Fuller, as published by Dr Bonavia, in order to place before the reader a comparative sketch of these groves to complete what has been said of the Khásia hills. These two localities represent the bulk of the orange production of India. Mr Fuller says — 'Within the last twelve years many new orchards have been planted in Nagpur, Kamptee, and other parts of the district, and orange cultivation is now spreading rapidly in other districts of the Province. There is a great demand for the Nagpur oranges in Bombay, and considerable quantities of the fruit are annually exported to this and other places. In the year 1885, 22 609 maunds of orange fruit were exported from Nagpur station, out of which 21,400 maunds were exported to Bombay alone."

re to repeat that the North-West Pro-
Nepal, Delhi, and to some extent also
and Burma are practically dependent
orchards, Madras drawing largely

Properties and Uses—

GUM

1245

Gum —The orange tree is said to yield a gum of no importance. A sample was sent from Masulipatam to be shown at the Madras Exhibition in 1855

MEDICINE
Rind
1246

Medicine —The *Pharmacopœia of India* treats the sweet and bitter

external applications

cel is useful for
Orange poultice

is recommended in some skin affections, such as psoriasis, &c. Oranges are considered to be alexipharmic and disinfectant, orange-water stimulating and refreshing. The essence is extracted by oil from the rind and flowers, and is used as a stimulating liniment' (*Dr Dy-mock, Ut Med W Ind*)

Ainslie makes the following remarks "Oranges are in great repute amongst the Hindu physicians, who suppose that they purify the blood,

C. 1246

The Bitter or Seville Orange

CITRUS
Aurantium.

allay thirst in fevers, cure catarrh, and improve the appetite. A sherbet Europeans made with the rind a valuable addition to bitter infusions in cases of dyspepsia and flatulence. The rind pulverised and added to magnesia and rhubarb affords a grateful tonic to the stomach in gout and dyspepsia. The roasted pulp is an

MEDICINE

FOOD
1247

grown in and about Delhi is on the average larger, but more spongy,

1200 The produce of one tree ranges from 500 to 6000 fruits a year, and the tree sometimes grows to a height of 50 feet, with a trunk 12 feet in circumference

Structure of the Wood —Yellowish white, moderately hard, close and even grained

TIMBER
1248

Var 2 Bigaradia, *Fl Br Ind*, I, 515 (For var 1st, see p 335 and for 3rd, p 347)

Botanical Diagnosis —Petiole short winged, flowers large, strongly scented, rind very aromatic, pulp bitter

Var 2
Bigaradia.
1249

THE BITTER OR SEVILLE ORANGE, BIGARADIER, *Fr*; ARANCIO FORTE, *It*; POMERANZE, *Ger*

Syn —*C VULGARIS* RISSO *C BUXIFOLIA*, *Poir*

Habitat —The bitter orange is very extensively grown in the warmer parts of the Mediterranean, especially in Spain and Malta. In India it does not seem to be cultivated except in gardens but it is believed by m Garhwal reas extends entirely from

Marmalade
1250

Marmalade is chiefly made from the rind of this species, but it is manufactured from the true rous bitter indigenous employed for grafting other oranges. Definite information cannot be obtained as to the extent the Seville orange is being cultivated in India

OIL OF NEROLI

Oil and Perfumery —Essential oils are obtained from most of the species of the Citrus family. Sir W O Shaughnessy, speaking of the sweet

OIL
1251

C. 1251

CITRUS
Aurantium.

The Bitter or Seville Orange.

PERFUMERY

1252

Bigarade, and the oil from the flowers of the sweet variety bears the name of *Essence de Nérols Pétale* or *Nérols Louce*. This statement is opposed, however, to the opinion given by almost every other writer, the neroli otto from the sweet orange being used only as an adulterant to that from the bitter. The fresh flowers of the *Bigaradia* orange yield on distillation *Essence de Nérols Bigarade*, and if the sepals are carefully removed from the flowers, the essence is known as *Essence de Nérols Pétale*. The latter is finer and much more expensive than the former. From the seeds *Essence de Petit Grain* used to be manufactured, but this is now entirely distilled from the *Essence de Petit* most species of

1253

orange leaf to adulterate neroli otto. The water which passes over with the oil during distillation constitutes, when separated from the oil, *Orange-flower Water* (see below).

The extraction of Neroli oil is chiefly carried on at Grasse, Cannes, and Nice, in South France, also in Algeria. In France, about 20,000 cwt of the flowers are annually distilled. The sweet variety yields but half the amount of oil which may be obtained from the bitter, as much as

Neroli Cam-

phor

1254

Eau de
Cologne

1255

1256

It are used to an enormous
and Eau de Cologne
" is mainly consumed

1257

among the distillers of essential oils. It is largely used in pharmacy. There are three sorts of orange-flower waters found in commerce. The first is distilled from the flowers, the second is made with distilled water

C. 1257

The Bergamot Orange.

CITRUS
Aurantium.

PERFUMERY.

and neroli, and the third is distilled from the leaves, the stems, and the young unripe fruit of the orange tree." (*Presse*) "As met with in commerce, orange-water is colourless or of a faintly greenish-yellow tinge, almost perfectly transparent, with a delicious odour and a bitter taste." (*Pharmacog*)

ESSENTIAL OIL OF ORANGE PEEL—"Largely made at Messina, and also the south of France. It is extracted by the sponge, or by the *écuelle* process, partly from the Bigarade and partly from the sweet or Portugal Orange, the scarcely ripe fruit being in either case employed. The oil made from the former is much more valuable than that obtained from the latter, and the two are distinguished in price-currents as *Essence de Bigarade* and *Essence de Portugal*.

"These essences are but little consumed in England, in liqueur-making and in perfumery." (*Pharmacog*)

Var 3. Bergamia, Fl Br Ind, I, 515

THE BERGAMOT ORANGE

Syn —C AURANTIUM, var. BERGAMIA, W & A Prodr, p8; C LIM-

Var. 3
Bergamia
I258

Lamiya-si, in tam bayu-si, BURN

References —Brandis, For Fl, 54, Dals & Gals, Bomb Fl, Supp, 13,
Voigt, Hort Sub Cal, 142, Pharm Ind, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935, 936, 937, 938, 939, 940, 941, 942, 943, 944, 945, 946, 947, 948, 949, 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, 984, 985, 986, 987, 988, 989, 990, 991, 992, 993, 994, 995, 996, 997, 998, 999, 1000.

Habitat—The Bergamot Orange is cultivated near Reggio in South Calabria, in Sicily, and in the south of France, but it is only rarely met with in India. It may be doubted how far the above vernacular names given to it are correct. The fruit, when full grown, is still unripe and green, they are sometimes known as green oranges. Some of the green oranges met with in India (and already alluded to, p 340) may belong to this variety.

BERGAMOT OIL

Oil—The rind of the fruit yields on expression the oil known under the name Bergamot. For this purpose the fruits are used, and one hundred of them are said to produce about three ounces of the otto. Formerly the oil was extracted by distillation or by expressing the rasped rind, but these processes have been superseded by the *écuelle*, a special instrument described in *Spons' Encyclopædia*, page 1457.

General Characters of the Oil—The oil, as produced by the machine

OIL.
I259

pen-
wity
rant
pro-
The oil, as produced by the machine

Chemical Composition—The authors of the *Pharmacographia* say: "If essential oil of bergamot is submitted to rectification, the portions

CHEMISTRY.
I260

C. I260

CITRUS
decumana.

The Bergamot; The Pumelo or Shaddock.

CHEMISTRY

that successively distil over do not accord in rotatory power or in boiling point—a fact which proves it to be a mixture of several oils, as is further confirmed by analysis. It appears to consist of hydrocarbons, $C_{10}H_{16}$, and their hydrates, neither of which have as yet been satisfactorily isolated. Oil of bergamot, like that of turpentine, yields crystals of the composition $C_{10}H_{16} + 3H_2O$, if 8 parts are allowed to stand some weeks with 1 part of spirit of wine, 2 of nitric acid (sp gr 1.2), and 10 of water, the mixture being frequently shaken.

Essential Oil
1261

Properties and Uses—The oil of bergamot is much employed in perfumery. It has stimulant properties, but is rarely used in medicine. It is sometimes employed to give an agreeable odour to ointments and other external applications.

Essent. Oil of Bergamot, *Pharm. Soc. Lond.*, *Pharm. Soc. New York*.

can be obtained.

MEDICINE
Juice
1262

Medicine—The JUICE of the fruit possesses properties similar to those of lemon juice (see under *Citrus Medica*, v. *Limonium*). It is often preferred to lemon juice, as the fresh juice can be readily obtained in nearly all parts of the tropics, and as the preserved lemon juice is less effectual. It is useful as a refrigerant drink in small-pox, measles, scarlatina, and other forms of fever. It may also be taken with advantage in cases of hemorrhage from the lungs, stomach, bowels, uterus, kidneys, and other internal organs. (*Waring, Bizar Medicines*.)

1263

Citrus decumana, Linn., *Fl. Br. Ind.*, I, 516

THE SHADDOCK, PUMELO, or POMPELMOS, THE FORBIDDEN FRUIT, PARADISE APPLE, *Eng.*, POMPELMOSE, *Fr.*, POMPELMOES, *Sp.*

The word Pumelo is a contraction of "pomum melo," the melon apple.

Vern.—*Maha nibu*, *chakotra*, *batari*, *nebu*, *sadaphal* HIND, *Bāṭi*, *nebu*, *mahā nambu*, *chakotra*, *bator*, *nebu*, BENG, *Chakotra*, *Pb.*, *Bjor*, *nebu*.

References—*Fl.*, 58, 572, *ice Citrus*, *Supp.*, 12, *Horl. Sub.*, *Mat. Med.*, *Cal.*, 141, *Fluck. & Land.*, *Indurmu*, 2, 111, *U. S. Disp.*, 127, *O. Shaughnessy*, *Beng. Dispens.*, 232, *Dalen*, *Lowell*, *Pb.*, *Pr.*, 334, *Cook*, *Gums and Gum resins*, 13, *Atkinson*, *Him. Dist.*, 710, *Atkinson*, *Gums and Gum resins*, 16, *Birtwood*, *Bomb. Prod.*, 141, *Lis*.

C. 1263

The Pumelo, The Citron

CITRUS
Medica.

boa, U Pl Bomb, 148, Smith Dict, 375 Treas of Bot Ure, Dict Arts and Manuf, III, 765 Kew Off Guide to the Bot Gardens and Arboretum, 64 65, Trop Agri, 117, Simmonds Trop Agri, 44

Habitat—A native of the islands of the Malay Archipelago, more particularly abundant in the Friendly Isles and Fiji. Introduced into India from Java and into the West Indies by Captain Shaddock, hence the name *Shaddock*. It is cultivated in most tropical countries. In India and Burma it is a common fruit tree. It is, however, more frequent in Bengal and Southern India than in the North-West Provinces. The vernacular name *Batavi nebu* suggests its having been originally brought from Batavia. "The fruit is very large, weighing sometimes ten to twenty pounds, roundish, with a smooth pale-yellow skin, and white or reddish sub acid pulp. When the fruits attain their largest size, they are called pompoleons, or sompilmousses, those of the smallest size form the 'Forbidden fruit' of all the English markets" (*Treasury of Botany*)

Gum—Said to yield scantily an unimportant gum. In 1855, Lieutenant Hawkes sent to the Madras Exhibition a sample of this gum (*Cooke*)

Medicine—Mr Baden Powell says that the FRUIT is nutritive and refrigerant. It contains sugar and citric acid, with much essential oil in the peel. The leaves are said to be useful in epilepsy, chorea, and convulsive cough.

Food—This tree is a favourite with the natives of India, as it gives fruit all the year round, flower unripe and ripe fruits may be seen on the same tree at once. There are two varieties, one with whitish, and the other with reddish pulp. Besides, the individual fruits differ from one another in size, shape, and colour, and also in quality.

□ Bonavia (in the paper to "The best pummelows of the Bombay market")

GUM
I264MEDICINE
Fruit
I265
Peel
I266
Leaves
I267
FOOD
I268

Citrus Medica, Linn, Fl Br Ind, I, 514

I269

may be found to be the mountain tracts of Eastern Bengal, more particularly of the Khásia and Garo hills, while the latter is of a more northern character, extending along the foot of the Himalaya to the Panjab

C. I269

CITRUS
Medica.

The Citron.

The sweet lime (*C. Limetta*) appears to be the southern manifestation of

frontier.

This species includes as varieties the Citron, the Lemon, the Sweet and the Sour Lime.

Var. 1. *Medica* proper.

THE CITRON, CEDRAT-TREE, ADAM'S-APPLE, *Eng.*; CEDRATIER, CITRONIER, *Fr.*; CISTRATO, CEDRO, *It.*; CIDRO, *Sp.*; CIDREIR, *Port.*; CEDRATEN, CITRONENBAUM, *Germ.*

Considerable difference of opinion prevails as to the origin of the word Citron. It is presumed that the Median apple was synonymous with the

Syn.—*C. AURANTIUM*, var. *MEDICA*, *W. & A. Prodr.*; *C. MEDICA*, var. *A.*, *Linn.*; *CITRUS MEDICA*, *Risso*

Vern.—*Bijaura*, *limbu*, *kutla*, *bara nimbu*, *turanj*, *nimbu*, *limu*, *HIND.*;

much resembles a small pumelo.

C. 1270

Var. 1
Medica.
1270

The Citron; The Lemon.

CITRUS
Medica.

Captivity According to Galesio it was introduced into Italy about the third or fourth century. The Jews cultivated citron when under the Roman rule, and used the fruit, as at the present day, in the Feast of Tabernacles; each person bringing a citron in his hand. Dr Royle found the species growing wild in the forests of Northern India, and, as already stated, it may therefore fairly be conjectured that the original home of the citron was in India. It has now spread over the whole of the civilised world, and even in cold regions it is cultivated under artificial heat.

Gum—Said to yield scantily an unimportant gum. Sent from Masulipatam to the Madras Exhibition in 1855.

GUM
I271
OIL
I272

(Pesse)

Medica Citron rind, khatái, used for dysentery, as a sedative (*Yerr-Book, Pharm.*, 1874, 623)

Special Opinions—§ "The rind is made into a marmalade and is an antiscorbutic" (*Surgeon-Major A. S. G. Fayakar, Muskat*) "It is made into preserve and is used for dysentery" (*Surgeon-Major J. Robb, Ahmedabad*)

Food.—The fruit is described in the *Flora of British India* as large, oblong or obovoid; rind usually warted, thick, tender, aromatic; pulp scanty, sub-acid. The rind makes good comfit, the pulp is also preserved in sugar. Both fruit and preserve are somewhat bitter to the taste. The rind of the fruit candied is well known as a delicate sweatmeat. Atkinson says the wild fruit is used for pickling (*khatái*).

D-

n

l.

-

lp "

Var. 2 *Limonum, sp. Risso.*

The word lemon is from the Arabic *limun* and *limon* is the

MEDICINE.
Rind
I273
Pulp
I274
Seeds
I275
Leaves.
I276
Juice
I277
Marmalade.
I278
FOOD
Fruit
I279
Comfit.
I280
Candied Rind.

I283
TIMBER.
I284
DOMESTIC
I285
Var 2.
Limonum.
I286

birá nimbu or large nimbu.

The Citron; The Lemon.

CITRUS
Medica.

According to Gallesio it
third or
rule, and
nacles;
species

stated, it may therefore fairly be conjectured that the original home of
the citron was in India. It has now spread over the whole of the civilised

important gum. Sent from Ma-

GUM
1271
OIL
1272

(Pesse)

Medicine.

SPEDS, LEAVES,
gent Accordi

To one who has taken a poison injurious to life, it may be given, producing
a strong action, drawn out. It also corrects
for of the fruit is used as a

se
an antiscorbutic" (Surgeon-Major A. S. G. Jayakar, Muskat) "It
is made into preserve and is used for dysentery" (Surgeon-Major
Robb, Ahmedabad)

MEDICINE.
Rind
1273
Pulp.
1274
Seeds.
1275
Leaves.
1276
Juice
1277
Marmalade.
1278
FOOD.
Fruit
1279
Comdt.
1280

Food.—The fruit is described in the *Flora of British India* as large,
oblong or obovoid; rind usually warted, thick, tender, aromatic; pulp
scanty, sub-acid. The rind makes good comfit, the pulp is also pre-
served in sugar. Both fruit and preserve are somewhat bitter to the
taste. The rind of the fruit candied is well known as a delicate
sweet.

othe
they

Var. 2 *Limonium, sp. Risso.*

The word lemon is from the Arabic *limūn*, and this, through the
Persian, is the Hindi *limu*, *limbu*, or *nimbu*, probably adopted by the
Sanskrit people. Much stress is by authors laid on the

DOMESTIC
1285
Var. 2.
Limonium.
1286

The Lemon.

CITRUS
Medica.Method of
extraction.
1288

in France. A brief account of the methods of extraction, as given in the *Pharmacographia* (p. 119), may be reproduced here:—

Sponge process.—"The workman first cuts off the peel in three thick longitudinal slices, leaving a little peel at either end in the middle, throwing it on one side. The latter are allowed to dry thus: the workman seates a piece of sponge, wrapping it round his fore-finger. With the other he places on the sponge one of the slices of peel, the outer surface downwards, then presses the rest side (which is uppermost), so as to give it for the moment a convex instead of a concave form. The vesicles are thus ruptured, and the oil which issues from them is received in the sponge with which they are in contact. Four or five slices are thus pressed, and then gives to each slice of peel, which done 1 bit of peel has attached to it a small trives to avoid pressing the latter. The workman wrings it forcibly, receiving its contents in a coarse earthen

hold-
which

oil - - - - - about

into a tube about an inch in diameter and five inches in length, closed at its lower end. This vessel, which is called an *écuelle à piquer*, has, therefore, some resemblance to a shallow, dish-shaped funnel, the tube of which is closed below. The workman takes a lemon in the hand, and rubs it

su
rec
lat
frc
ar
pr
w

ec
sa

by which the portion of peel richest in essential oil is removed, on a coarse grater of tinned iron, grated 1 This
essence ourless

Dest

yellow
says the
more

a faint
Presse

CHEMISTRY.
1290

the foregoing boils at 176°C . Lastly, a small quantity of cymene and of a compound acetic ether, $\text{C}_8\text{H}_8\text{O}$ ($\text{C}_{16}\text{H}_{17}\text{O}$), would appear to occur also in oil of lemons. The crude oil of lemons already yields the crystalline compound $\text{C}_{10}\text{H}_{16} + \text{Z H Cl}$, when saturated with anhydrous

CITRUS
Medica.

The Lemon.

PERFUMERY
1291

hydrochloric acid, the sol
the sol
Prec
a flav
combination with rosemary, cloves, and caraway, for perfuming powders for the nursery. From its rapid oxidation it should not be used for perfuming green as it has a tendency to turn black.

MEDICINE
1292

Valued as an antiscorbutic and refrigerant—primarily antalkaline; secondarily, antacid. It forms the best remedy for scurvy, and an excellent drink in fever and inflammatory affections. It has met with success in acute rheumatism, dysentery, and diarrhoea. It also forms an antidote to acro-narcotic poisons. (*Pharm. Ind.*) Mr. Baden Powell says that it is considered by

In bilious
with port wine
Medica fruct.
the relief
such as p
hip joints, &c.,
Saragadhiara recommends the use of lemon juice with *yarakshara* and honey (*U. C. Dutt*).

The best substitute for lemon juice is a solution of about eight drachms of citric acid in sixteen ounces of water, with the addition of a few drops of lemon oil. Lemon juice may also be used in preparing effervescent diaphoretic and diuretic draughts. The relative proportions of lemon juice and citric acid with the alkaline carbonates, for the formation of effervescent draughts, are as follow:—

Lemon juice—	or Citric acid—	to 20 grains of
℥i drs iijss	grs xiv	Bicarbonate of Potash,
℥i drs vi	grs xxiv	Carbonate of Ammonia,
℥i drs iv	grs xvii	Bicarbonate of Soda.

The lemon juice, being liable to spontaneous decomposition, speedily becomes unfit for medical use. "One of the best methods of preserving the juice is to allow it to stand for a short time after expression, till a coagulable matter separates, then to filter, and introduce it into glass bottles, with a stratum of almond oil or other sweet oil on its surface. It will keep still better if the bottles containing the filtered juice be suffered, before being closed, to stand for fifteen minutes in a vessel of boiling water. Another mode is to add one-tenth of alcohol and to filter. The juice may also be preserved by concentrating it either by evaporation with a gentle heat, or by exposure to a freezing temperature, which congeals the watery portion, and leaves the juice much stronger than before." (*U S Dispens.*, 15th Ed., 849)

Dr. Charles Rice of New York states that the bark of the root has been used in the West Indies as a febrifuge and the seeds as a vermifuge.

The Lemon, The Sour Lime

CITRUS
Medica.

§ 'Lemons as well as other fruits of the same order, contain a principle—*hesperidene* By some chemists this substance is described as bitter and crystalline and by others as tasteless. *Chladstone* b. 3 f.

of orange pee
A glucoside
mana. (Sur

Citric Acid

It occurs in colourless crystals, is very soluble in water, less soluble in rectified spirit, and insoluble in pure ether. The chief use of citric acid in medicine is in the preparation of effervescing draughts and refrigerant drinks.

§
be s
and
(Sur
L

are given for the preparation of this substance "Take of fresh lemon peel two ounces, lemon juice, strained, one pint, refined sugar, two pounds and a quarter Heat the lemon juice to the boiling point, and having until with and should have the specific gravity 1.34"

Special Opinions—§ 'Lime juice—Most useful in dysentery with sloughing of the mucus membranes I have given 12 ounces a day in apparently hopeless cases with success" (From a Contributor) "Lemon oil mixed with glycerine is applied on the eruption of acne" (Surgeon R Gray, Lahore) Lemon juice and gunpowder used topically for scabies" (Surgeon Major E C Bensley, Rajshahye) "The fruit in the form of pickle is useful in hypertrophy of the spleen" (Surgeon J C Penny, Amritsar)

Food—The lemon juice is used largely in sherbets and cooling drinks The fruit is also pickled

Var 3 acids.

THE SOUR LIME OF INDIA

Syn—C ACIDA R A E T T E S C H A C A

Vern—*Lebu, nebu, limbu, nimba* 1 = = = 14 1 = 1 1 1

Limbu, nimbu, pati, nebu

nebu BENG, *Nimba*

limbu GUJ, *Limbu*, 1

mich-cham-pasham, eli

pandu, nemniapunda

janatam, naranna, jerd

Jambira, limbika, ni

Limun, limue-kamis

Thandaya, samya si, tambiya si BUEN Dehi, SINGH

References—Brandis, For Fl 52, Stewart, Pb Pl, 29 DC Origin, Cult Pl, 179 U C Dutt Mat Med Hind, 225, Ainslie, Mat Ind, 1, 193 Atkinson Him Dist 710, McCann Dyes and Tans Bengal, 159 New Off Guide to the Museum, 25, New Off Guide to the Bot Gardens and Arboretum, 64

MEDICINE

Citric acid
1293Syrup.
1294FOOD
1295
1296

CITRUS
Medica

The Sour Lime.

Habitat—Wild in the warm valleys of the outer Himálaya, from Garhwal and Sikkim to the Khásia and Garo hills, Chittagong, and probably also the mountain tracts of the Central Provinces and of the Western Peninsula and the Satpura mountains of Central India. It

according to native gardeners. There are many minor cultivated forms, differing chiefly in size. The fruits of all are more or less round, smooth, with a shining rind, green, or only tinged with yellow when ripe.

Dye—The leaves of this plant are used in tanning in Mánbhum. This seems to be doubtful; at most, the leaves can be used only as an

DYE
I297MEDICINE
I298

inferior to a superior, it is beautiful to behold, cooling and fragrant to the smell, the juice of it rubbed upon the head will soothe the ravings of frenzy, and the rind of it dried in the sun has the power, when laid

swelling caused by mosquito bites (*Brigade Surgeon J. H. Ingham, Monghyr*)

FOOD
I299

Food—The Sour Lime of India has "flowers small, fruit usually small, globose or ovoid." U. C. Dutt says "The fresh juice, squeezed

Pickles
I300

and salt is a popular and effectual remedy by excess in eating, or by indigestible food rubbed over a stone, or their rind

from other fruits of the so the addition of common salt are preserved in porcelain

of Jaunpur and Azamgarh. The value is the *patli nibu* or *patli*, the *kamarali nibu* or *patli*, the small sour lime, and the larger ones

Dr. Bonavia remarks that they are to be found everywhere, and even where no other Citrus occurs, some kind of lime is sure to be seen. Nevertheless, it is astonishing that so common a thing so useful a fruit, and a tree so easily raised from seed, is not to be found in the villages of the North-West Provinces. There is probably

C. I300

The Sweet Lime, The Sweet Lemon.

CITRUS
Medica.

not a *village* in the whole of India where the *highest nimbu* would not readily grow" "Although they are called limes, I believe them to be an

FOOD.

Var. 4. *Limetta*, W. & A., *Fl Br Ind*, I, p 515.

THE SWEET LIME OF INDIA

Syn—*C NOBILIS* Lour, as in *Kurz, For Fl Burm.*, I, 197, *Wight Ic t 158* *C LIMETTA* Risso It might be asked has the *C LIMETTA*, Risso, sweet or bitter fruits? if the latter, it might be viewed as a synonym of var *acida*.

Vern—*Mitha nebu*, *nembá*, *mitha amrit phal*, HIND, *Mitha nebu*, BENG *Mita nimbu*, PA, *Mitha limbu*, GUJ, BOMB *Elemitchum* TAM, *Nemba pandu gajanumma*, TEL, *Erumitchi narracum*, MALA, *Madhukarkatika*, SANS, *Thanbaya* BURM, *Dekhi*, SING

Reference—P. J. F. F. A. O. I. S. C. I. B. L. E. C.

I301

Habitat—Commonly cultivated in most parts of India and Burma. Most probably a native of Southern India, Wight says it is indigenous at Kolagberry in the Nilgiri hills

Botanic Diagnosis—Leaves with winged petioles, flowers small, white, fruit globose or ovoid, shortly mamillate, rind with concave vesicles

The limes approach much nearer to the true oranges than do any of the other forms of *C. Medica*. Indeed, it is difficult to say how far the published accounts of *C. Limetta* have become mixed up with *C. Bigaradia*, and the vernacular names given to both these forms, as

preceding variety.

Var. 5. *Lumia*, W & A., *Fl Br Ind*, I, 515

THE SWEET LEMON, Eng., LUMIE, Fr. & Germ.

Vern—See *C LIMETTA*

MEDICINE
I322
FOOD.
I303

I304

Botanical Diagnosis—Leaf petioles simply margined, flowers tinged with red, fruit bright yellow, ovoid oblong, with a long curved mamilla, rind with convex vesicles, pulp sweet.

C I304

CITRUS
Medica.

The Sour Lime.

Habitat—Wild in the warm valleys of the outer Hímálāya, from Garhwal and Sikkim to the Khásia and Garo hills, Chittagong, and probably also the mountain tracts of the Central Provinces and of the Western Peninsula and the Satpura mountains of Central India. It

is distinguished as the large lime, but this is the lime itself in minor cultivated forms, more or less round, smooth, with a shining rind, green, or only tinged with yellow when ripe.

Dye.—The leaves of this plant are used in tanning in Mánbhum. This seems to be doubtful; at most, the leaves can be used only as an adjunct to the tans, imparting an odour to the leather.

Medicine.—"Lime-juice is much used in medicine by the native prac-

DYE.
1297

MEDICINE.
1298

FOOD
1299

Monghyr).

Food—The Sour Lime of India has "flowers small, fruit usually small, globose or ovoid." U. C. Dutt says: "The fresh juice, squeezed

Pickle
1300

with lime and salt is a popular and effective remedy for indigestion, by excess in eating, or by indigestible food. The fruit is first rubbed over a stone, or their rind is scraped off, and they are then steeped in juice obtained from other fruits of the sort, and exposed to the sun for a few days with the addition of common salt. When crisp and of a brown colour, they are preserved in porcelain vessels or glass jars. This preparation is called 'pickled lime' (or 'pickled lemon') in the vernacular." Atkinson

in the Punjab and Azamgarh is the 'páti-nibu' or 'kamaráti-nibu'.

The small sour lime is the larger ones.

See also remarks

and

of

on

to

ly

The Sweet Lime, The Sweet Lemon.

CITRUS
Medica.

FOOD.

1301

Var. 4. Limetta, W. & A., *Fl Br Ind*, I, p 515.

THE SWEET LIME OF INDIA.

Syn—C NOBILIS Lour, as in *Kurs, For Fl Burm.*, I, 197, Wight
l.c., t 558 C LIMETTA, Risso It might be asked has the C LIMETTA,
 Risso sweet or bitter fruits? if the latter, it might be viewed as a synonym
 of var. *acida*

Vern—*Mitha nebu*, *nembā*, *mitha amrit phal*, HIND, *Mitha nebu*,
 BENG *Mitha nimbu* PB, *Mitha limbū* GUJ, BOMB, *Elemitchum*
 TAM; *Nemma pandu*, *gajanimma* TEL, *Erumitchi narracum*, MALA,
Madhukarkatika, SANS, *Thanbaya* BURM, *Dehi*, SING

Habitat—Commonly cultivated in most parts of India and Burma.
 Most probably a native of Southern India, Wight says it is indigenous
 at Kolagberry in the Nilgiri hills

Botanic Diagnosis—Leaves with winged petioles, flowers small,
 white, fruit globose or ovoid, shortly mamillate, rind with concave
 vesicles

The limes approach much nearer to the true oranges than do any of
 the other forms of C. Medica. Indeed, it is difficult to say how far
 the published accounts of C Limetta have become mixed up with
 C Bigaradia, and the vernacular names given to both these forms, as

y of

340

bilis

hers

ce

(Surgeon & Lenny, *Amritsar*)

Food—The fruit is both eaten fresh and after being preserved or
 cooked in various ways, but the juice is not so much valued as that of the
 preceding variety

MEDICINE

1302

FOOD.

1303

Var. 5. Lumia, W & A. *Fl Br Ind*, I, 515

THE SWEET LEMON, Eng.; LUMIE, Fr. & Germ.

Vern—See C LIMETTA

Habitat—This form is very little known in India, and occurs only occa-
 sionally in gardens. It is probable that, with the lemon, this is not an Indian
 form. Atkinson and many Indian writers use the terms "sweet lime"
 and "sweet lemon" as synonymous

Botanical Diagnosis—Leaf petioles simply margined, flowers tinged
 with red, fruit bright yellow, ovoid oblong, with a long curved mamilla,
 rind with convex vesicles, pulp sweet

1304

C 1304

CLAUSENA
indica.

The Mandarin or Maltese Orange.

OIL.
1305

Essential Oil.—Dr. Rice says that this oil is prepared at Squillace in Calabria by mechanical means.

1306

Citrus nobilis, Lour.

THE MANDARIN ORANGE, sometimes also called the MALTESE ORANGE

Syn.—*CITRUS CHINENSIS* and *C. MYRTIFOLIUS*

Vern.—Probably the same as for *C. LIMETTA*; it is the *kán* of China

Habitat.—Cultivated in China and Cochín-China, where it appears to

1307

has been greatly ext
the blood oranges of
gardens at the begir
tifully in Sicily and

Botanical Diagn
face, spherical but
yellow, pulp almost blood red with a peculiar flavour, both leaves and
fruit have the same odour.

ENCOURAGE-
MENT OF
CULTIVATION
IN INDIA.

1308

sour, and juicy lemon known in the Panjab as *guigui*; and that Bombay
should prepare to meet the Indian demand for its excellent pomelos. In
this way, with extended railway communication, free interchange
might be made with the various provinces and a more constant and uni-
form supply
stricting th
which they
thoroughly
grow up conversant with the best modes of dealing with it, not only with
regard to the cultivation and propagation, but also with the best modes of
packing and preserving the fruit for a long time."

CLAUSENA, Linn.; *Gen Pl*, I., 304

1309

Clausena indica, Oliv.; *Fl. Br. Ind*, I., 505, *Beddome*; *RUTACEÆ*.

Syn.—*PISTOSTYLIS INDICA*, *Dals*; *Dals. & Gibs*, *Bomb. Fl.*, 29; *BER-*
GERA NITIDA, *Tam*, *Flum. Ceylon Pl.*, 41.

Vern.—*Migong-karapichi-gass*, *SINO*

Reference.—*Lisboa*, *U Pl of Bomb*, 33.

C. 1309

Ergot of Rye.

CLAVICEPS
purpurea.

Habitat —A shrub or small tree, met with in the Western Peninsula from the Bombay Ghats to the Anamally Hills, and also in Ceylon
Structure of the Wood.—Close-grained and hard, adapted for the lathe.

Clausena pentaphylla, DC, *Fl Br Ind*, I, 503

Syn —*AMYRIS PENTAPHYLLA*, Roxb • *Fl Ind*, Ed C B C, 321

Vern —*Rattanjote, surymukha, teyrur*, HIND

Defacement — D J F E C L M T

TIMBER
1310
1311

CLAVICEPS.

Claviceps purpurea, Tulasne, FUNGI

THE ERGOT OF RYE, HORNED OR SPIKED RYE (*Secale Cornutum*), BUNT

Syn —*SCLEROTIUM CLAVUS* DC. *ERGOTÆTIA ABORTIFACIENS*, Quak, *OMIDEUM ABORTIFACIENS*, Berk & Br

References —*Pharm Ind*, 251, O Shaughnessy, *Beng Disp*, 631, 673, 76, Balfour *Agri Pests of India*, 61, 115, Flück & Hanb, *Pharmacog*, 740, Benl & Trim, *Med Pl*, IV, 303, U S *Dispens*, 15th Ed, 5567

Dr R Tytler (in the *C J Med DI* • *Tr* • *So* • *and V A* • *1888*) reports that barley in a disease very similar

MEDICINE
Leaves
1312

1313

produced within the paleæ of the common rye, *Secale cereale*, forms the officinal part "In medicinal doses ergot acts principally upon the mus-

MEDICINE.
1314

tids, from the uterus

"In overdoses ergot produces nausea, vomiting, colicky pains, headache, and sometimes delirium, stupor, and even death Taken for a

CLAY.	Ergot of Rye Clay
1315	<p>seem of good quality but which contain a fungus, most probably an ergot. It seems probable that Indian wheat rust may be due to a species of <i>Æcidium</i> reared on a <i>Euphorbia</i>.</p>
1316	<p>Some writers have attributed to an ergot the poisonous qualities which <i>kesari</i> (<i>Lathyrus sativus</i>) is said to possess. An indulgent use of this produces a paralysis of the lower limbs which is generally incurable. See under Fungoid Pests.</p>
	<p style="text-align: center;">CLAY.</p>
1317	<p>Clay is a hydrated silicate of alumina, which is expressed in mineralogy by the formula $H_2Si_2O_5 + H_2O$ which may be said to be Si 52, 46.40, Al, O 39.68, Water 13.92.</p>
	<p>Vernacular and other names —</p>
	<p>Properties and Classification — The pure clay, defined above, when it occurs, is fine, however, clay, shale, or these would, or less clay on, the peculiar superficial deposits in river-basins, estuaries, or dried up lakes. Pure clay is derived from a decomposition of felspar, from which the silicates of potash, soda, &c., have been washed out. The purer forms of clay are</p>
	<p>the former makes red clays, and the latter dark or even almost black. The former is usually much of silicate, and the latter is a form imparted termed "clay."</p>
	<p>These facts naturally lead to an industrial classification of the clays, and in dealing with those met with in India we shall, as far as possible, take them up in the alphabetical order of their better known names in preference to attempting a scientific assortment.</p>
	<p style="text-align: center;">I.—BRICK CLAYS</p>
1318	<p>In the early part of the present century, it was thought necessary to import bricks into India from England. It was soon discovered, however, that in almost every district clays suitable for this purpose existed.</p>
C. 1318	

Brick-Clay.

CLAY.

abundance, for bricks were employed in many buildings in India long anterior to the arrival of the English. Some of an enormous size are found in the ancient monuments, and in more recent times others much

is to blame. Of course there are some clays so impregnated with lime as to be valuable to the manufacture of the large rivers from these impurities at Akra near Calcutta, which is exported annually. India see the Rurki

II—EDIBLE AND MEDICINAL CLAYS AND FULLER'S EARTH.

1319

In most bazars in India a fine unctuous or oily clay is sold as a drug or as an article of food eaten by *enceinte* women, or used by ladies as a cosmetic. Allied to this is the clay used to effect caste markings on the forehead. Balfour says such a clay "is excavated from a pit near Koluth in large quantities, and exported as an article of commerce,

Manipur, which he was informed was regularly eaten by the women

Multani.
1320

further comments on an imported earth known as *Sang-i-Basri* (a Persian name) "This is generally imported from Bassorah and the Persian Gulf, as its name implies. It is used in tonic preparations and in irregular menses and with benefit from the iron it contains." He states that the earth in question is a silicate of alumina with lime and iron. U. C. Dutt (*Sans. Mat. Med.*) after dealing with red and yellow ochre (which see) or the *geru māl* in Beng., and *gairika* in Sans., adds: "besides *gairika* several other varieties of earth are described

for relieving bleeding from internal organs. If this earth be a natural product of Surat it is nowhere (so far as the writer can discover) de-

CLAY.

Edible Clay.

the source of a product may be inferred from its name. Under his

1321

1322

1323

which bore the name of *gagní* or *garí*; the shop-keeper could, however,

as indicat-

il or quasi-

are most

oned earth

' Fuller's

570) gives

pposition

r's earth.

His account is of so much interest that we may reproduce here the main facts from it. "Being of detrital origin fuller's earth does not possess

Sabun Miti.

1324

1325

1326

ong in the Bhagalpur
al mitti, a comestible

earth, the precise source of which is not known.

Ajmir

itions that fuller's

Over 2,000 camel-

Fire Clay

CLAY.

Bombay and Sind—A pale greenish clay is found in Western Sind, which is used for washing, and is also eaten by pregnant women.

Panjab—Dera Ghazi Khan and Multan already alluded to, in the Salt range at Nilawan, Mr. Wynne says a lavender-coloured clay is found which is used as a fuller's earth.

III.—FIRE CLAYS.

These derive their name from their refractory nature—that is to say, they are capable of resisting great heat without fusing.

lays

, or

In

ider-

and

that

of 11

I329

clays are procurable at Streepermatoor, Tipasool, Chingieput, Mitnapo-
liam, and Cuddapah, indeed, are very common in many parts of India, and
bricks can be made that resist the action of great heat. A clay found at
Bey pore, 20 to 30 feet below the surface, is used for fire-bricks and for

I331

as follow —

"(1) First experiment in September 1874 by Theodore W. H. Hughes, Esq., F.G.S., A.R.S.M., Officiating Deputy Superintendent, Geological Survey, India.

"The fire-bricks tested by me were furnished by the firm of Messrs Burn and Company. The materials from which they are made are very refractory and capable of resisting high temperature, without sensibly fusing. That, compared with Stourbridge fire-bricks, they are somewhat superior.

C. I331

CLAY.

Pipe Clay.

Whitelaw, Manager of the Bengal Iron Company's proposed works and others, who agreed in the favourable estimate formed of the quality of these bricks.

"In addition to the foregoing we beg to quote you the opinions of D. W. Campbell, Esq., Locomotive Superintendent, East Indian Railway, and J. Blackburn, Esq., Engineer and Manager of the Oriental Gas Company. The former, in a letter to us, dated 23rd February 1875, writes:—

"(2) I have had the fire-bricks and fire-clay tried here, they are both very good; I will send you a requisition as soon as present stock is exhausted."

"And Mr. Blackburn, in his letter of 2nd March 1875, states as follows —

"(3) The Gas retorts made for the Company by your firm two years ago have since been kept in constant use at a temperature of about 2,000° Fht., and they have been found fully as durable and effective as those of the best English manufacture."

"We trust that the above extracts will be found to contain the information required by Dr. Watt for the Dictionary of Economic Products, but in case he wishes to analyse the clay himself, we have pleasure in sending herewith a few sample pieces obtained from the coal measures of the Raniganj District."

IV.—PIPE CLAYS.

This is known as *Namam* in Tamil and *Kharra* in Dukhni; its English name is taken from the fact of its being used to manufacture tobacco-pipes. It much resembles China-clay, only that it possesses more silica. Balfour says "This is found in abundance in several parts of India, the Hindus employ it for making the distinguishing marks on their foreheads, and (moistened with water) it is often applied

between Terany and Kauray in Trichinopoly.

V.—POTTERY CLAYS.

These might be popularly referred to three sections or degrees of purity: (a) porcelain or kaolin clays, (b) ordinary white or glazed pottery clays, and (c) red or tile and flower pot clays. In every province, indeed in almost every district of India, one or other of these

Pottery Clay.

CLAY.

Bengal, is attempting to compete with European imported articles
 under the name of the School of Art + as off + he has made to + the

glazed pottery is less known than is the case in many parts of India
 Mr. Kipling (*Journal of Indian Art*) says: "No substance resembling
 the fine clays of Dorsetshire, Devonshire, and Cornwall, is known to the

social status, no craft, excepting, perhaps, that of the leather-dresser, is
 held in lower esteem than the potter's trade in Hindustan, the Deccan, and
 South India." Mr. Kipling next distinguishes the two classes of
 workers in earth, viz., *Kumhars* and *Kashigars*. The former are the
 common village potters who "produce wares which, though of little techni-
 cal value as pottery and of small commercial importance, are often good in
 colour and form, and perfectly fitted for the purposes they are intended to
 serve." The latter, the *Kashigars*, are "makers of glazed earthenware
 who are only to be found in the Panjáb and in Sind, and within the last
 few years in the town of Bombay and at Khurja in the North-Western
 Provinces. The name of the trade is Persian, derived probably from

into India by the Mussulman invasion, and not by means of the friendly
 intercourse which there seems reason to believe subsisted at various times
 with Tibet and the further East." Sir George Birdwood (*Indian Arts*)

CLAY.	Pottery Clay.
1335	<p>par and kaolin are obtainable in different parts of the district." "In the South Arcot district a fine plastic clay occurs in the Cuddalore beds near the south bank of the Guddalum," but it contains small quantities of lime and iron, the latter giving it a pinkish tint. In North Arcot the granite beds of the district are decomposed to a certain extent, and, according to the report of the Geological Survey, furnish a very considerable supply of pottery clays which may enjoy some reputation, but they are not so abundant as the pottery clays exist in great abundance in the district of Chingleput, more especially at Sririperimattur. From the beds exposed at Coopum a supply has been taken for the Madras School of Art.</p> <p><i>2nd, Mysore.</i>—For many years it has been known that kaolin earth occurs in the hills near Bangalore. It has been sent from</p>
1336	<p><i>3rd, Mangalore.</i>—As early as 1811 Dr. Christie discovered, in association with the laterite, an extensive deposit of what he conceived to be pure porcelain clay</p>
1337	<p><i>4th, Bengal.</i>—In Orissa white clays occur in the Mahanadi valley of Rajmahal age. These clays are used by the natives for ornamenting</p>
1338	<p>suitable for the manufacture of many articles of hard pottery, and which, with proper treatment, would afford suitable material for fire-bricks. But the best known clays of this series are the refractory and other clays now being worked by Messrs. Burn and Co. of Raniganj. The</p>
1339	<p></p>
1340	<p><i>7th, Assam and Burma.</i>—Rich deposits of porcelain clays have been reported to occur in Upper Assam near the Bhramakhund, known locally as <i>rukmanipitha</i>, and a fine clay for pottery purposes is also said to be found near the base of the cretaceous rocks at the western end of the Garo hills. In Burma the ordinary alluvial clay, mixed with sand, affords the material for common pottery, but a dark-coloured seam in the Irawadi valley is much sought after by the potters. Some of the upper beds in the nummulitic group are said to consist of China clay and would answer</p>

Glazing and Colouring Pottery.

CLAY.

well for pottery, owing to their freedom from iron. Kaolin is also reported to exist in Tenasserim. Of the clays experimented with by Sir William O'Shaughnessy that from Singapore was said to be the best.

VI.—MATERIALS USED FOR GLAZING OR PAINTING POTTERY IN INDIA.

I341

The indigenous art of glazing pottery, as practised in India is crude and unsatisfactory. Ball says "The varnish or imperfect glaze used for the sugar-boilers' pans, known in Bengal as *kolas*, is thus described by Mr Piddington. There are two kinds of earth used, one of which is called *belutti*, it is a silicious and ochreous earth, the best being found at

use, the p
Uporom,
20 miles
Kulna

obtained from one maund of the earth, two varieties of the *uporom* are

of lime. The black colour of pottery is often obtained from the smoke of oil-cake thrown into the kiln when the baking is complete. At other times an organic varnish is used for this purpose, except when, as mentioned in connection with Azimgarh, the clay itself contains the necessary organic matter to cause it to burn black. Artificially blackened pottery is produced at Monghir, Patna, Sarun, Chunar, and Surat. In the younger rocks of the Rajmahal series certain clays occur called *khari*. These are used as pigments. According to Buchanan the potters of Rajmahal use this *khari* for giving a white surface to pottery made of ordinary clays. Cheap pottery is often painted after having been baked, such as that seen at Kota, Lucknow, Benares, &c., at other times it is powdered with mica, or by other mechanical means has a colour imparted to it. Black pottery is, for example, often etched, and a preparation of tin and mercury rubbed into the patterns in imitation of metal-brown ware. With the exception of these miserable attempts the *kumhar*

I342

I343

the material is put into a furnace until it melts, when clean-picked *shora*

C. I343

CLEIDION
javanicum.

Glazing and Colouring Pottery,

1344

kalmi or saltpetre is stirred in. A foam appears in the surface, which is skimmed off and set aside for use. The latter is similarly made of quartzose rock and borax or siliceous sand and soda. "A point is made of firing the furnace in which the *kanch* is melted with *kikar*" (*Acacia arabica*)
 oxides
 of the *l*
sikka s
 made b
 reducin
 reducing with zinc instead of tin, *sikka lal* in the same way, oxidising

been roasted and powdered, mixed with a little powdered flint. Sir
 of the deep blue another piece of *nila* or indigo
 1 flint 4 parts
 5, zinc 5, and

1345

The yellow glaze used as the basis of the greens is made of *sikka sard*, white oxide 1 seer, and *sang safed* a white quartzose rock or mill stone, or burnt and powdered flint, 4 chittaks, to which, when fused, 1 chittak of borax is added.

1346

"The green colours produced are (1) *Zamrud*, deep green (1 seer of glaze and 3 chittaks of *chihul tamba* or calcined copper), (2) *Sibz*, full by smaller
 urning 1 seer
 George Bird-

wood, in his most interesting account of Indian pottery, after having described the glazes and colours used proceeds "The colours after being reduced to powder, are painted on with gum or gluten. The vessel to receive them is first carefully smoothed over and cleaned, and, as the pottery clay is red when burnt it is next painted all over with a soapy, whitish engobe, prepared with white clay and borax and *Acacia* or *Anogeissus* gums called *kharya mutti*. The powdered colours are ground up with a mixture of *nishasta*, or gluten and water called *mawa*, until the proper consistence is obtained when they are painted on with a brush. The vessels are then carefully dried and baked in a furnace heated with *ber* (*Zizyphus*), or, in some cases, *Capparis* wood."

1347

VII—CLAYS OR EARTHS EMPLOYED AS PIGMENTS
OR DYES

See "Pigments" for further information as to colouring of pottery

Clearing Nut, see *Strychnos potatorum*, Linn., LOGANIACEÆ

CLEIDION, *Blume, Gen Pl, III, 320*

1348

Cleidion javanicum, Bl., Fl Br Ind, V, 444, EUPHORBIACEÆ

Syn.—*ROTTLEIRA URANDA* Dals & Gids Bomb Fl, 230

Vern.—*Okaragass okuru*, Sing

References.—*Kurs For Fl, Burm, II, 300, Beddome, Fl Syl, t cclxxii, Gamble Man Timb, 348, Thwaites, En Ceylon Fl, 271, Lisboa, U Pl Bomb, 123*

C. 1348

The Clematis.		CLEMATIS grata
Habitat — An evergreen tree met with in the tropical forests of North- rather heavy, durable in		TIMBER. 1349
[EUPHORBIACEÆ.		
CLEISTANTHUS, Hook f. Gen Pl, III, 268,		
Cleistanthus malabaricus, Mull-Arg, Fl Br Ind, V, 276		1350
References — Gamble, Man Timb, 357 Lisboa, U Pl Bomb, 120		
Habitat — A small tree found in the Konkan and Malabar districts of South India		
Structure of the Wood — Lisboa mentions this plant amongst his useful timbers		TIMBER 1351
C. myrianthus, Kurz, For Fl Burm, II 370, Fl Br Ind, V, 275		1352
Vern — Mo man tha BURM		
Reference — Gamble Man Timb, 357		
Habitat — A moderate sized evergreen tree of the tropical forests of Burma and the Andaman Islands		
Structure of the Wood — Moderately hard, reddish grey Weight 41lb per cubic foot		TIMBER. 1353
CLEMATIS, Linn, Gen Pl, I, 3		
Clematis barbellata, Edgew, Fl Br Ind, I, 3, RANUNCULACEÆ		1354
Reference — Gamble, Man Timb, I		
Habitat — A woody climber of the western temperate Himalaya, Garhwal, and Kumaon		
C. Buchananiana, DC, Fl Br Ind, I, 6		1355
References — Kurz, For Fl Burm, I, 17 Gamble, Man Timb, I, Royle Ill Him Bot, I, 51		
Habitat — A large woody climber, occurs throughout the temperate Himalaya at 6,000 feet		
C. Gouriana, Roxb, Fl Br Ind, I, 4, Wight, Ic, t 933 4		1356
References — Roxb, Fl Br Ind, I, 4 Wight, Ic, t 933 4		
Balfour, Cyclop		
Habitat — An extensive climber found in the hilly districts from the Western Himalaya, rising up to 3000 feet, to Ceylon and the Western Peninsula		
		MEDICINE. Leaves. 1357 Stems 1358
C. grata, Wall, Fl Br Ind, I, 3		1359
Vern — Ghantali, bilari, HIND		
References — Gamble, Man Timb, I, Voigt, Hort Sub Cal, 2, Royle, Ill Him Bot, I, 44, 45, 51, Balfour, Cyclop		

CLEOME
VISCOSA

Wild Mustard

Habitat—A climber of the sub-tropical and temperate Himalaya at 2,000 to 3,000 feet

1360 *Clematis montana*, Ham ; Fl Br Ind , I , 2

Vern—*Ghantidli*, HIND

References—*Gamble, Man Timb, I Royle, Ill Him Bot I, 45, 51*

Habitat—A woody climber of the temperate Himalaya, from the Indus to the Bramaputra ascending to 12,000 feet, always above 8,500 in Sikkim, and in the Khasa Hills, Manipur, above 4,000 feet

1361 *C. napaulensis*, DC ; Fl Br Ind , I , 2

Vern—*Pawanne birri, wandok*, PB

References—*Stewart, Pb Pl 3 Royle Ill Him Bot, 23*

Habitat—Found in the temperate Himalaya from Garhwal to Bhutan

Medicine—In Kanawar the LEAVES are said to act deleteriously on the skin

MEDICINE
Leaves

1362

1363 *C. triloba*, Heyne, Fl Br Ind I, 3

Vern—*Moravele, moriel, moriel, ranjee, ranjai*, BOMB, *Moravele*, MAR

References—*Dals & Gbs Bomb Fl, I Dymock, Mat Med W Ind, 2nd Ed 21, S Arjun Bomb Drugs 2*

Habitat—An extensive climber met with in the mountains of the

MEDICINE
Plant

1364

FIBRE

1365

Distillate

1366

CLEOME, Linn , Gen Pl I, 105, 968

Cleome pentaphylla, see *Gynandropsis pentaphylla*, DC , CAPPARIDEE

1367 *C. viscosa*, Linn , Fl Br Ind , I, 170 Wright, Ic, t 2

Sometimes called WILD MUSTARD

Syn—*C. ICOSANDRA* Linn *POLANISIA VISCOSA*, DC , *P. ICOSANDRA*, W & A

Vern—*K* *lia* HIND, *Tinnaru*, *Ka fod*, *jangli kul*, *AM, Kuka*, *ALA, Naya*, *ina-b urbdra*

References—*Royle Fl Ind Ed C B C 501 U C Dutt Mat Med Hind 289 Dymock Mat Med W Ind 2nd Ed 61 Ainslie Mat Ind II 223 O Shaughnessy Beng Dispens 206 Murray Pl and Drugs Sind 52 Drury, U Pl 351 Baden Powell Pb Prod 330, Cooke Ols and Oilseeds 37 Atkinson, Him Dist 732, B rdwood*

C. 1367

or Hurhur

CLEOME
VISCOSA.

Bomb Pr, 276 Lisboa, U Pl Bomb, 145, Spens Encyclop, 1415
Balfour, Cyclop

Habitat—A common weed throughout the greater part of India, appearing in the rainy season, very common in Bengal and South India.

Oil—The seeds yield a light olive-green coloured limpid oil when subject to a great pressure. It seems likely that this oil would prove serviceable where a very liquid oil is required. The oil could be prepared to any extent.

Medicine—The JUICE of the leaves is poured into the ear to relieve ear-ache. According to Rheede, it is useful in deafness. Dr Dymock writes that the juice mixed with oil is a popular remedy in Bombay for purulent discharges from the ear, whence the Bombay name of the plant *Kanphutti*. "The LEAVES boiled in *ghis* are applied to recent wounds, and the juice to ulcers" (*Drury*). In Cochin China the whole plant, bruised,

OIL
1368MEDICINE
Juice
1369Leaves
1370Seeds
1371

to relieve ear-ache and as an astringent in cases of atorrhœa the ear should be syringed well before its application" (*Brigade Surgeon J H Thornton, Monghyr*). "Alterative, useful in secondary syphilis and enlargement of the liver and spleen" (*Surgeon-Major J McD Houston Travancore, and John Gomes, Esq, Medical Storekeeper, Trevandrum*). "The seed made into chutney has strong digestive power" (*Native Doctor Ummegudien, Mettapolissam, Madras*).

"The seeds of *Cleome viscosa* are anthelmintic, rubefacient, and vesicant, and the leaves rubefacient, vesicant, and a useful remedy for a few diseases of the ear. The seeds are valuable in expelling round worms, and also as a rubefacient and vesicant in all the complaints in which mustard is used. The leaves are also useful in the same way as a local stimulant, and, in addition to this, the juice possesses a curative influence over some cases of otalgia and otorrhœa, but the smarting it produces in

according to their age. As a drug the leaves of *Cleome viscosa* are much superior to those of *Gynandropsis pentaphylla*. It is the former which possess a distinct fetid smell and efficient rubefacient and vesicant properties, and not the latter. The above plants are frequently found growing together and are often confused partly from a general botanical similarity between them, and partly on account of their native synonyms being almost the same. The close similarity of their seeds adds greatly to this confusion. There will be, however, no difficulty in

A Mild Antiperiodic

distinguishing the two plants if due attention is paid to the following botanical characters -

Cleome viscosa — Siliqua flat, striated, pubescent, and sessile or short stalked, flowers yellow, stem and branches quite covered with viscid

strongly

"As the seeds of both of these plants are very similar, I need not describe them separately. They are as follows: small, flat, and slightly acid or bitterish in taste. They yield a small quantity of fixed oil on expression.

remedial value' (Honorary Surgeon Moodten Sheriff, Khan Bahadur, Triplicane Madras)

Food — The SEEDS of *Cleome viscosa* are much used by the natives, chiefly the Brahmans, in their curries, they are sold in all the bazars at a trifling price (*Roxb*). Lisboa says that the PLANT is eaten boiled with chillies and salt as salad.

CLERODENDRON, *Linn* ; *Gen Pl*, II, 1155

This name alludes to the variable properties of the species *kleros*, lot, and *dendron*, a tree

†VERBENACEÆ

1374

Clerodendron Colebrookianum, Walp , Fl Br Ind, IV, 594,

Vern — Ladungb; ЛЕРЧА

Reference.—Gamble, *Man Timb* 299

Habitat—An evergreen shrub with silvery grey bark, met with in Sikkim and the Khasia Hills, 2 000 to 6 000 feet also in Burma.

Food—The young LEAVES are eaten by the Lepchas.

Structure of the Wood — Grey, soft

C. inerme, Gærtm Fl Br Ind, IV, 586

SVH —VOLKAMERIA INERMIS Linn

Vern —Sang-kupbi: sang k bi: la i jai, HIND Bun jumat, bun joi: don

11 MAR 3

THE *NEW YORK*

*154 mg/dl

La prima

Pirola[illegible]

Habitat — A large, ramous often scandent evergreen shrub, common in tidal forests in Bengal, Burma, and the Andamans.

Perfumery—An exquisite perfume is said to be derived from the flowers of this plant (*Presse*)

Medicine — Dr Dymock says that the PLANT has a reputation as a febrifuge in remittent and intermittent fevers. This fact is supported by Dr Sakharam Arjun, who, upon the authority of Dr Hojel, states that

C. 1379

PERFUMERY

1378

MEDICINE

Plant

1379

A Substitute for Chiretta

CLERODENDRON
infortunatum

"the thick succulent leaves are very bitter, and on expression yield a large quantity of thickish somewhat mucilaginous juice with a slightly saline but intensely bitter taste. Although not generally known, it has of late been used as a febrifuge and antiperiodic with marked benefit."

[Wight, Ic, t 1471

Clerodendron infortunatum, Garl'n, Fl Br Ind, IV, 594.

1380

Syn — VOLLAERIA INFORTUNATA, Roxb, Fl Ind, Ed C B C, 478, G VISCOSUM, Vent

Vern — Bhant bhat HIND, Bhant, glentu BENG, Kharbari, barni or varni SANTAL, Kula narsal KOL, Chitu NEPAL, Adung, LECHA, Likunah, MECHI, Kali basiti PB, Kari BOMI, Bhandira, kari, MAR, Bockada TEL, Peragu MALA, Bhandira, bhanti bhitaka, SANS, Ka aunggyi, buiphyi, khaoung gyi BURM, Gas pinna, SING

References — Brandis For Fl 363, Kura For Fl Burm, II, 267,

74, S Arjun
Pb Pr, 364,
Mal, II, t 25

Habitat.—A pinkish-white-flowered shrub, common in waste places throughout the greater part of India and Burma and in the damp forests of Ceylon up to an elevation of 5000 feet. Grows gregariously, forming a dense under-vegetation, specially associated with the Bamboo. On passing into fruit the calyx becomes scarlet, and the plant is then even more attractive than when covered with its foetidly-scented flowers.

Medicine.—Dr Bholanath Bose calls attention to the LEAVES of this plant as a cheap and efficient substitute for chiretta as a tonic and antiperiodic" (Pharm Ind). According to Dr Kanny Lal De, OIE, the fresh JUICE of the leaves is employed by the natives as a vermifuge, and also as a bitter tonic and febrifuge in malarious fevers, especially in those of children. Dr Dymock states that he has not seen the leaves used medicinally in Bombay, but they are bitter. Dr Hong Berger mentions the use of the BARK in medicine by the Arabian and the Indian physicians.

Special Opinions.—§ "The expressed juice is an excellent laxative, cholagogue and anthelmintic. It is used as an injection into the rectum in cases of ascarides. It is also a valuable bitter tonic, and the natives believe that its presence cures scabies in the locality" (Brigade Surgeon J H Thornton B A, M B, Monghir). "Is said to be a very useful antiperiodic" (Surgeon Major E Sanders Chittagong). "The juice of

MEDICINE.
Leaves,
1381

Juice
1382
Bark
1383

"Decoction of the leaves is used as an antiperiodic" (Surgeon Anund). "The leaves are used as a" (Officer W). " (Surgeon-

Decoction,
1384

Domestic Uses.—Edgeworth mentions that this plant is used in the Ambala district to give fire by friction.

DOMESTIC.
1385

C. 1385

CLERODENDRON
serratum

Medicine for Cattle

1386

Clerodendron phlomoides, Linn Fl Br Ind IV 590 Wight Ic t 1473
Vern.—Hm A H n D C C C

References—Roxb Fl Ind Ed C B C 477 Brand s For Fl 363
Gamble Ma Tmb 298 Thwa tes E Ceylon Pl 243 Dals &
G bs Bomb Fl 200, At h son Cat Pb Pl 120 Vo gt Ho t Sub
Cal 465 Dymock Mat Med W Ind 498 Ansl e Mat I d II
408 Murray Pl and Drugs Snd 174 S Arj n Bomb Drugs 04
Royle Ill Hin Bota y 299 Balfour Cyclop

Hab tat—A tall pubescent shrub common in many parts of Ind a
pr nc pally n the dr er regions of the Panjáb Sindh Mar wara the Dekkan
Behar Bengal Oudh Central Prov nces and also in Ceylon

MEDICINE
Root
1387

Medicine—Dr Dymock says that the nat ves of Western Ind a sup
pose the ROOT of the plant has lterat ve propert es but he has never
seen it used as such
valescence o mersles
to Ansl e considered k

plant to the r cattle
to cu e them of diarrhoea and worms or when the stomach s ells Mr
Campbell also says the Santals rub the plant over the r bod es in dropsy

1388

C serratum, Spreng Fl Br Ind IV 592 Wight Ic t 1472
V n "

SANTAL
bharang
addakk
ra g or
nāpālu
a (root)

References—Brand s For Fl 364 Kurs For Fl Burm II 267
Gamble Man Tmb 299 Dals & G bs Bomb Fl 200 At h son
Cat Pb Pl 12 Vo gt Ho t Sub Cat 466 Pharm Ind 164
Med W I d
le Cat Raw
ury U Pl
168 Balfour

Cyclop

Hab tat—A blue flowered shrub common in the Sub H malayan tract
e m h C d l C C

MEDICINE
Root
1389

h ck
re
ely
"

Leaves
1390

Seeds
1391

C. 1391

**CLITORIA
Ternatea.**

A Powerful Cathartic.

Indonésien, *gokarna mal*, *nilaghiré kurni*, *nila-g'hiria*, *khurne*, *SANS*, *Mdsariyune-hindi* (Indian Mezereon), *basrulmasari*, *yune-hindi* (seeds), *ARAB*, *Darakhte-bikhe-hayat*, *tukhme bikhehayat* (seeds), *PERS*, *Bakyu*, *poi noung ni*, *oung mai phya*, *BURM*, *Kattarodu*, *nil katta-rodu*, *nil-katarolu*, *SING*

References.—*Park*, *Fl. Ind. Es. C. C. C.*, *Fl. Ind. Es. C. C. C.*

Habitat.—A common garden flower, also occurs in every hedge-row all over India. The seeds were first taken to England from the Island of Ternate, one of the Moluccas, hence the specific (and former generic) name of the plant.

Dye—*Bidie* remarks that the SEEDS are said to be used by dyers. "The corollas of the blue variety are said to afford a blue dye in Cochin China, but it is not permanent; and *Rumphius* says that they are used for colouring boiled rice in Amboyna" (*Treasury of Botany*)

Medicine.—The ROOT is a powerful cathartic like jalap, and has been recommended to be used along with other laxatives and diuretics in ascites and enlargements of the abdominal viscera (*Dymock*). *Ainslie* recommends it in croup as an emetic, but *O'Shaughnessy*, in *Bengal Dispensa-*

DYE.
Seeds
I404

MEDICINE
Root.
I405

Seeds.
I406

Leaves
I407

as a diuretic, and in some cases as a laxative. The SEEDS are, however, more useful, and have gained a certain reputation in Europe as a safe medicine, especially for children. The powdered seeds are purgative and aperient. Combined with acid tartrate of potash and ginger, they are administered in the same doses as jalap. The infusion of the LEAVES is

used in cases of cholera, and in some cases of dysentery.

..

ing and to act as an antidote to poisons. The roots are used as emetics and in rheuma

orders, also in "The juice of in cases of colliquative sweating in hectic fever" (*Taylor, Med. Ind. Dacca*, 52, 53)

Special Opinions.—§ "There are two varieties of *Clitoria Ternatea* distinguished by the colour of their flowers, as blue and white, and the blue

C. I408

Jules
I408

Clitoria Seeds—a Medicine used in Croup, &c

CLOVES.

again has a sub variety, in which the flowers are double There is no distinct difference between the action of the seeds of these varieties, or if any at all, it is in favour of the white one The plants are in flower

MEDICINE.

thus precaution are nearly round or slightly compressed along the edges oblong, dull green, greenish brown, or brown in colour, and minutely mottled The ends of some seeds are round, and of others flat, as though cut off clean by a knife, taste d sagreeable and acrid, and no smell The thicker and rounder the seeds are, the more active they prove The immature seeds are flat and dark brown in colour, the matured thick and round seeds are an efficient purgative and produce five or six motions in one drachm or one drachm and a half doses Their action is increased in proportion to the increase of their quantity up to two drachms, when the

seeds are one of those but they may also be in equal proportion, he compound powder

The dose of the compound powder is from a drachm and a half to two drachms The following is a list of the seeds of Clitoria

dome
childr
It act
thus
doses
the sy
scaldi

gonorrhoeal discharge itself is much abated under its use One small root is generally a dose for children under two years, and one large root or two small ones for those between three and six years For adults

1409

The roots of the blue species are used as an antidote in cases of snake-bite" (Brigade Surgeon J H Thornton, B A, M B, Monghir) "The seeds are used as a mild purgative for children" (Surgeon Major J

white flowers and the st" (Nati e Doctor a drastic purgative geon Shib Chunder owdered root of this dropsy" (Surgeon

1410

Major John North, Bangalore)

Sacred Uses —The flower is held sacred to the goddess Durga

SACRED
USES.
1411

Clover, see *Trifolium pratense*, Linn , LEGUMINOSÆ

Cloves, see *Caryophyllus aromaticus*, Linn , MYRTACEÆ

C. 1411

COAL.	Coal
I412	<p>CNICUS, <i>Linn</i>, <i>Gen Pl</i>, II, 468</p> <p>Cnicus arvensis, <i>Hoffm</i>, <i>Fl Br Ind</i>, III., 362, COMPOSITÆ</p> <p>Syn — <i>CARDUUS LANATUS</i> <i>Roxb</i>, <i>Fl Ind</i>, Ed C B C, 595</p> <p>Vern — <i>Bhur bhur</i>, N W P</p> <p>Reference — <i>Smith, Dictionary</i>, 410</p>
OIL Seeds I413	<p>Habita — <i>W A L A N I A</i></p> <p>Gangetic</p> <p>Oil —</p> <p>The seeds them for their own use It burns with smoke, is otherwise of good quality</p>
Cnidium diffusum , see <i>Seseli indicum</i> , IV & A, UMBELLIFERÆ	
I414	<p>COAL.</p> <p>Coal</p> <p>CHARBON DE TERRE, <i>Fr</i>, STEINKOHLN, <i>Germ</i>, CARBONI FOS SILI, <i>It</i>, CARVES DE PEDRA, <i>Port</i>, CARBONES DE PIEDRA, <i>Sp</i></p> <p>Vern — <i>Kôyelah</i> or <i>kuela</i> HIND, <i>Kôyala</i>, BENG <i>Kôlsa</i> DUK <i>Kari</i> or <i>Simai karri</i>, TAM, <i>Boggu</i> or <i>Sima boggu</i> TEL; <i>Kari</i>, MAL, <i>Iddallu</i> KAN, <i>Koelo</i>, <i>koiso</i> GUJ, <i>Aiguru</i> CING <i>Fahm</i>, ARAB, <i>Zughal</i>, PERS, <i>Angaraha</i> SANS, <i>Misu</i>, <i>midu ye</i> BURM</p> <p>References — So much has been written regarding Indian Coal that an enumeration of the publications would occupy many pages. The reader is referred to <i>Ball's Economic Geology</i>, pp 599 604, to the <i>Memoirs</i> <i>Records of the Geological Survey</i>, and to the <i>Journals of the Asiatic</i> <i>Society of Bengal</i>. The following works may, however, be specially men- tioned —</p> <p><i>Final Report of the Coal Committee</i> Dr T. Oldham's <i>Report on the</i> <i>Coal Resources of India</i> <i>Sel Rec Govt Ind</i> LXIV</p> <p><i>Ball's Coal fields and Coal productions of India</i> <i>Annual Adminis-</i> <i>tration Reports on Railways in India</i></p>
REGIONS OF INDIAN COAL	
The following account of the coal fields of India has been furnished by Mr H B Medlicott for this publication —	
I415	<p>ABSTRACT OF THE FEATURES OF INDIAN COAL</p> <p>"India possesses extensive stores of coal, though none of it belongs to the so-styled carboniferous period, and in India itself the coal measure rocks are not all of one formation. All the coal of peninsular India occurs in the rocks known as the Gondwana system, the fossil flora of which has a mesozoic facies, and all the coal of extra peninsular India occurs in rocks of cretaceous or tertiary age. In both cases the distribution is partial — and north inces an North-W Madras f margin of the Indo-Gangetic plains from Sind to Pegu but it is only in Assam and Upper Burma that valuable measures have been found where a cretaceous coal occurs in workable quantity</p>

Coal fields of India

(H B Medlicott)

COAL.

' In both regions the quality of the coal varies much as in all coal-ndard, almost if not Gondwanan (Bengal) some an excess of ntrige of ash is low, producing a lighter

fuel The following tabular statement exhibits these facts —

	BENGAL		ASSAM	
	Average of 31	Best	Average of 23	Best
Fixed carbon	53 20	66 52	56 5	66 1
Volatile exclusive of moisture	25 93	28 12	34 6	33 5
Moisture	4 80	9 6	5 0	
Ash	16 17	4 40	3 9	4
	100	100	100	100

In Bengal only the Raniganj and Karharbari fields have as yet been largely worked and to a small extent the Daltonganj field. Several other large coal-fields are still quite untouched, owing to difficulty of communication.

"In the Central Provinces the Mohpani mines in the Narbada valley, and the Warora mines in the Wardha valley, have been for some time in work and the* Umaria and Sohagpur fields in the Rewah State are being opened up.

"In the* Singareni and Sastri fields of the Nizam's Territories some preliminary mining has been carried out pending the establishment of railway communication.

"Vigorous mining enterprise has recently been started in the Makum coal field in Upper Assam."

MORE DETAILED STATEMENT OF THE COAL-YIELDING DISTRICTS

' The mineral is more particularly developed in the central eastern portion

SOUTH INDIA
1416

the field about 38 miles
tent, and contains four
is the most southern

36', Long 81°7' Has its
the River Godavari, on
ons of coal, of which only

seams, neither of which ex-
e Godavari, and another,

"Singareni —The best field as yet known for Madras, but still in the Nizam's Dominions, is that near Singareni, lat 17°30'30", long 80°20'. There are five seams the thickness of one was not ascertained, those of the

* Since opened out

COAL	Coal-fields of India.
	<p>others are respectively 6, 3, 3, and 34 feet. This coal answers well for was found to be a serviceable way communication is now being started, coal report-</p> <p>Kamuram—Lat. 18° 5', Long 80° 14'. Two seams of fair coal, 9 and 6 feet in thickness respectively. The available coal is estimated at 1,132,560 tons, its position is, however, unfavourable to its development.</p> <p>Tandur—Lat 19° 9', Long 79° 30'. This village is situated about the centre of a strip of Barakar rocks, extending from Kaigura to Aksa pati, and contains a 15-foot seam of fair coal.</p> <p>Antergaon—Lat. 19° 32' 30", Long 79° 33'. South of this place a 6-foot seam occurs, 9 inches of which are shale.</p> <p>Sasti and Paoni—In the Nizam's Dominions, included in the Wardha area, a 50-foot seam occurs here, a considerable portion of which is of good quality. 30,000,000 tons of coal are estimated to be available from this source.</p>
ORISSA 1417	<p>valley of the Brah- coal is of an inferior</p>
BENGAL. 1418	<p>in the western margin of the Rajmahal hills, coal measure rocks are exposed, and these doubt- less extend over a vastly greater area under the younger formations. Separated by these overlying rocks, there are five distinct fields, namely, Hura, Chaparbhitia, Pachwara, Mohowgurhi, and Brahmini. There is no continuity of the seams in each of these, while the data about them are very vague and incomplete. If the coal measures extend below the trap to the east, they would be close to the water carriage of the Ganges and hence transport would be cheap, but on the other hand the coal of this region is for the most part stony and bad.</p>
	<p>Deogarh—In the Janti, Sahajori, and Kandit Karaiah fields, coal of different qualities occurs. Some in the Janti field is excellent, but that known from the Sahajori area is inferior.</p>
	<p>Karharburi or Kurhurbali, in the district of Hazaribagh—This small field, having an area of 8 square miles, is of great importance on account of its position (about 200 miles from Calcutta by rail) and the good quality of its coal. The coal occurs in three principal seams, with an average total thickness of 16 feet, the estimated amount of coal is about 136,000,000 tons, while the available portion is estimated at</p>
	<p>14 000,000,000 tons. The total area exposed is about 500 square miles, but the real area is possibly even double that, as the beds dip to the east under the alluvium. This is the largest and most important coal field in which coal is worked in India, its proximity to the main line of railway, and to the port of Calcutta, tending to give it pre-eminence over other less favourably situated localities. The principal Companies en- gaged here in the extraction of coal are—the Bengal, Barakar, Equit- able, New Birbhoom, and Raniganj Association, besides many minor firms and native associations. Many of the seams are of considerable thick-</p>
	<p>C. 1418</p>

Coal fields of India

(H B Medlicott)

COAL.

ness, one containing from 70 to 80 feet of coal. As a rule, however, the best coal is not found in the very thick seams.

"*Jharia or Jeriah*—This field is situated in the valley of the Damuda river, 16 miles west of the Ranigunj field, and is nearly all included in the district of Manbhum. The thickness and quality of the seams vary a

Nothing has been done to develop the resources of this field.

"*Ramgarh*—This field, situated to the south of the Bokaro field, has an area of about 40 square miles. The coal is for the most part of poor quality and limited in extent.

There are probably 5 million

tons of the field is close to

and it is believed that some

by the natives and carried to Ranchi for sale.

"*North Karanpura*—Situated at the head of the Damuda valley, has an area of about 472 square miles, and the estimated amount of coal is 8,750 million tons.

"*South Karanpura*—Situated to the south-east of the northern field, has an area of 72 square miles, and the estimated amount of coal is 75 million tons. The assays of some of the coal indicate a high calorific power.

"*Choke*—Is a small field of less than a square mile in extent. Situated on the Hazaribagh plateau.

"*Itkuri*, 25 miles north west of Hazaribagh. A few seams of inferior coal are exposed.

"*Aurunga*—In the district of Jharkhand, in the valley of the Koel, a tributary of the Son. The area is 97 square miles, and the estimated amount of coal is 20 million tons, but the quality of the coal as taken from the outcrop is poor.

tons

"*Tatapant, Iria, and Morne*—Situated in the valley of the Son and tributaries. These fields are portions of a large tract stretching far to the westward. Several coal seams of workable thickness and many thin ones exist.

NORTH-WEST
PROVINCES.
1419

COAL.

Coal-fields of India.

CENTRAL INDIA.
1420

"*Koror*—Three miles north of Umaria. The area is 9 square miles, and a thick seam of good coal has been proved.

"*Jhilmili*—Is another area of about 41 square miles, in which seams of some promise have been observed.

"*Bisrampur*—Has an area of about 400 square miles occupying the central basin of Sarguja; it contains some good coal suitable for locomotives.

CENTRAL PROVINCES.
1421

With the other associated rocks, these occupy an area of at least 1,000 square miles, some of the seams are very thick, two being respectively 90 and 168 feet, but though including good coal they often contain a large proportion of shale, and the horizontal extension of the seams is sometimes irregular and uncertain. These fields will probably assume importance in connection with the line to connect Calcutta with the Central Provinces. The recent boring experiments show that the Korba area has proved most worthy of consideration, particularly at Ghordewa, 9 miles to west north-west of Korba, where there is a 5-foot seam of good coal.

"*Satpura Basti*—The *Mahadeo* field is of importance in Indian Peninsula.

"*Shahpur* (or *Betul*) on the south of the Tawa valley—This field

HYDERABAD.
1422

Warora basin	14	} million tons
Chugus	45	
Wun	1,500	
Between Wun and Papur	50	
Between Janara and Chicholi	75	
Sasti and Paoni (Nizam's territory)	30	

BOMBAY.
1423

of which are of

near the village of District, contains 38 feet.

other areas, 81st 1 to exist. There

The only pits worked in this wide area are at Warora, whence a special branch line conveys the coal to the Nagpur branch of the Great Indian Peninsula Railway.

"*Cutch*—There are a few thin shaly seams at Trambal (Tromba or Trombow), about 5 miles north east of Buj, in a stream north of Sis-agad, and in a stream west of Guneri near Lakhpat. Besides these jurassic seams, there are some tertiary carbonaceous layers of no promise.

C. 1423

Coal-fields of India.	(H B Medlicott)	COAL.
		...
		...
		...

purposes. The latest reports give a 6 foot seam of coal near Kosht, but the dip is said to be as high as 45° which will militate greatly against its profitable extraction.

"Chamarling, in the Luni Pathan country, about 75 miles from Dera Ghazi Khan—There are several seams of tertiary coal, of which the principal one has a thickness of 9 inches.

"Kamgram, in the " al
exists near this place, e ur
in the Ghilgai country a at

containing coal

PANJAB
1426

in places. As the locality is near a good road a fair amount of fuel might be obtained, for Bhaganwalla, the only place within 2 miles, the coal is By means of suitable machinery, and though it is in this respect. The available coal is estimated at 16,20,000 maunds (60,000 tons).

"North-West Himalayas—At Dandli, near Kotli, on the Punch, and at the north-west shoulder of the Singar Marg Mountain, there are

HIMALAYAN.
1427

COAL.

Coal-fields of India

CENTRAL
INDIA
I420CENTRAL
PROVINCES
I421HYDERABAD
I422BOMBAY
I423

28 million tons of coal, of great importance.

its command-
on the East
supply of the

square miles,

and a thick seam of good coal has been proved

"*Jhulmili*—Is another area of about 41 square miles, in which seams of some promise have been observed

"*Bisrampur*—Has an area of about 400 square miles occupying the central basin of Sarguja, it contains some good coal suitable for locomotives

"*Lakhanpur*—South of the Bisrampur area, holds some seams of good coal, the area is 50 square miles

"*Raigarh, Hingir, Udaipur and Korba* fields in the Mahanadi valley—With the other associated rocks, these occupy an area of at least 1,000 square miles, some of the seams are very thick, two being respectively 90 and 168 feet, but though including good coal they often contain a large proportion of shale and the horizontal extension of the seams is sometimes irregular and uncertain. These fields will probably assume importance in connection with the line to connect Calcutta with the Central Provinces. The recent boring experiments show that the Korba area has proved most worthy of consideration, particularly at Ghordewa, 9 miles to west north west of Korba, where there is a 5 foot seam of good coal

"*Satpura Basin south of the Nerbada Valley*—The *Mohpani* field is of importance in consequence of its position with reference to the Great Indian Peninsula Railway (95 miles by rail, west south west from Jabalpur). The coal is worked by the Nerbada Coal Company and supplied to the railway, but the supply falls short of its requirements

"*Shahpur (or Betul)* on the south of the Tawa valley—This field contains seams of coal with large quantities of lignite

of which are of

Chimur, 30 miles north east of Warora in the Chanda District, contains three seams of coal, with a maximum total thickness of 38 feet

"*Wardha (or Chanda), &c*—Includes, with several other areas, Sisti and Paoni in Hyderabad in which coal has been proved to exist. There are about 1 714 million tons of coal available, viz—

Warora basin	14
Ghugus	45
Wun	500
Bet veen Wun and Papir	50
Bet veen Janara and Ch choli	75
Sast and Paoni (Nizam's territory)	30
million tons	

The only pits worked in this wide area are at Warora whence a special branch line conveys the coal to the Nagpur branch of the Great Indian Peninsula Railway

"*Cutch*—There are a few thin shaly seams at Trambal (Tromba or Trombow), about 5 miles north east of Buj in a stream north of Sis agad and in a stream west of Guneri near Lakhpat. Besides these jurassic seams there are some tertiary carbonaceous layers of no promise

Coal-fields of India.	(H B Medlicott)	COAL
		cryst
		...
		..

purposes The latest reports give a 6 foot seam of coal near Kohat, but the dip is said to be as high as 45° which will militate greatly against its profitable extraction

"Chamarlang, in the Lum Pathan country, about 75 miles from Dera Ghazi Khan — There are several seams of tertiary coal, of which the principal one has a thickness of 9 inches

containing coal

"At
alum sha
bed of ju
"Salt

Sunglewar, Chami Kutta, Sowa Khan, Derwat, Nurpur (Nawab), and Karuli, but only in small quantities, presenting no prospect of being profitably worked At Dandot, in the neighbourhood of which coal is seen at three localities, and where thickest is 2 feet 6 inches The later develop-

PANJAB.
1426

can be delivered At Pid there is a seam of good bright fuel 3 feet thick in places As the locality is near a good road a fair amount of fuel

(60,000 tons)

"North West Himalayas — At Dandli, near Kotli, on the Punch, and at the north west shoulder of the Singar Marg Mountain, there are

HIMALAYAN.
1427

COAL.

Coal-fields of India.

its coalfield is
on the East
supply of the

square miles,

and a thick seam of good coal has been proved.

"*Jhulmili*—Is another area of about 41 square miles, in which seams of some promise have been observed.

"*Bisrampur*—Has an area of about 400 square miles occupying the central basin of Sarguja, it contains some good coal suitable for locomotives.

"*Lakhanpur*—South of the Bisrampur area, holds some seams of good coal, the area is 50 square miles

CENTRAL
INDIA.
1420

"*Raigarh, Hingir, Udaipur and Korba* fields in the Mahanadi valley.—With the other associated rocks, these occupy an area of at least 1,000 square miles, some of the seams are very thick, two being respectively 90 and 168 feet, but though including good coal they often contain a large proportion of shale, and the horizontal extension of the seams is sometimes irregular and uncertain. These fields will probably assume importance in connection with the line to connect Calcutta with the Central Provinces. The recent boring experiments show that the Korba area has proved most worthy of consideration, particularly at Ghordewa, 9 miles to west-north-west of Korba, where there is a 5 foot seam of good coal.

"*Satpura Basin, south of the Narbada Valley*—The *Mohpani* field is of importance in consequence of its position with reference to the Great Indian Peninsula Railway (95 miles by rail, west-south-west from Jabalpur). The coal is worked by the Narbada Coal Company and supplied to the railway, but the supply falls short of its requirements.

CENTRAL
PROVINCES
1421

"*Shahpur (or Betul)* on the south of the Tawa valley.—This field

of which are of

the village of
Chimur, 30 miles north-east of Warora in the Chanda District, contains three seams of coal, with a maximum total thickness of 38 feet.

"*Wardha (or Chanda), &c*—Includes, with several other areas, Sisti and Paoni in Hyderabad, in which coal has been proved to exist. There are about 1,714 million tons of coal available, viz—

Warora basin	14	} million tons
Ghugus	45	
Wun	500	
Between Wun and Papur	50	
Between Janara and Chicholi	75	
Sasti and Paoni (Nizam's territory)	30	

The only pits worked in this wide area are at Warora, whence a special branch line conveys the coal to the Nagpur branch of the Great Indian Peninsula Railway.

"*Cutch*—There are a few thin shaly seams at Trambal (Tromba or Trombow), about 5 miles north-east of Buj, in a stream north of Sis-agad, and in a stream west of Guneri near Lakhpat. Besides these jurassic seams, there are some tertiary carbonaceous layers of no promise

BOMBAY.
1423

C. 1423

Coal and Coal-mining in India.	(W. Saise)	COAL.
abandoned This seam was 11 feet 8½ inches thick, of which 6 feet 8 inches were true coal At Hienlap (or Hienlat), about 6 miles from the last locality, there is a seam from 17 to 18 feet in thickness, and the coal is of pretty uniform character with conchoidal fracture. Three		
western banks therly is 10 the seam is		
ANDAMAN. 1430 far val Part III)		
INDIAN MINES.		1431
Indian coal up to present date— INDIAN CONSUMPTION OF COAL. "The coal used either imp of coal a first, the under—		1432
Imported (1883-84) Raised in India (1884) about		Tons, 678,000 1,556,400 <hr/> 2,216,000
"The value of the former is stated to be Rs1,09,96,047. The value of the latter at the pit's mouth may be taken at Rs8,45,000. The imported coal is chiefly large or steam coal. The coal used in the taken at 1,200,000 allowed to go to wa steam and rubble		

* See page 388.

COAL.	Coal-fields of India.
<p>ASSAM, 1428</p>	<p>to Dalingkote, the coal is of Gondwana age and is much crushed, some of it is in the form of a powder, and has assumed the character of graphite</p> <p>of access</p> <p>"Upper Assam—There is an important field at <i>Makum</i> which is being worked by the Assam Trading Company, it contains several seams of coal, one of which is over 100 feet thick, 75 feet being good coal. The beds are disturbed and the coal seams lie at an average angle of about 40°, so that some difficulty may be met with in working them. An approximate estimate gives 18,000,000 tons as available, supposing the workings to be nowhere carried more than 200 yards from the face or 400 feet to the deep.</p> <p>For the most part workable seams is estimated to have 10,000,000 tons of coal available, this is exclusive of what may be proved by borings, but is mostly of poor quality</p> <p>"Nasira, in Upper Assam—Some of the seams in this field are of considerable thickness, 30 feet and over, the estimated quantity available is 10,000,000 tons</p> <p>"<i>Fanjit and Disai</i>—Two small and unimportant fields in Upper Assam</p>
<p>BURMA, 1429</p>	<p>on the Cheduba Land</p> <p>"Pegu—Coal was discovered in 1855, and a mine opened at <i>Thayetmyo</i>, but after a few cwts had been extracted, the work was abandoned recently and in fallacious hopes of a source of fuel, have been met with division of the <i>Henzada</i> district several seams of coal occur</p> <p>"Tenasserim—C of localities those a Thoo-hte khyoung (o where a mine was</p>

Coal and Coal mining in India	(W Suse)	COAL.
abandoned This seam was 11 feet 8½ inches thick, of which 6 feet 8 inches were true coal At Hienlap (or Hienlat) about 6 miles from the last locality, there is a seam from 17 to 18 feet in thickness, and the coal is of pretty uniform character with conchoidal fracture Three		

taceous coal it is well situated for transit purposes On the Paulwing river there are numerous irregular thick seams of tertiary coal

In the Andaman and Nicobar Islands coal is known to exist, but so far as they have been examined there are no grounds for belief that a valuable deposit of coal occurs (See *Manual of the Geology of India, Part III*)

ANDAMAN

1430

INDIAN MINES

1431

Dr Walter Saise Manager, E. I. R. Company's Collieries, has obligingly furnished the following note on Coal and Coal mining in India which, it may here be remarked, is based on the results of 1883-84 but on returns some of which are not accessible to Government This explanation accounts for the apparent discrepancies between the returns of production and consumption published by Government for that year and the figures here given by Dr Saise On a further page will be found more recent figures abstracted from Government returns which bring this brief note on Indian coal up to present date —

1432

INDIAN CONSUMPTION OF COAL — The coal and coke used in India are either imported or raised and made in the country The foreign sources of coal and coke supply are Europe, Australia and Africa Taking coal first, the proportion of coal raised in the country and that imported is as under—

Imported (1883-84)	Tons, 678 000
Raised in India (1884) about	1 556 400
	<u>2,216 000</u>

The value of the former is stated to be Rs 1 09,96 047 The value of

COAL

Coal and Coal-mining in India

to a smaller extent. The small kinds of rubble or smithy are used in

I433

Below is a table of ultimate analyses of specimens from Karharbari and Raniganj coal-fields with analysis of English and Welsh coals for comparison —

COAL FIELD	Carbon	Hydrogen	Oxygen and Nitrogen	Sulphur	Ash	
Karharbari	78.20	4.34	7.89	0.42	9.15	Main Seam Upper Seam
E I Railway	70.93	4.10	12.49	0.52	11.96	
Raniganj (N. B. Coal Co.)	74.31	5.12	9.67	0.47	10.43	
England { Newcastle	82.83	5.32	7.13	1.17	3.55	
{ South Wales	88.47	4.59	3.02	1.25	3.09	

"It will be noticed that in several particulars Indian coal is inferior to English, *1st*, in containing more ash, and *2nd*, less carbon and hydrogen

"In the table below the commercial analyses of many Indian coals by the writer and Mr T H Ward, FGS, are given, as also commercial analyses of Newcastle and Welsh coals, for comparison —

COAL-FIELD	Spec gravity	Ash	Fixed carbon	Volatile matter	Sulphur	Heating power by Thomson's calorimeter	REMARKS.
BENGAL { Karhar						13.20	Not worked
{ Raniganj						12.50	
						12.89	
						13.89	
						12.35	
Tindana						12.40	Not worked
CENTRAL { (1885)							
Welsh	1.312	3.68	82.66	13.66	1.59		
Newcastle	...	3.49	63.25	33.26	1.07		

"The above table shows that there is great diversity in the chemistry of the coals of India, and the variations in physical features are just as marked. With the exception of Tindana and Assam coal, all Indian coals are remarkably laminated in structure, the laminae consisting of a dark highly

Coal and Coal mining in India

(W. Sasse)

COAL

carbonaceous shale, a bright pitch looking matter, and a mineral charcoal — a very dull charcoal looking substance. When these laminae are very

volatile matter

COMPARISON OF INDIAN WITH IMPORTED COAL FOR RAILWAY PURPOSES — The Indian and imported coals have been tried on Indian Railways with the following results —

EAST INDIAN RAILWAY

COAL	Gross weight of trains	lb per ton of coal consumed	lb per ton mile
	Tons cwt		
Ka ha bar	207 19	30 12	145
Ran ganj Sanctor a	212 17	32 21	151
Equ table	208 1	33 68	161
O d nary	204 14	36 98	181
North Wales	215 9	31 90	148
South Wales Ca d ff	203 11	32 64	160
New South Wales	207 14	31 43	151

D W CAMPBELL

Locomotive Supdt, East Indian Railway

COAL	Gross weight of trains	lb per ton of coal consumed	lb per ton mile
	Tons cwt		
Ka ha bar	166 12	25 76	155
Ran ganj	181 7	33 33	184
Barakar	170 3	30 04	177
Lothargills (S W)	183 12	30 45	165
North Wales	174 9	27 12	156
Austral a	180 4	27 43	133
Duckfield			
Merthyr			
Godavari	171 12	33 48	196

F H TREVETHICK,

Locomotive Supdt, Madras Railway

COAL.

Coal and Coal-mining in India.

"It will be seen from these results that Karharbari coal is a good steam coal, little inferior to imported coals, and that the other Indian coals (except Godavari) are of fair quality. Umaria coal, tried on the Great Indian Peninsula, gave 42.63 lb per train mile with a gross load of 410 tons. This is nearly but not quite as good as Karharbari coal.

INDIAN PRODUCTION—"The sources of Indian coal supply and the estimated yearly output are as under:—

CENTRAL PROVINCES	{	Warora	100,000
		Narbada	28,000
		Umaria	7,290
BENGAL	{	Karharbari	520,000
		Raniganj	890,000
ASSAM	50,000
								<u>1,595,200*</u>

As the newer fields develop this estimate will have to be increased.

DISTRIBUTION OF INDIAN SUPPLY—"The Warora coal-field is connected with the Nagpur branch of the Great Indian Peninsula by the Wardha Coal State Railway, the Mohpani (Narbada) coal-field by a branch from Gadawara with the Great Indian Peninsular. The Umaria coal-field has been tapped by the new line from Kutni through the East Indian Railway, Jubbulpur line. The Assam coal field is connected

the fol-
Wardha
al going

"The Bengal coal finds its way to the Panjab railways and the railways of Bengal, as also into the manufactories of Calcutta and the large cities along the line of railway. Some is used in the steam ship lines. Small coal is largely employed for brick making. Comparatively little is utilized for domestic purposes. The Colliery Companies should endeavour to create a want by teaching the people how to use small coal in large towns, such as Allahabad instead of wood and cowdung. Agencies like those in English cities could probably do this in a few years, and the large waste of small coal that goes on at present would thus be obviated.

MINING IN INDIA.

"Has made considerable progress during the past few years, machinery and well-appointed heapsteads and pit frames are coming generally into use

which is 402 feet deep

"The system of working varies very much. At Warora, Central Provinces, where 100,000 tons per annum is wound by direct acting engines out of two shafts 200 feet deep, the system most nearly approaches the

* It may be noted that it is the marketable coal that appears in the Government returns, not the actual amounts raised. In 1883-84 these were 1,200,957 tons. Conf. with p. 385.—Ed

Coal and Coal-mining in India (H' Suse)	COAL.
---	-------

English
day morn
shifts of 8½
thus Gai

in height, leaving the roof coal, and pillars 40 feet square. The coal is so hard, it has to be nicked and undercut and then blasted down. The pillars are worked by splitting each from one headway to another and then taking the far end off in slices. The roof coal comes with it.

"At the Mohpani collieries a similar system is worked. The difficulties met with in these mines, owing to the faulted and disturbed nature of strata, are probably unequalled in India.

Karharbiri coal-field—"Is the smallest field in Bengal. It is mainly worked by three Companies, the Karharbiri Coal Company, and connected with the main line worked by locomotives.

a scene of great activity. As much as 50,000 tons of coal and coke have been raised and despatched in one month. The coal field is connected with the East Indian Railway Chord line by a branch from Madhupur to Girdi, the terminus or colliery station. In mechanical arrangements for raising coal, this coal field is well advanced. The old fashioned gin is almost obsolete and bullock-carts have little to do.

"The system here is similar to that obtaining all over Bengal. The

and make are now universal, the crowbar and single pick having been ousted. The workings are on the bord and pillar system. Pillars vary from 12 feet to 40 feet square and 40 feet X 60 feet. In the shallow mines and thin seams (7 to 8 feet) the former size obtains, in the thick seams (from 12 to 20 feet thick) the latter. Pillars are worked in the 8 feet seam in the following manner. A 4 feet chock is placed between each pillar in the row of pillars (generally six in number) that are to come out. A chock is also placed in front of each pillar. The pillar is then attacked from the front side. When pillars are taken out the chocks are withdrawn and the roof falls.

"The system here is similar to that obtaining all over Bengal. The

the Bauris are not in such requisition as formerly.

"Drainage is effectively carried out by Tangye's special and lifting and forcing pumps, worked by bob levers from horizontal engines. The machinery is of good type, and winding and hauling are done by good engines.

"Ventilation is attended to in the deep mines, mainly by furnaces or steam jets.

Coal and Coal-mining in India.	(W. Saise)	COAL.
<p>considerable -- Coal -- of the seam 12 feet to 16 the roof, r insists (anc mening o height of the seam has been excavated. His chief and dearly-prized weapon is a 'sabal' or crowbar with a sharp point at one end. With this he smashes the coal, standing always when at work. He never grooves beyond the first 'cleat,' gangs of 4 or 5 men occupy each gallery; they are paid b tom caste (</p>	<p>izes to support ie native coolie field) on com- until the full</p>	
<p>ployed tram or bucket The women often take their babes, 2 and 3 months old, down the mine, taking with them also a small cot on which the child sleeps or plays while its parents are at work</p>		

coal get 'won' being from much less depths Some fire damp has been met with in the western part of the district Chanch colliery (west of the Barakar) belonging to the Bengal Coal Company was abandoned some years ago after the coal had been burnt, some of the coal was burnt, some of the coal was burnt, some of the coal was burnt.

idly been men-
the outcrops

merely, of these magnificent seams, and thousands of tons remain still to be worked without in

"The 'Bauri' is the district In some amusing like those of drunk, especially at week on Mondays For the a difficult matter to per (contract) rate for his do more than will, with his wife's contribution, keep the household 'in rice' and himself in drink for the day. The nearly universal and very bad custom in this district is to pay each evening for the work done during the day The collier or coolie has often to wait about 8 or 9 P.M. for his money. He then goes to his home and takes a

house is often substituted). In this he places a small quantity of oil and

COAL.

Trade in Coal.

a portion of wick. Any oil he can save from his 'allowance' is his

1438

The ignorant native has not yet recognised that his health and longevity is in question, and he has besides helped much to prevent ventilation becoming a necessity by the wonderful power of endurance he has shown. This power of endurance enables him to work for hours at the bottom of a sinking shaft with water pouring over his naked body or to work all

India employs about 30,000 persons, the quantity of coal raised per annum per person employed, surface and underground, being 51 tons.

"In Europe the numbers are different, varying with the thickness of seams and nature of difficulties met with :

England (average)	348 tons per person employed under-ground and surface per annum.
Belgium	134 Ditto
Saarbrücken	187 Ditto

There is no Government regulation of the coal industry ; any person can manage a mine on any system he likes, whether or not he has experience or training. Interest has a great deal with the appointment of the managing staff, and it is to be feared that the best is not made of the splendid coal deposits, the favourable roof, and the moderate depths and inclinations of the seams "

1439

TRADE IN COAL.

The following brief note, prepared by the Revenue and Agricultural

YEAR	COAL.		Coke	Patent Fuel	Wood
	English	Country			
	Tons	Tons	Tons	Tons	Tons.
1887	212,529	479,210	9,564	30,029	292,808
1886	240,063	460,948	9,132	26,212	259,513
1885	225,721	476,277	10,439	23,117	255,178

C. 1439

Trade in Coal.

COAL.

	however, 37 were in Umeria in Rewa district worked The
	Tons
Bengal	1,187,000
Central Provinces	117,300
Assam	70,800
Central India	13,500
TOTAL	1,388,600

Assam has since increased its output, the figures for 1886-87 being returned at 72,000 tons. It is stated in the Railway Administration Report for 1886-87 that—

"Coal continues to enjoy the confidence of the public. Its sale to the river steamers and tea factories for by the Dacca State Railway, the Eastern Bengal State Railway. It is being largely enquired for by Calcutta, also by the Eastern Railway. It has been found suitable to the engines and the Northern Bengal State Railway, but the difficulty of access to these two railways from the river Brahmaputra prevents its extensive use by their administrations. The coal continues dusty, though it is being mined deep in the hill sides. But its nature is beginning to be understood, and its friability is not found to be a drawback to its use as a steam fuel."

"The coke is found to be saleable to the tea factories of Lakhimpur to an extent of about 3,000 tons per annum. The Company is preparing by means of an increased labour force to enlarge the output of coal to 100,000 tons yearly."

Collieries have recently been opened out at Dandot (Panjab) and Singarem (Nizam's Territory). The coal in these mines has been pronounced of good quality, and in Upper Burma coal has been found (in the Kali Valley on the Chindwin River), but arrangements have not as

The commencement of this industry appears to date back to 1820, when a mine was opened in the Raniganj district in Bengal. For twenty years no new mine seems to have been opened, and then only three mines were opened down to 1854. In that year the commencement of the Eastern Indian Railway line which was laid to run through the coal bearing regions of the Damodar basin.

In the paragraph above the number of mines in 1886-87 is stated to

COAL

Trade in Coal

doubled themselves since 1866-67, having risen from 341 000 tons, valued at Rs55 lakhs, in that year to 765 000 tons valued at Rs130 lakhs in 1886-87. The United Kingdom supplies nearly all the imported coal, though Australia, which ranks next to it as a source of supply, is now sharing more largely in the imports, the value of its consignments in 1886-87 being Rs 22 lakhs. The percentage taken by each province in these imports is noted on the margin.

Bombay	10.4
Lower Burma	9.5
Bengal	4.7
Madras	4.2
Sind	

INTERNAL
TRADE
1441

INTERNAL TRADE—Statistics may now be given regarding the internal movements of coal by rail during 1886-87 between the different blocks (i.e., provinces, chief towns and Native States). The total trade amounted in quantity to 1 097 800 tons and in value to Rs158 83 lakhs. The position of each block as a net exporting or importing centre may be thus indicated—

Exports,	Tons	Imports	Tons
Bengal	743 000	Calcutta	304 000
Bombay Town	162 000	Bombay Pres. dency.	167 000
Central Provinces	44 000	North Western Pro-	
Karachi	7 000	vinces and Oudh	161 000
Assam	4 000	Rajputana and Cen-	
Madras Town	2 000	tral India	66 000
Madras	1 000	Punjab	35 000
		Benar	23 000
		Sind	5 000
		Mysore	4 000
		Nizam's Territory	3 000

As might be expected, Bengal, where the most extensive mines in India are situated takes the lead among the exporting centres. Of its

Mysore from Madras and the Nizam's Territory from Bombay Town.

The development of the coal industry in India is indicated by the fact that the gross exports from Bengal to other provinces and Calcutta have increased from 641 807 tons in 1882-83 to 755 831 tons in 1886-87 and those from the Central Provinces from 26 451 tons to 56,125 tons during the same period. Assam for the first time shows a net export (4,000 tons) in referring to which the Director of Land Records and Agriculture writes—'This is entirely due to the increased output of the Alakum coal mines near Dibrugarh, which now supply nearly all the coal used in the Assam Valley besides furnishing large quantities for export.'

1442

Coke (A note contributed by Dr W. Sasse.)

Coke is imported and also made in India. In 1883-84 the imports amounted to 16 700 tons valued at Rs4 to 738. Coke, however, is now made to a very large extent in Bengal. It is a most important industry in

C 1442

Cobalt

COBALT.

its relation to coal raisings as the manufacture of coke means the utilization of small and otherwise useless coal. The industry is of recent and very rapid growth having increased fourfold since 1875. There are two kinds of coke called respectively hard and soft. The former is dense and is

less expenditure of coal. Soft coke is incompletely burnt coal, made for the purpose of the chart.

except

and a show that in a year about 55,000 tons of coke exclusive of foreign coke are led over the line, add to this the own consumption, the respect

1443

Hard coke for foundry blast furnaces locomotive, &c.
Soft coke

65,800

11,200

per annum. The point of great plant of washing of the market collieries

COBALT

1444

Cobalt; Ball, Econ Geol, 324 & 616, also Mallet, Mineralogy, 27

Cobalt metal is never met with in the native form, except in small proportions as a constituent of —

chiefly in primitive rocks and
to a nickel iron and often by
sulphur or by arsenic or by
Spe ss Cobalt or tin white Co
Linnæite or Cobalt Pyrites $\text{CoS} + \text{Co}_2\text{S}_3$

Vern — The

a

t

SOURCE — A complex mineral (*selita*) is found in various mines in Rajputana especially in those of Babul and Bagor near Khetr. Mr Mallet says of this substance that it has the specific gravity of 6.00. On analysis it yielded the following composition —

Sulphur	19.46
Arsenic	43.87
Antimony	a trace
Nickel	a trace
Cobalt	25.30
Iron	7.83
Gangue	50

100.26

1445

C 1445

COBALT.

Source of Cobalt.

This substance is generally known as Cobaltite. In the *Rajputana Gazetteer*, and in the *Fury Reports* of the Exhibition of 1862, occur accounts of the Jeypore enamels, but in a recent publication, Dr. T. H. Hendley (*Journal of Indian Art*), gives more precise details. Sir George Birdwood (in his *Industrial Arts of India*) under Enamels (pages 163-165) and also under Pottery (pages 301-324), gives most instructive parti-

scribed Cobaltite, in the Records of the Geo'ogical Department, seem to be unanimous in their opinion that Cobalt is only rarely met with in India, and that, too, in the mines of Rajputana alone (as far as peninsular India is concerned), and that the oxide is artificially prepared; in other words, that it does not occur naturally in Central and Southern India.

1446

will be found some account of the uses of cobalt in the ceramic industry while

Dr Hendley says that the colours used by the Jeypore enamellers "are obtained in opaque & treous masses from Lahore, where they are prepared by Muhammadan *manihars* or bracelet makers. The Jeypore workmen state that they cannot make the colours themselves. The base of each colour is vitreous and the colouring matter is the oxide of a metal such as cobalt or iron. Large quantities of cobalt are obtained from Bhagore near Khetri, the chief town of a tributary State of Jeypore, and are used in producing the beautiful blue enamel." In these passages Dr Hendley does not make it quite clear whether the Jeypore enamellers prepare their own material for the blue colour, though unable to prepare the other colours, or whether the entire mass of the crude material is conveyed to Lahore and other centres to be prepared and returned in its manufac-

cusses the Multán enamel industry and furnishes particulars regarding the *Mina* blue vitreous enamel. In the *Multán Gazetteer* (p. 107) this subject is enlarged upon, and reference is also made to the Bahawalpur enamels, where, in addition to opaques, a semi-translucent sea green and also a dark blue are produced.

Cocculus

COCCULUS
villosus.

In Europe Cobalt is largely used as a pigment and to colour ordinary glass

Coccinia indica, W. & A., see *Cephalandra indica*, Nand, CUCURBITACEÆ

COCCULUS, DC, Gen Pl, I, 36, 961

I447

Cocculus cordifolius, DC, see *Tinospora cordifolia*, Miers, MENIS- [PERMACÆE

C. indicus (see *Fluck and Hanb, Pharm, p 31*), a commercial synonym for *Anamista Cocculus*, W & A., see Vol. I., A. 1037.

C. Leæba, DC, Fl Br Ind, I, 102

I448

Vern — *lallur, illar billar, parwatti, vehri*, PB, *Ullar billar*, SIND

References — *Gamble Man Timb, 11 Brandis, For Fl o Stewart, Pb Pl 6 Aitchison, Cat Pb and Sind Pl, 3, Murray, Pl and Drugs, Sind 38*

Habitat — A large climber of the dry and arid zones, especially of Western India — the Panjab, Sind, and the Carnatic

Medicine — Stewart says the stems often become as much as 3 or 4 feet in girth. It is used in Sind and Afghanistan in the treatment of intermittent fevers and as a substitute for *Cocculus indicus* (*Murray, Dymock*)

MEDICINE,
I449

Food and Fodder — In the Trans Indus, Stewart says, it is browsed by goats but by no other animals. Said to be used as a partial substitute for hops in the manufacture of Indian beer (*Murray*)

FOOD and
FODDER
I450
Hop
Substitute.
I451

C. palmatus, DC, see *Jateorhiza palmata*, Miers

C. villosus, DC, Fl Br Ind, I, 101.

I452

Vern — *gamti ki bel, hier, dier*, HIND, *Kursan, samir*, SIND, *Vasana vela*, MAR, *Wassanwel parwel*, BOMB, *Kattuk kodi*, TAN, *Dusari tige chipuru tige, kattle-tige*, TEL. In the Concan the Vaidas give this plant the Sans. name of *Vanatiktika*

Th. plant — — — — —

by the leaves of *Cocculus villosus*

References — *Gamble, Man Timb, 11 Roxb, Fl Ind, Ed C B C, 732, (under Menispermum hirsutum, Willd), Drury, U Pl, 145, Dymock, Mat Med W Ind, 2nd Ed, 32*

Habitat — A large climber of the dry and arid zones, Sind, Panjab, Deccan, extending into Madras and Bengal

MEDICINE.
Leaves
I453

rheumatic and old venereal pains, half a pint every morning is the dose. It is reckoned heating, laxative, and sudorific. By more recent writers the root is said to be alterative and to be a good substitute for sarsaparilla. Dymock remarks that in the Concan the roots rubbed with Bonduc nuts in water are administered as a cure for belly-ache in child-

Roots.
I454

C. I454

COCCUS
cacti.

The Cochineal Insect.

FOOD.
I455

this as a Sind drug under its bazar name of *zamir*, and remarks that it is employed in pains of the head

Food.—The leaves are made into curry and eaten by patients under treatment, with the roots or the jelly from the leaves. If suffered to stand for a few minutes, the jelly clears, "the gelatinous or mucilaginous parts separate, contract and float in the centre, leaving the water clear like Madeira wine, and almost tasteless" (*Roxb*). With regard to this property the remark under the vernacular name *Farid-biti* should be read. In Eastern Bengal the writer repeatedly observed the milkmen carrying milk to market with a few leaves of this plant and the spine-like leaflets of the date-palm placed in the vessel. On enquiry he was told these prevented the milk from getting bad through the heat and the shaking to which it was subjected. It has been observed that the plant is not eaten

FODDER.
I456
DOMESTIC
I457

during the famine of 1877-78 in the Khandesh district, and that it is

e plant.
good,

outcrop, bluish purple ink (*Roxb*)

COCCUS; Packard, Guide to the Study of Insects, 526.

A genus of Insects belonging to the Coccidæ of the Order Hemiptera. Several species are, by Entomologists, referred to this genus, but two only are of commercial importance,—the one a native of Southern Asia and the other of the

will be found

spherical scale

I458

Coccus cacti, Linn

THE COCHINEAL INSECT; COCHENILLE, *Fr*; KOCHENILLE
SCHARLACHWURM, *Germ*; COCCINIGLIA, *It*, COCHINILLA, *Sp*

Vern — *Airmdana*, BENG, *Kirmas*, BOMB, *Kiranda*, N.-W. P., *Kirm*, PB

References — *Royle, Prod Res of Ind*, 57, *Encyclop Britannica*, VI, 97;
Balfour, Cycl of India; *Listard Dyes and Tans of India*, *Wardle*,
Report on the Dyes of India, *Buck, Dyes and Tans of N.-W. P.*

The Cochineal Insect.

COCCUS
cacti.

Official Papers on Pigments used in India, Crookes, Dyeing and Calico Printing, 350, Hummel, the Dyeing of Textile Fabrics, 348;

Habitat.—The Cochineal insect was first discovered by the Spaniards in Mexico in the year 1518, but it was not made known to Europe until 1523. At first it was supposed to be a seed, but in 1703 Leeuwenhoek showed it to be an insect. In Mexico it is particularly abundant in the provinces of Oaxaca and Guerrero. It occurs in many localities in Central America, and for long has been one of the most important articles of export from Guatemala, but it is met with also in South America, and recently it has been found (or perhaps only an allied insect) in the West Indies and in the southern portions of the United States.

HISTORY AND INTRODUCTION.—The immense importance of the trade, early established in this insect, led to efforts for its propagation in other countries, and for many years this has been profitably prosecuted in Tenerife, the Canary Islands, Java, Algeria, and to some extent even in Spain. According to some writers the best quality now comes from Honduras. The attention of the Court of Directors of the East India Company was directed to this subject by Dr James Anderson of Madras in 1756. He forwarded to Sir Joseph Banks samples of a dye-yielding insect which was proved to be a species of *Coccus*, but not Cochineal.

HISTORY.
1459

species of *Cactus* or *Opuntia*. On the China and Manilla species of the Nopal, and even on that from Kew, the survivors began to die fast. It is fortunate that they have not all perished, for they are the only plants seen growing on the plantations. The plant is now being propagated.

COCCUS
cacti.

The Cochineal Insect.

HISTORY.

to the discovery of America, and therefore no *Cactus* can be called indigenous to India. This is more than a quibble as to the correct usage of a scientific term. If the *Coccus* sent to Sir Joseph Banks, one hundred years ago, was found feeding on a *Cactus*, it must be regarded as but an earlier introduction than the Cochineal brought to India by Captain Neilson. It therefore seems probable that the Portuguese (or whoever introduced the *Opuntia*) may have intentionally or unintentionally brought the *Cactus*-feeding *Coccus* also. In 1848 Dr Dempster addressed a letter to the Governor General of India which afterwards appeared in the Journal of the Agri-Horticultural Society. He there extols the superior quality of the dye obtained from "the native" or "indigenous" insect as compared with the imported. "The quality," he says, "of native Cochineal which I found capable of dyeing a certain weight of woollen cloth proves that the indigenous insects contain an

Jullunder Doab "as to become a nuisance, and rewards were offered for its extermination, which, however, were rendered unnecessary shortly after, as a large number of insects of some kind of *Coccus* appeared and soon effected the destruction of the plant, which is now only occasionally to be met with."

"The *Coccus* of India enters into

" "

species of *Opuntia*; but as we have abundance of the South American plant, *O. cochimifera*, that species may also be tried along with the several sorts of our own."

"The *Coccus* of India enters into

"The *Coccus* of India enters into
ara
real
the
ind
the
nd
uld
the

to feed upon a sp
be allied to the *C*
feeding on Tamar
exudation known as Manna

THE INTRODUCTION OF THE *OPUNTIA* OR PRICKLY-PEAR.—The above remarks may be accepted as disposing of the question of "the indigenous cochineal insect which feeds on the common prickly-pear." If not indigenous then, as an acclimatised insect, has it deteriorated after

C. 1460

Reintroduction of the Cochineal Insect.

COCCUS
cacti.

the lapse of 100 to 150 years? Perhaps the further question may also be suggested—was the insect derived from the best stock? If unfavourable answers have to be given to these enquiries, then it would remain to be ascertained by actual experiment whether an improved and fresh stock could be re-introduced. We shall return to this question later on, but it may

Madras
Cochineal
Plant.
I461

Europe, and at the same time the head quarters of the acclimatised *Opuntias*. The sudden appearance and disappearance of a *Coccus* in the Panjab, mentioned by Mr. Baden Powell, would justify the conclusion

Panjab
Cochineal
Plant.
I462

Bombay
Cochineal
Plant.
I463

MODERN EFFORTS TO REINTRODUCE THE COCHINEAL INSECT.

I464

Grana sylvestris. A voluminous correspondence has ensued since 1795 as to the desirability of introducing the superior quality, which fetches (from its greater amount of the tinctorial principle) three times the price

C. I464

COCCUS
cacti.

Forms of Cochineal.

paid for the wild insect. As late as 1882, the Madras Government had this subject brought to its attention, and instructions were given that

FORMS OF COCHINEAL.

definite opinion than that a superior or an inferior cochineal was found in certain districts. The old system of having a scientific establishment to settle the system of the

1845, sundry attempts were made by the late M. Bunt and others, after considerable expense incurred, and a heavy amount of correspondence, as usual in such cases, the whole ended in smoke" (*Fl Bomb Supp*, 40)

GRANA FINA AND GRANA SYLVESTRIS.—Humboldt was, perhaps, the earliest observer to distinguish "the *fine* from the *silvester* or wild sort of cochineal." The former insect, he says, is mealy, or covered with a white powder, while the latter is enveloped in a thick cottony substance which prevents the rings of the insect being seen. The *Grana fina* is reported to be a native of Mexico, and the *Grana sylvestris* of South America. Dr. Balfour remarks, "It has been mentioned that at Vizagapatam

Grana Fina
I465
Grana sylvestris
I466

Red-flowered
Opuntia
I467

Yellow-
flowered
Opuntia.
I468

yellow flowering prickly pear or 'Opuntia' I have seen it tried at Benary and fail." Commenting on this, Mr. Liotard remarks (and he has been followed by several more recent writers) "Regarding the future in India, it may be well to lay stress on the statement made by Dr. Balfour that

C. I468

Peculiarities of the Cochineal Insect.

COCCUS
cacti.

the true cochineal insect only destroys the prickly pear plant with red flowers and few prickles, and will not propagate on the yellow-flowered plant or *Opuntia*." Again, "as regards the Peninsular, we learn from Dr. Balfour that not only the *variety* (*sic*) of plant required but the superior *species* (*sic*) of the insect also exists in parts of the Madras Presidency." Although Dr. Balfour's remark as to the existence of the true

can
be
t on
If
Balfour be correct in the statement that the latter insect does actually

Steps to be

on the red-flowered cactus is or is not a race derived from the true cochineal insect, perhaps more ancient than Captain Neilson's stock. The position assumed by Mr. Liotard of urging the extended cultivation of

1469

fostered in anticipation of the arrival of a fresh importation. Degeneration, if established, might be accounted for by an originally semi-domesticated creature having been allowed to run wild for a century or more, or from having been forced to feed on the wrong plant. Mistakes may thus be made, but the course indicated would most probably prove the most direct, and it may happen that we possess a long-acclimatised stock which, under careful treatment, would prove more hopeful than any insect that might now be introduced

1470

COCCUS
cacti.

Propagation of the Cochineal Insect.

Male.
1471

"The male also adheres to the plant, and in about 12 days becomes enveloped in a cottony cylindrical purse, open at the bottom; the insects huddle together one upon another to form a mass, with their heads out."

Female.
1472

They appear generally mouth are quite sunk in as are almost covered by

may be detected from the at the C. I. 1. 1. ed on her ex-
cording to the
indicate the

at the moment of her fixing upon the plant,
of a mouth she has
introduces into the
uch is her excessive
again. After shed-
becomes a mere shell,
emale commences to

Cochineal
nesting.
1473

shed her young that measures are taken to remove the young to other cactus leaves. A nest is formed, in the shape of a sausage or purse, of cotton gauze or other tissue pierced with small holes, in which 8 or 10 of the females are put, and the purse is fastened at the bottom of a leaf of cactus. The nest is then covered with a layer of cotton wool.

cochineal mother produces above a hundred young ones; but the mortality is great, and three or four mothers are required to cover one side of a cactus leaf with sufficient young for cultivation."

1474

PROPAGATION.

In an interesting pamphlet written by I. S. C. D. and published by the Government, much useful information has been brought together regarding the various systems pursued in America and other countries in the propagation both of the insect and the plant. We cannot afford space to deal with this subject, and must accept the above abstract of the

C. 1474

The Cochineal Dye.

COCCUS
cacti.

life-history of the insect is indicating the great governing factors with
 The following
 useful "The
 to which the

Collection
I475

leaving only one or two of these insects on the branches is fatal to the health of the plant " "The cactus cannot bear much water when not strengthened with manure " "When a plantation is reserved for the production of a winter crop, the leaves should be covered with cochineal in the month of October or November, by planting the young cochineal at this season it ripens, and is ready for gathering at the latter end of February or of March Another part of the plantation is reserved for receiving the seed at this season, but as the plants cannot be forced to bud during the winter, the seed must be planted in March upon last year's leaves, which have the disadvantage of being tough for the insect, and this renders a winter crop more precarious than one obtained in summer " Wind and rain are very destructive hence a region with a pronounced rainy season would either be unsuitable or the seed-stock at least

Propagation.
I476Suitable
Climate
I477

COCHINEAL DYE.

Mr Wardle, in his recent *Report on the Dyes of India*, mentions experiments performed by him with several samples. Of a Hyderabad sample he says, it "appears to be very good" "The Government report, in which reference is made to it, is by Major W Tweedie " "It would be interesting to ascertain whether the cochineal is produced in the Hyderabad Residency, or is imported from South America " Of

Treatment of
Crop
I478DYE
I479

sists of insects matted together by some dark-coloured substance Both samples small and poor " Reference has already been made to Dr Dempster's report on cochineal from the lower North-Western Himalayas He says "It is beyond all doubt a true *Coccus cacti*, and

and further, it is the same as the cochineal is not the Mexican insect Dr Dempster continues "In the month of December the young brood were extremely numerous, very lively, and ready to leave the mother and spread themselves over the plant Sulphate of alumina, added to an alkaline solution of the colouring matter of the native (*sic*) cochineal,

C. I479

COCCUS
cacti.

The Cochineal Dye.

1480

Europe-dyed scarlet broadcloth" "I find here an imported cochineal
from Bombay the same as the one used in Europe."

cloth proves that the indigenous insects contain an amount of colouring
matter not inferior to the fine Mexican cochineal." This statement is
so completely at variance with the opinions of all other European writers,
that the

in this village are lined with magnificent specimens of the cactus, far
superior to any I have seen since I left Ludianah, and their leaves are
covered with the cochineal insect, which, it strikes me, attains here, prob-
ably from good feeding, a larger size than I have ever seen it do before.
As I passed these hedges of the prickly pear, numerous Kashmiris were
scraping the cochineal with a blunt iron instrument from the surface of
the leaves into
asking them what
to the Amritsar
ser (2lb) of the

1431

seen by Dr Fleming was the *Grana sylvestris*.

1482

that weight. These two figures are almost alternately given by different
writers—a fact which may be accounted for by the larger or smaller size
of the different breeds of insects

It is little
erly much em-
ced by the use
duction of the
Two different
shades of red are obtained from cochineal, namely, a bluish red, called

C. 1482

Cochineal as a Medicine.

COCCUS
cacti.

crimson, and a yellowish or fiery red, called *scarlet*. Wool mordanted with 2 per cent of bichromate of potash and dyed in a separate bath receives a good purple, the colour being darkened by the addition of sulphuric acid to the mordant. Mr Hummel gives particulars of the dyeing for crimson or scarlet. Wool to be dyed the former colour is mordanted with aluminum sulphate and tartar, the dyeing being effected in a separate bath. There are other methods, but the above is perhaps the best. Lime-salts are not beneficial. The latter shade is produced by the acid of stannous salt and cream of tartar or oxalic acid. The mordanting may be performed separately or along with the cochineal.

Wool dyeing
1483

For *silk* the mordant is alum, to be worked into the fabric for half an hour and steeped overnight. The fabric is then washed and dried and dyed in a separate bath. This gives the crimson. For the scarlet, after boiling and washing the silk is first grounded with a light yellow produced with soap and arnatto and thereafter washed. For darker shades soap should not be used. In both cases the fabric should be mordanted by the same process as described or the crimson, only using nitro-muriate of tin in place of alum. By the aid of iron mordants fine shades of lilac may be obtained.

Silk dyeing
1484Pigments.
1485

part amber, and 2 parts linseed oil

(For Ammoniacal Cochineal see under paragraph of Chemistry)

COCHINEAL AS A MEDICINE

Medicine.—Cochineal is used mainly as an agent for colouring drugs, but it is supposed by some to possess anti-spasmodic and anodyne properties.

MEDICINE
1486

Chemical Composition—As far as has been determined, cochineal and lac owe their tinctorial properties to an acid apparently identical in character. This is formed within the body of the female insect. The chemical examination of this substance has revealed somewhat conflicting results—a fact which has led certain writers to presume that its composition varies. Pelletier and Caventon isolated the acid from cochineal and called it carmine, a nitrogenous compound which they expressed by the formula $C_{14}H_{14}NO_8$. Subsequent observers (Arppe, Warren de la Rue, Hugo Müller, &c.) showed it to be an acid, and found that, in a perfectly pure state, it does not contain nitrogen, though accompanied by nitrogenous matter which it is difficult to separate from it. John named the colouring principle cochineal. The acid of the authors named has been expressed as $C_{14}H_{14}O_{10}$, but the crystalline carminic acid isolated by Dr Schützenberger is given as $C_9H_8O_6$, the same substance being expressed by Dr Schaller as $C_9H_8O_6$. Most recent writers give its formula as $C_{17}H_{11}O_{10}$ (Crookes). It may be separated from cochineal by precipitating its aqueous extract with plumbic acetate and decomposing the washed precipitate with sulphuric acid. The solution thus obtained is alternately precipitated, and the precipitate decomposed, a second and a third time in a similar manner, employing, however, hydric sulphide to effect the final decomposition. The filtered solution is evaporated to dryness, the residue dissolved in alcohol, and the crystalline nodules of carminic acid

CHEMISTRY.
1487

C. 1487

COCCUS
cacti.

Trade in Cochineal.

obtained on allowing this solution to evaporate treated with water (*Miller, Elements of Chemistry, P III, 690*) This same substance has been found in the flowers of *Monarda didyma* and probably in other plants. Pure carminic acid is a purplish-red substance, which, when reduced to a very fine powder, is bright red. Its crystals taste decidedly acid, it is very soluble in water. It may be heated to 136° without melting. It may be dissolved in ether and treating the residue with successive portions of boiling alcohol, which on cooling deposits a part of the carminic acid and yields the remainder by

solution unless some ammonia be next added, when carmine lake is thrown down. Neutral alkaline salts turn carminic acid to violet, while the acid salts of alkalis (bitartrate of potash, for example) render the shade more of an orange.

The chemical history of the carminates is, however, incomplete. The alkaline carminates are soluble, the others, as far as has been ascertained, are amorphous substances. The different results obtained with cochineal

1483

For further particulars see Carmine

1489

TRADE IN COCHINEAL

The Madras Government exported in September 1797, 21,744lb. From the reports of the sales of Indian Cochineal during the years 1797-1807, 55,196lb were sold at an average of 1s 6d the more than the prime cost. The price fell in 1807 that during the past seven years in England, but that from the London Company's article of profit to the Company. The propriety of discontinuing the purchase or reducing the price to be paid to the producers. The home authorities, with the view of still further fostering the industry, directed the

C. 1489

The Lac Insect.

COCCUS
lacca.

I490

trade been destroyed by aniline that a large quantity of lac-dye was recently thrown into the Thames as worthless and unsaleable (For the trade in lac-dye see a further page)

Coccus lacca, Kerr

I491

THE LAC INSECT, *Ing*, LAQUE, *Fr*, LACK, *Germ*, LACCA, *It*
Vern —*Lakh* HIND, *Guld*, BENG, *Laksha* SANS

India and occurs
especially Butea
a complete list of

I492

DESCRIPTION AND MODE OF GROWTH —Lac is the resinous incrustation formed on the bark of the twigs through the action of the lac insect. When the larvæ or grubs of the *Coccus lacca* escape from their eggs they crawl about in search of fresh sappy twigs. When satisfied, they become fixed and form a sort of cocoon by excreting a resinous substance. The male cocoon is oval in shape, the female circular. For about 2½ months the insects remain within their cocoons in the lethargic state but

it at once commences to crawl over the females. The impregnated female after depositing her eggs below her body, commences to construct cells round each with as much precision as the bee forms its comb.

the resinous excretion—lac—which it encrusts around itself. As time advances further changes are visible, the body of the female enlarges considerably and becomes brilliantly coloured. The red colour is due to the formation of a substance intended as food for the offspring. The eggs germinate below, and the larvæ, eating their way through the body of the mother, make their escape to repeat this strange history.

C. I492

COCCUS
lacca

Trees on which the Lac Insect feeds

1493

TREES ON WHICH THE LAC INSECT IS REPORTED TO FEED

- 1 *Acacia arabica*, Willd (LEGUMINOSÆ) The *Bibul* or *Kikar* (*Gamble*, 151) 'In Sind and Guzerat yields large quantities of lac'
- 2 *Acacia Catechu*, Willd. (LEGUMINOSÆ)
- 3 *Albizzia lucida*, Benth (LEGUMINOSÆ) *Silkori*, Beng
- 4 *Alenrites moluccana*, Willd (EUPHORBIACEÆ) The *Akrot* of the plains, introduced from Malay, now almost wild, especially in South India
- 5 *Anona squamosa*, Linn (ANONACEÆ) The *Ata*, a tree introduced from the West Indies
- 6 *Butea frondosa*, Roxb (LEGUMINOSÆ) The *Dhak* or *Palas*
- 7 *Butea superba*, Roxb (LEGUMINOSÆ) A climber, scarcely distinguishable from the tree *B frondosa*, except by its habit
- 8 *Carissa Carandas*, Linn (APOCYNACEÆ) Var. *spinarum*, sp, A DC
- 9 *Celtis Roxburghii*, Bedd. (URTICACEÆ) Eastern Bengal, Central and South India
- 10 *Ceratonia Siliqua*, Linn (LEGUMINOSÆ) The *Carob Tree*; now almost naturalised in the Panjab and South India.
- 11 *Croton Draco*, Schlecht (EUPHORBIACEÆ)
- 12 *Dalbergia latifolia*, Roxb (LEGUMINOSÆ)
- 13 *Dalbergia paniculata*, Roxb (LEGUMINOSÆ)
- 14 *Dichrostachys cinerea*, W & A (LEGUMINOSÆ) The *Virtul*, a shrub of Central and South India
- 15 *Dolichandrone Rheedii*, Seem (BIGNONIACEÆ). A small tree of Burma and the Andaman Islands
- 16 *Eriolana Hookeriana*, W. & A (STERCULIACEÆ)
- 17 *Erythrina indica*, Linn (LEGUMINOSÆ)
- 18 *Feroma Elephantum*, Correa (RUTACEÆ)
- 19 *Ficus bengalensis*, Linn (URTICACEÆ)
- 20 *Ficus comosa*, Roxb, in Assam
- 21 *Ficus cordifolia*, Roxb (*Gamble*, 335) Assam Lac
- 22 *Ficus elastica*, Bl The India rubber Tree (the *Bir*)
- 23 *Ficus glomerata*, Roxb
- 24 *Ficus infectoria*, Willd The *Pikar* or *Keol*
- 25 *Ficus laccifera*, Roxb (URTICACEÆ) A native of Sylhet the *Ruthal But*
- 26 *Ficus religiosa*, Linn The *Aswat* or *Pipal*.
- 27 *Garcia pinnata*, Roxb (BURSERACEÆ) The *Giruga* or *Kaikar*
- 28 *Kydia calycina*, Roxb (MALVACEÆ) A small tree the *Pola*
- 29 *Lagerstromia parviflora*, Hook f. (LYTHRACEÆ) The *Bikli* or *Sid*
- 30 *Mangifera Indica* Linn (ANACARDIACEÆ) The *Mango*, in its wild state, often yields lac
- 31 *Nephelium Litchu*, Camb (SAPINDACEÆ) The *Lichi*
- 32 *Ougeima dalbergioides*, Benth (LEGUMINOSÆ) The *Sandan*
- 33 *Prosopis spicigera*, Linn (LEGUMINOSÆ) The *Jhand* of the arid zones of the Panjab and Guzerat
- 34 *Pterocarpus Marsupium* Roxb (LEGUMINOSÆ) The *Bija* or *Lino* tree, a native of Central and South India
- 35 *Pithecolobium dulce*, Benth (LEGUMINOSÆ) The *Dikhun babli*, a tree introduced from Mexico
- 36 *Schima crenata* North (TERNSTROMIACEÆ) An evergreen tree of Burma

C. 1493

Uses of Lac.		COCCUS lacca.
37. §	issumb of the	
38. §	ease rapid	
39.	lacci-	
40. §	(Mangrove) The Tree-wood a native of	
41. §		
42. §		
43. <i>Zizyphus zizyphora</i> , Willd. (RHAMNACE). The Kat-ber.		
PROPERTIES AND USES OF LAC.		
After the larvæ escape, the old encrusted twigs are removed and cut up into pieces 4 to 6 inches long. These form <i>stick-lac</i> . They are spread upon a flat floor and a roller passed over them by which the resinous		Stick lac. 1494
		Lac-dye. 1495
		Seed-lac. 1496
		Shell-lac. 1497
		Sheet-lac. 1498
		Button-lac. 1499 D C 1500 Liver. 1501 Native Orange 1502 Garnet. 1503 Native-leaf. 1504
		Adulterated Lac 1505

COCHLOSPERMUM
Gossypium.

Lac Dye White Silk Cotton Tree.

smell on crushing the lac. The writer was once informed by a merchant that his firm in the usual course of business imported very largely resin which he believed was used up by the native dealers in adulterating the lac which they and other merchants exported. The gentleman in question condemned strongly the process of adulteration, but justly remarked that resin was an ordinary article of trade used for other purposes which if they discontinued to import would only be more largely imported by other firms.

USPS OF LAC—In India lac is dissolved in native spirits and coloured, in this form it is used as a varnish for carpentry and furniture, mixed with sulphur and some colouring agent, it is formed into the sticks *batti* like sealing wax, which are used by the toy makers to coat their wooden wares. In Europe it is largely made into *sealing wax* and dissolved in spirits, it forms *spirit varnish*. It is made into *cement* and into lithographer's ink, and is used to stiffen hats and other articles constructed of felt.

Varnish

1506

Batti

1507

Sealing-wax.

1508

Cement

1509

Dye

1510

LAC DYE

Having now indicated the main features of the lac industry collectively, the present article may be concluded by dealing in greater detail with the subject of the dye extracted from *Coccus lacca*. The reader is referred for further particulars regarding the European industry and trade in the Resin to the article **LAC**.

information regarding its use in the North West Provinces. Owing to the existence of the resinous matter mechanically mixed with the dye, lac is not so easily worked as cochineal. All the reactions and processes

1511

extent in India, the article is scarcely, if at all, exported

COCHLOSPERMUM, *Kunth ; Gen Pl, I, 124, 971.*

1512

Cochlospermum Gossypium, *DC, Fl Br Ind I, 189, Bixineæ*

SOMETIMES CALLED WHITE SILK COTTON TREE!

C. 1512

COCO or COCOA.

The White Silk-Cotton Tree.

GUM.

FIBRE.
Floss
1514Bark
1515OIL.
1516MEDICINE.
Gum
1517Floss.
1518TIMBER.
1519

R4 per maund, retail or bazaar, 3 annas per pound of the worst or black variety, wholesale, R3 per maund, retail or bazaar, 2 annas per pound "

Fibre.—The seeds possess a short but very soft and elastic floss, from which fact the plant has received its specific name. This floss is much too short to be of any service as a textile fibre, but, with the flosses of *Bombax malabaricum*, *Eriodendron anfractuosum*, and *Calotropis gigantea*, it has been classed as a "silk cotton." By some writers these have recently been designated "kapok fibres," but there is every reason to believe that the true kapok of the Dutch upholsterers is the floss of *Eriodendron anfractuosum* (see Vol I, B 641). In some parts of India the floss of this tree is collected and used for stuffing pillows, for which purpose it would seem better suited than the floss from *Bombax malabaricum*, as it is not so liable to get matted. It might be found serviceable as a gun-cotton. (Conf with C 175 and Kapok in a further volume.)

The Rev A Campbell states that the Santals prepare a good, useful cordage fibre from the bark of the tree. In the report of the Conference held on Indian fibres, at the late Colonial and Indian Exhibition, it is stated that Mr Campbell's fibres from this tree were much admired, the floss being viewed as possessing the merit of elasticity—a merit which might allow of its competing favourably with the true kapok.

Oil.—The Rev A Campbell, Santal Mission, Chutiá Nagpur, de-
 abundance
 seeds is well
 his Oil and
 and the fact
 of the

Medicine.—The gum has the properties in a mild degree of *Tragacanth*, for which it is proposed by Moodeen Sheriff and others as a substitute. It is also used as a mild demulcent in coughs. The floss has been recommended as admirably suited for padding bandages, splints, &c, being soft and cool. On this account it has been suggested as suitable for pillows and cushions used in hospitals, &c. Irvine (*Mat Med*, Patna, p. 78) says the dried leaves and flowers are used as stimulants.

Structure of the Wood.—Extremely soft, grey, but has no heart wood, and is not apparently put to any useful purpose, weight 17lb per cubic foot.

Cockles, see Molluscs (edible)

Coco or Cocoa, see *Cocos nucifera*; Coca, see *Erythroxylon*
 and *Cocoba Nibs*, see *Theobroma*

C. 1519

The Cocoa nut Palm	COCOS nucifera.
COCOS, Linn , Gen Pl, III, 945	
Cocos nucifera, Linn , Brandis, For Fl, 556, PALME	1520
THE COCOA-NUT PALM, THE COIR OF COCOA NUT FIBRE; PORCUPINE WOOD; COCOSER, Fr , CocosnuSS, KAIR, Germ	
Vern —Narel, ndriyal nar ndriyal, dab, narakel BER yal, jhada naryal, GUJ mar, naural, BOMB , Na	
.	
Dr JAVA Cocos (Kondal) - Cocos -	
OIL, COCOA NUT OIL— A	
WATER— Yelnur kapani, DUK , Yella nir, TAM , Yella niru, TEL	
TODDY— Nardli, HIND , Nardl ki-sendi, narillie, DUK , Ténga kallu, tennan- kallu, tennang kallú, TAM , Tenkaya kallu, tenkala, TEL , Nargilie, nargilli, ARAB ; Tariye-nargil, PERS	
FIBRE— Coir ? (See first paragraph of chapter on Coir), HIND ; Tennam nar, TAM , Tenkaya nar, TEL	
COCOA NUT CABBAGE— Tennam kurtu, TAM , Tenkala gurtu, TEL , Naril ka krute, ARAB	
COTTON OF TOMENTUM— Tenna maruttu pungie, TAM , Tenkeia-chettu puthie, TEL , Tennam puppha, MAL	
References—C	

COCOS
nucifera.

The Cocoa-nut Palm.

A 1012 COLUMBIENSIS, 11, 566.

Habitat.—A pinnate-leaved palm, with a straight or often gracefully curved stem, marked by annular scars; cultivated throughout tropical India and Burma, especially near the sea-coast. On the eastern and western coasts it is particularly abundant, more so towards the south. There are

Indian
Region.
1521

for example, is little more than half a mile from the beach. In very exceptional circumstances, or under the most careful garden cultivation, it

Brahmaputra, and the Malabar and Coromandel coasts. In the Brahmaputra valley it ascends to a greater distance from the sea than in the Gangetic; but in both it is an introduced tree, as it nowhere occurs in forests far away from human dwellings. On the Malabar coast, and on the islands off the coast of India, it may be different; but even in these localities it rarely exists as a forest tree, although it is self-sown. It is abundant on the Laccadive Islands, and on the Nicobar group in the Bay

The Cocoa-nut Palm.

COCOS
nucifera.

geographical and physical conditions were different from those of our day."

CULTIVATION OF THE COCOA-NUT.

CULTIVA-
TION

It is commonly reported that there are in India 480,000 acres under the cocoa-nut. A number of passages from Indian authors will be found scattered through the present account of the palm, which every now and again recur to the question of its cultivation. It may, however, be desirable to give here a brief abstract of the opinions published by the better known European writers, since from these may be gathered the results of scientific experiments.

Sowing—Ripe nuts, carefully collected, should alone be employed as seed, and for this purpose they are usually gathered from February to May. Seed from very young or very old trees should be avoided. After having been kept for a month to six weeks they should be planted

Sowing
1522

inches of their surface exposed. Ashes, or ashes and soil, should be freely placed in the trenches, these act both as a manure and as a preventative against insects. The seed-bed thus prepared should be kept moist, but not soaked. The germinated seeds may be transplanted when they are in their second to their sixth or even twelfth month. In the Godavari district they are placed in their permanent positions when three to four years old. In damp localities the transplanting may be done in the hot

planted
in the
deepTransplant-
ing
1523

In cold dry soils these pits should be lined with sand. In marshy land, walls should be constructed around them. Ashes are often recommended to be freely mixed with the prepared soil to be put into the pits, as this is supposed to prevent the attacks of the beetles that prove so destructive to the trees. Cultivation of turmeric, arrowroot, &c, in the pits, along with the cocoa-nuts is believed to be beneficial. The soil round the seedlings is also often kept damp by a bed of leaves, particularly such as will not en-

rains, the soil being replaced and levelled about the close of the rains. By the fourth year the stem begins to appear and has about 12 leaves; it is distinctly visible by the fifth year, when the tree has about 24 leaves. The spathes commence to be formed by the sixth year, and the stem is then 1 to 2 feet above the ground, but in exceptionally favourable climates

or not till the tenth year. About six months after

The Cocoa nut Palm

COCOS
nucifera.CULTIVA-
TION

of the ashes and salts of ammonia from the urine, &c, deposited day by day in the soil "

Simmonds further says "The nuts for seed should not, on being gathered, be allowed to fall to the earth, but be lowered in a basket or fastened to a rope. If let fall, the polished cover to the fibres will be injured and collect damp about the nut, or the shell inside may be cracked and the water disturbed. These are fatal injuries, or even if the plants

"Nurseries should be somewhat exposed to the influence of the sun, though not too much heat. plants thus grown will even, though deficient in stature, be strong, and when transplanted will not fail, nor suffer from heat. The planting of the nuts should take place in January to April, and also in August provided the rains are not heavy and then the planter may expect fruitful trees to be produced when grown, but nurseries formed during the heavy monsoon will generally fail, or produce trees

is recommended to be thrown into the pits when the earth is being returned around the plants. Half sand half earth is considered the best material to fill up the pits with "

PECULIARITIES OF INDIAN CULTIVATION

The following passages from the Gazetteers will be found instructive and of value to intending cultivators as having a special bearing on India

1 In Bombay (Kolab District) — Of the liquor yielding trees of this dis

1
Bombay
1529

plant the ground is hollowed 3 or 4 inches deep, and during the dry

COCOS
nucifera.

The Cocoa-nut Palm.

CULTIVA-
TION.

in the garden is set apart for growing seed-nuts. The nuts take from

If the nuts are left to drop from the tree, which is the usual practice in Bassein, they are either kept in the house for some time and then left to sprout in a well, or they are buried immediately after they have fallen. When the nuts are ready for planting they are buried either entirely or from one half to two thirds in sweet land, generally from 1 to 2 feet apart, and sometimes as close as 9 inches. A little grass, rice-straw, or dry plantain leaves are spread over the nuts to shade them. If white-ants get at the nuts the grass is taken away, and some salt or saltish mud mixed with wood ashes and a second layer of earth is laid over the nuts. Nuts are sometimes planted as late as August (*Shrāvan*), but the regular season is from March to May (*Chaitra* and *Voishākli*), when, unless the ground is damp and their inner moisture is enough for their nourishment, the nuts want watering every second or third day until rain falls. The nuts begin to sprout from four to six months after they are planted, and

the ditch round the tree, 22 pounds (4 *dhālis*) of powdered dry hsi

1530

mixture of cow-dung and wood-ashes covered with earth; or night-soil, which on the whole is the best manure. Palms suffer from an insect named *bhonga* which gnaws the roots of the tree, and from the large black carpenter-bee which bores the spikes of its half-opened leaves. When a palm is suffering from the attacks of the *bhonga*, a dark red juice oozes from the trunk. When this is noticed, a hole 3 inches square is cut in the trunk from 4 to 6 feet above where the juice is coming out, and is filled with salt, which drives away or kills the insect. To get rid of the boring bee, it is either drawn out by the hand, or it is killed by pouring into the spike assafoetida water or salt-water.

C. 1530

The Cocoa nut Palm

COCOS
nuciferaCULTIVA-
TION

"When the tree begins to yield, a sprout comes out called *por* or *pogi* at the bottom of which is a strong web like substance called *pisundi*. After about a fortnight the tree flowers, though few blossoms come to perfection. Many of the young nuts also fall off, and only a few reach maturity. A young nut is called *bonda*, a nut with a newly formed kernel is called *slale*, and a fully-formed nut *ndrel*. A good tree yields three or four times a year, the average number of nuts being about seventy-five (*Gas*, XIII, I, 295).

In the report of the Kathiawar District (*Bomb Gas*, VIII, p 95), there occurs a short but interesting account of the cocoa nut. At Ma

feet in diameter is cut in the rock and filled with mould. "All the trees" at

II
Madras
1531

nearing the Madras Presidency from Bombay it becomes more and more plentiful. Of its abundance on the Malabar coast an opinion may be

that there are 80000 acres under the cocoa nut. Indeed, the Malabar coast and the Laccadive and Maldive Islands are pre-eminently the seats of the Indian cocoa nut industry. The enquirer after Indian cocoa-nuts, for, or cocoa nut oil need practically concern himself with no other part of the country unless he add to these the Nicobar Islands. The last-mentioned islands furnish a very large number of cocoa-nuts, but apparently the islanders are ignorant of making coir or expressing the oil.

ports from these islands are treated as if they were produce of the main-

C. 1531

COCOS
nucifera

The Cocoa-nut Pa'm

CULTIVA-
TION

land, while the imports from the Maldives are returned as from foreign territory. Last year the Maldives the Nicobar Islands 4 510,000 it is not reported that they man only a small amount of copra below that which prevails on the mainland of India

1532

Impersal Gazetteer as "possessing no important trade by sea or land" It seems impossible to believe that all the coir returned under the name of "Cochin Coir" could therefore come from Cochin. Indeed the ex-

Cochin by sea amounted to only 689 cwt, valued at R4,134, and manufactured coir 2,777 cwt, valued at R25,339 these were all sent to Bengal or Bombay, how much may have gone by land to Madras cannot be dis-

1533

hin coir. pages, to the from these ref notice of tion given r six ally

bear fruit about the ninth year after transplantation. The expenses of cultivation are stated to be R668 for a *putti* of land,—namely, R140, being the price of 600 young plants, R48 being the value of the labour required for planting them, and R480 being the wages of labourers employed to water and tend the trees until they come into bearing. When the trees begin to bear fruit, the value of the produce of a tree, exclusive of the fibre, is estimated at about 12 annas a year, making the total value of the produce in a *putti* of land R300 (*p. 70*)

III
Mysore
1534

III *In Mysore* "there are four varieties of the cocoa nut 1st, red, 2d, red mixed with green, 3rd, light green, and 4th, dark green. These varieties are permanent, but although the red is reckoned somewhat better than the others, they are commonly sold promiscuously. Their produce is nearly the same

"The soil does not answer in the Bangalore district unless water can be had on digging into it to the depth of 3 or 4 cubits, and in such situations a light sandy soil is the best. The black clay, called *ere*, is the next best soil. The worst is the red clay, called *kebbe*, but with proper cultivation all the three soils answer tolerably well

"The manner of forming a new cocoa nut garden is as follows. The nuts intended for seed must be allowed to ripen until they fall from the tree, and must then be dried in the open air for a month without having the husk removed. A plot for a nursery is then dug to the depth of 2 feet, and the soil is allowed to dry three days. On the *Ugadi* feast (in March) remove 1 foot of earth from the nursery and cover the surface of the plot with 8 inches of sand. On this, place the nuts close to each other, with the end containing the eye uppermost. Cover them with 3 inches of sand and 2 of earth. If the supply of water be from

C. 1534

The Cocoa-nut Palm.

COCOS
nucifera.CULTIVA-
TION

a well, the plot must once a day be watered, but if a more copious supply can be had from a reservoir, one watering in the three days is sufficient. In three months the seedlings are fit for being transplanted. By this time the garden must have been enclosed, and hoed to the depth of 2 feet. Holes are then dug for the reception of the seedlings at 20 feet distance from each other in all directions, for when planted nearer they do not thrive. The holes are 2 feet deep and a cubit wide. At the bottom is put sand 7 inches deep, and on this is placed the nut with the young tree adhering to it. Sand is now put in until it rises 2 inches above the nut, and then the hole is filled with earth and a little dung. Every day for three years, except when it rains, the young tree must have water.

saline substances. Other soils, however, are employed, but black mould is reckoned very bad. The cocoa-nuts intended for seed are cut in the second month after the vernal equinox. A coco-nut is then dug out of the

then the young palms are fit for being transplanted. Whenever, during the two months following the vernal equinox, an occasional shower gives an opportunity by softening the soil, the garden must be ploughed five times. All the next month it is allowed to rest. In the month following the summer solstice, the ground must again be ploughed twice, and next month, at the distance of 48 cubits in every direction, there must be dug pits a cubit wide and as much deep. In the bottom of each a little previously well watered each pit. The shell be filled with earth so ng. For three months

the young plants must be watered every other day, afterwards every fourth day, until they are four years old, except when there is rain. Afterwards they require no water.

at any rate be ploughed, as the manure must be given, and as no rent is paid for the grain. On this kind of ground the cocoa-nut palm begins

COCOS
nucifera.

The Cocoa-nut Palm.

CULTIVA-
TION.

to bear in twelve or thirteen years, and continues in perfection about sixty years. It dies altogether after bearing for about a hundred years. They are always allowed to die, and when they begin to decay a young one is planted near the old one to supply its place.

"In this country, wine is never extracted from this palm, for that operation destroys the fruit, and these when ripe are considered as the valuable part of the produce. A few green nuts are cut in the hot season, on account of the refreshing juice which they then contain, and to make coir rope; but this also is thought to injure the crop. The coir made from the ripe nuts is very bad, and their husks are commonly burned for fuel.

"The crop begins in the second month after the summer solstice, and continues four months. A bunch is known to be ripe when a nut falls
 six bunches,
 om 60 to 70
 huts, raised
 id is removed

at his expense, by a man who fixes an iron rod in the ground, and forces its upper end, which is sharp, through the fibres, by which means the

IV
Nicobar
Islands.
1535

IV *On the Nicobar Islands* the cocoa-nut palm is very abundant, although, as already stated, it exists only under recent cultivation on the Andaman Islands, but reappears still further to the north on the group of the Cocos Islands. Sir W. W. Hunter gives an interesting account of the Nicobar trade in cocoa-nuts which may be here quoted. "At present the principal product of these islands is the cocoa-nut palm, and its ripe nuts form the chief export." "The northern islands are said to yield annually 10 million cocoa-nuts, of which about half are exported. The estimated number exported in 1881-82 was 4,570,000. As this important product is six times cheaper here than on the coast of Bengal or in the Straits of Malacca, the number of English and Malay vessels that come to the Nicobars is every year increasing." "The trade in cocoa-nuts is carried on chiefly by native craft from Burma, the Straits Settlements, Ceylon, &c. Forty vessels of an aggregate tonnage of 6,270 tons visited the islands for cocoa-nuts in 1881-82." "The Administration Report for 1885-86 gives the exports as 4,510,000 nuts and 5,730 bags of copra. In that year 49 vessels, with an aggregate tonnage of 8,218 tons, obtained permission to trade with the Nicobar Islands for cocoa-nuts, &c. The same report states that there are now 112,000 cocoa-nut palms under cultivation at Port Blair.

V.
Burma.
1536

V. *Of Burma* it is reported that the cocoa-nut is "largely cultivated, and might be much more so in many places along the Arakan coast as it is in Ceylon, and as doubtless it would be but for the sparseness of population, the difficulties of approaching the coast except at a few spots, and the absence of the means of land communication between the ports and the sites fitted for the production of the trees." In the Bassein district of Pegu it

VI
Bengal.
1537

and the 24 Parganas.

The Cocoa-nut Palm

COCOS
nucifera.

VII. In Upper India the cocoa-nut is alluded to in many works, but

CULTIVA-
TION

VII
Upper India.
1538

forth no branches to face its violence, the cocoa, on the contrary, loves

tropical zone "

VIII Ceylon —Speaking of Ceylon cultivation Mr Treloar says — "The ripe nuts are first planted in a nursery, where they are covered an inch deep with sand and sea-weed or soft mud from the beach, and watered daily til they germinate. In two or three months a white shoot containing the foliaceous rudiments springs from one of the three holes in the end of the nut, the radicals emerging from the other two orifices opposite to the shoot, and penetrate the ground." This is not quite a correct description of the germination. The leaf-stalk of the cotyledon elongates and pushes the embryo bodily out of the seed. The blade of the cotyledon remains within the nut forming a sort of arm of attachment. The lower point of the projected embryo elongates and forms the roots, and from a slit in the cotyledonar sheath the plumule or stem makes its appearance. The "three holes" on the nut are all close together, not "opposite" as in the above description and are only spots not holes. But Mr. Treloar pro-

VIII
Ceylon
1539

ENEMIES TO THE COCOA-NUT.

It is commonly stated that if the soil be too rich a large grub with a reddish-brown head soon finds its way to the roots and into the stem. This eats its way through the tissues until the leaves turn yellow, the terminal bud withers, and the tree is killed. This appears to be the beetle known as *Butocera rubus*. "In the Straits of Malacca, the chief natural enemy of the tree is a species of elephant-beetle, which begins by nibbling the leaves into the shape of a fan, it then perforates the central pithy fibre, so that the leaf snaps off, and lastly, it descends into the folds of the upper shoot, where it bores itself a nest, and, if not speedily extracted or killed, soon destroys the tree. A similar kind of beetle is known on the Coromandel coast, and is extracted by means of a long iron needle or probe, having a barb like that of a fish-hook. By using this and by pouring salt or brine on the top of the tree, so as to

1540

1541

1542

COCOS
nucifera

The Cocoa-nut Palm: Coir Fibre.

CULTIVA-
TION

1543

more formidable is the *cooroominyo* beetle (*Butocera rubus*), which waits to pierce the tender trunk near the ground, and to deposit its eggs in the cavity whence the young grubs, directly they are hatched, begin to eat their way up through the centre of the tree to the young leaf-ends at the

1544

ted by
mity is

1545

e and

GUM

1546

GUM.

The stem of this well-known tree is in Tahiti said to yield gum. It forms large stalactitic masses, red-brown, translucent or transparent (*Spons' Encycl*) Cooke, in his report on Gum and Gum-resins, says that this gum was sent to the Madras Exhibition of 1855 from Travancore. No other author appears to allude to this gum however, and it therefore, seems probable that if produced it is met with only in certain localities. The writer cannot recollect ever having seen a gum adhering to the stems of the palm.

DYE

1547

DYE

"In a patent obtained by Mr. J. H. Baker (No 5139, March 29th 1825) the whole or every part of this tree is claimed as a dye-ware, especially the husk enclosing the fruit, and the foot-stalks of the leaves. The dye was to be extracted by water, cold or boiling, or by solutions of lime, potash, ammonia, &c., and was to serve for dyeing nankeens, blue-blacks, &c. The infusion was likewise to serve as a substitute for nut-galls in Turkey-red dyeing. The material does not appear ever to have come into practical use." (*Crookes*.)

1548

Mr. Liotard says of this dye property - "Produces a dirty-brown (*khaki*) colour, and is a good deal used from its abundance. Lime and *chaula* are added as mordants." Drury remarks that "the shell when burnt yields a black print which in fine powder and mixed with *chunam* is used for colouring walls of houses." Cocoa-nut oil is frequently employed in certain processes sulphate of iron colour to silk pose of the coc special properties that assist the tinctorial actions

1549

The natives of India generally do not seem to be aware of the dye properties. The milk is, however, said to be used by plasterers both in India and Ceylon for a wash that has mixed with lime or colour washes it sh. For this purpose cements (see No. 1625

1550

COIR FIBRE

1551

COIR FIBRE.

The thick pericarp or outer wall of the fruit yields the valuable coir FIBRE of commerce. The SHEATHS of the leaves are used to wrap up articles, and as paper to write upon. At the Colonial and Indian Exhibi-

Leaf-Stalks.

1552

Tomentum

1553

Coir

1554

C. 1554

The Cocoa-nut Palm Coir Fibre

COCOS
nucifera.

COIR FIBRE.

of this fibre is said to come from the Malayalam *layar* (from the verb *kayiru*, to twist) through the Portuguese corruption *coiro*. "The word

and Matting Co., Highworth, Wilts., and Messrs. W. I. Sly and T. Wilson of Lancaster, who were the patentees of improved machinery for making

quality of the fibre,—soil climate, and proximity to the sea being important influences. But there are other considerations. Certain varieties or cultivated forms of the cocoa nut are better suited than others for the

accurate system of steeping, beating, and cleaning the fibre, completes the manipulation calculated to produce the superior qualities of coir (*Conf. with Mr Jackson's report in next para*). "The fibre appears in the market in various degrees of fineness, depending on the age at which the cocoa nut was cut and husked, and the care bestowed in steeping and cleaning" Mr. Treloar says. "The usual indications are that the commoner and coarser fibre comes from the old nuts, and the finer, lighter quality from the new, but there are, of course, essential differences in the qualities brought from each locality, and the Cochin are usually the best." "Here let it be parenthetically but emphatically remarked that any attempt to give to cocoa-nut fibre a finer hue by the process of bleaching is to destroy its quality if it be good, and if it be of common quality to make it almost worthless."

Properties of the Fibre and Season when Mature—"The Cochin has the purest hue and fetches the best price." On this account it has been customary to imitate this by bleaching. "Cocoa nut fibre is tough, elastic,

PROPERTIES
OF
COIR
1555

of the sea, but it will not stand bleaching. It gives up when confronted with sulphuric acid, chloride of tin, or any other chemicals which are

COCOS
nucifera.

The Cocoa-nut Palm Coir Fibre.

PROPERTIES
OF
COIR.

If cut earlier than this, the fibre is weak, if later it becomes coarse and hard, requires a longer soaking, and is more difficult to manufacture” Dr Buchanan Hamilton in his journey across Mysore states (I, 155) the green cocoa nuts are sold for their husks, from which fibre is extracted, but the husks of the ripe cocoa-nuts are commonly burnt for fuel (II, 50) At the same time immense quantities of apparently ripe cocoa nuts, in husk are sent to Europe, the coir from the husk being there separated, cleaned, and manufactured Mr Jackson of Ken, in the *Planters' Gazette*, describing a visit to Messrs. Chubb, Round & Co's factory, gives an interesting account of the process of husking there pursued He says “The enormous heap of husks—which, indeed is known in the locality

£
t
£
£
Honduras), all round the coast of America and the Fiji Island Nearly all the nuts are imported in the husk or outer covering, from which, on arrival, they are stripped by men using two fine-pointed steel chisel, and who, by constant practice, become so skilful in the art that many are able to open 1,000 to 1,200 nuts per day The nuts themselves after being removed from the husks are generally sold to wholesale fruit dealers, who, in turn, supply the the above passage Mr new ideas India is not furnish cocoa nuts to Eng cocoa nuts is actually use England attained a vast apparently having been kept for years on the nut These facts open up a new field of trade of which with a little assistance the Nicobar and Laccadive Islands might profitably and without fear of any rival hope to enjoy a large share

SEPARATION
OF COIR.
1556

Separation of Coir in India.—“The removal of the fibre from the shell is effected by forcing the nut upon a pointed implement stuck into the ground, in this way a man can clean 1,000 nuts a day The fibrous husks are next submitted to a soaking, which is variously conducted In some places they are placed in pits of salt or brackish water, for 6 to 18 months, in other places fresh water is used, but it becomes foul and injures the colour of the fibre. The chief point to be considered is the duration of the soaking, if it be continued too long, the fibre will be weakened, if it be curtailed, the subsequent extraction and cleansing of the fibre will be rendered more difficult The most approved plan of conducting the soaking is in tanks of stone, brick, iron, or wood, steam is admitted to warm the water By this means the operation is rendered very much shorter,

The Cocoa nut Palm Coir Fibre.

COCOS
nuciferaSEPARATION
OF COIR.

cellular substance is separated from the fibrous portion. When quite clean it is arranged into a loose roving preparatory to being twisted, which is done between the palms of the hands in a very ingenious way, so as to produce a yarn of two strands at once."

"As the husk gets hard and woody if the fruit is allowed to become quite ripe, the proper time for cutting it is about the tenth month. If cut before this, the coir is weak, if later, it becomes coarse and hard, and more difficult to twist, and requires to be longer in the soaking pit, and thus becomes darker in colour. When cut, the husk is severed from the nut and thrown into soaking pits. These, in some of the islands, are merely holes in the sand, just within the influence of the salt water. Here they lie buried for a year, and are kept down by heaps of stones thrown over them to protect them from the ripple. In others, the soaking pits are fresh water tanks behind the crest of coral. In these, the water, not being changed, becomes foul and dark coloured, which affects the colour of the coir. When thoroughly soaked, the fibrous parts are

"taken out of the pits too
ies, if left in too long, the
with that soaked in fresh

In the Maldives (neigh-
vornor of Ceylon) cocoa-

nuts are very plentiful, and enormous quantities of both the nut and the fibre are exported to India and Ceylon. (See the further paragraph on trade in nuts.)

From what has been said in an early paragraph regarding the cultiva-

som-shoots for the manufacture of jaggery during the first two years of its production after which it may be discontinued." In the Konkan the opinion is held that "if tapped the trees become unproductive much sooner."

The B... .. of the fibre

successfully prosecuted in the reclamation of salt-impregnated land, where

COCOS
nucifera.

The Cocoa nut Palm Coir Fibre.

nothing else would thrive (*Gen Admin Report, p 95*) A curious fact in regard to cocoa nuts grown on salt marshes is conveyed by the

beetle, and are found to bear much sooner than those planted in a sandy soil" (*p 182-83*)

TRADE IN
COIR
1557INTERESTING FACTS CONNECTED WITH THE TRADE IN INDIAN COIR
(*Conf with p 435*)

Although as suggested, the better class fibre is most likely not produced where tapping for the juice is practised, still it should not be for-

from

style-

ready

Cochin

coir is

a mere commercial term for all good coir wherever obtained In the Indian regions alluded to above, cocoa nut cultivation is prosecuted to a considerable extent Of Cochin (Madras), it may be said, coir is perhaps the most important article of export from that Native State, but Dr Shortt (*in his Monograph*)

mentions Cochin coir He Laccadives, Amindivi, Ka

the passage quoted above

Round & Co do not, it would seem, use any Cochin fibre but prefer a husk

In

indust

averag

1860-

of the coir

own The

ars ending

five years

This idea is borne out by the statement made by Royle that "the Laccadive Islands are famed for the good quality of the coir which is made there and exported to the Malabar coast" Again, speaking of the peculiar form of the palm grown in the Island of Kiltan, Royle observes "It requires no attention and comes into bearing early The tree is not so

quite ripe How far the exports of coir from the Malabar coast to

northern Laccad-

Southern by

Mal Gazetteer

users at fixed

reference can

stitutes the re
respectively

C. 1557

The Cocoa nut Palm Coir Fibre.

COCOS
nucifera.

Government on account of the islands which he manages. No change has been made for many years in the price which is given by Government for the coir produced in the islands attached to Kánara. The returns of the coasting trade of India do not specify the amounts of coir sent from the Laccadives to Malabar, so that the somewhat interesting subject of how far the juice-extracting industry of the coast is combined with the preparation of fibre cannot be definitely learned. The following facts are however, instructive.

IMPORTS of coir (manufactured and unmanufactured) into Madras from other Indian ports—

	Cwt	R
1884-85	14,745	95,884
1886-87	13,750	81,386

EXPORTS to other Indian ports—

	Cwt	R
1884-85	186,862	12,66,356
1886-87	178,228	7,98,255

TRADE IN
COIRImports.
1558Exports
1559

Turning to the tables that give the details of these figures, it is shown that of raw or unmanufactured coir Madras receives none from British or foreign Indian ports, so that unless the Laccadives, which (as stated

Malabar district alone that "the value of exported cocoa nut products is estimated at nearly a mill on sterling annually."

In a previous page some indication of the extent of the Nicobar trade in cocoa-nuts has been given. There does not, however, appear to be any trade in coir, although it seems possible that one of the inducements that

YIELD PER NUT OF FIBRE AND PRICE

YIELD OF
FIBRE.
1560

Mr Robinson, in his Report on the Laccadives, states that the difference in the quantity of coir produced in the islands is said to yield 6 cwt of coir nuts will fine island produce 2 lb of such yarns, of which there are 14 to a bundle, averaging about a maund of 28 lb. A

C. 1560

COCOS
nucifera

The Cocoa-nut Palm. Coir Fibre.

PRICE
1561

Mangalore candy of 560lb will thus be the produce of 5,600 nuts, and should contain about 20,000 fathoms of yarn. The actual price of coir received by the islanders is about R13 per candy. The value of the coir produce of a tree is calculated to be from 2 to 2½ annas, and that of the produce of 100 trees from R13 to 15. "The average value of the total raw produce of a tree bearing fruit would then be seven annas to half a rupee, and that of a plot of 100 trees, R45." For the nuts which they export to the Malabar coast they get from R7 to 10 per thousand, or rather 1,100, as 10 per cent is always allowed for luck in these sales. The islanders export from 300,000 to 400,000 nuts annually. The natives bring their coir to the Government for at the rate of R21-1 candy of 640lb. After 1

Since then the average price paid for a Mangalore candy of Amceendevy and Kidamut coir has been R20-2-0 (or R23 per Calicut candy of 640lb). But for the Kilian and Cheilat coirs, which are the best, an average of R20-12-7 or R23-12-0 per Calicut candy is paid. Up to A.D. 1825-26, the Bombay and Bengal Governments took almost the whole of the coir brought from these islands, and credited the Mangalore Collectorate with R25 per candy. The price has since fallen very much during the last twenty years. It has been frequently below the price paid to the islanders, and at best has never yielded above 12 to 20 per cent profit. The average imports of coir have been from 500 to 600 candies. Mr Morris, in his account of the Godavery district, Madras, gives the following brief statement regarding the production and yield of coir—

"The cocoa-nut tree yields an excellent fibre. The quantity of fibre

Dist 70)

Spon's Encyclopadia gives the London prices of coir as "Cochin—good to fine, £19 to £25 a ton, coarse, £16-10s to £19-15s. Yarn—good to fine, £26-10s to £46 a ton, medium, £21-5s to £28-10s, common, £14 to £22 10s, roping, £18 to £24."

USES OF
COIR
1562

USES OF COIR

"The fibrous husk of the cocoa-nut is not its least valuable product, and gives rise to a very large trade, both in the East and in Europe. At first it was only used in this country (England) for stuffing mattresses and cushions, but its applications have been enlarged and its value greatly increased by mechanical processes, and in a small pamphlet issued by Mr Treloar, more than twenty years ago, he stated that its natural capabilities having been brought out, coir has been found suited for the production of a variety of articles of great utility and elegance of workmanship—table-mats, fancy baskets, and bonnets, &c. Instead of being formed into rough cordage only, and mats made by

1563

the fibre is rendered into pleasing textures and used in carpeting and for sheep-folds, pheasantries, and poultry yards, church cushions and hassocks, hammocks, clothes lines, cordage of all sizes, and string for nurserymen

C. 1563

The Cocoa-nut Palm. Coir Fibre.

COCOS
nucifera.

and others, for tying up trees and other garden purposes, nosebags for horses, mats and bags for seed-crushers, oil-pressers and candle-manufacturers, are only a few of the varied purposes to which the fibrous coating of the cocoa-nut is now applied" (*Simmonds, Trop. Agri.*, 234) The uses of coir are of course so varied and extensive that it is scarcely necessary to enter upon them in greater detail than indicated in the above passage. To the natives of India it is invaluable as lasting in a damp climate. It is accordingly universally employed in tying the bamboos used in the construction of their huts.

FIBROUS SHEATHS OF THE LEAVES AND COCOA-NUT COTTON.—A brief reference has been made to these in an early part of this article. The finer ones are used as filters and sieves, but the coarser are apparently put to no purpose, although they have been proposed as suitable for paper-making. They might be used to strengthen saddlery, and even for ladies' corsets and splints. Knox says of Ceylon that "the filaments at the

USES OF
COIR.

1564

Fibrous
Sheaths.
1565

flabelliformis (B 680) This is sometimes collected and used by the natives to stop bleeding from wounds. A good sample of it was shown at the Colonial and Indian Exhibition.

CADJANS—"The leaves are plaited into mats and screens and also made into baskets, and combs are said to be made of the midrib of the leaflets in the Friendly Islands. In the Laccadive Islands mats are made of the cocoa-nut leaf. These mats are of fine quality and much esteemed when exported. In the islands they are employed for the sails of the smaller boats." "The Singa neatly, so as to make they form the usual the Europeans."

Cadjans.
1566Fronds
1567

for fuel, their midribs, tied together, are sometimes used as brooms for the decks of ships, as the fibres of the stalk are woody, brittle, and difficult to clean" (*Royle*).

COLLECTIVE TRADE IN COCOA-NUT PRODUCTS.

enters so largely into the daily life of the people, that little or nothing can be ascertained of the actual consumption. The returns of road, river, and rail traffic throw some light on this, and the coasting trade affords another means of arriving at an approximate estimate of a certain proportion, but even these returns fall far short of establishing a tangible conception of the total local consumption. Wherever the palm grows,

COCOS
nucifera.

Trade in Cocoa-nut Palm Products.

TRADE.

growth of the trade in the cocoa-nut palm it will not be necessary to go further back than the year 1850. Royle, in his *Fibrous Plants of India*,

the following statement :—

All published Imports and Exports for 1850.

	Imports.	Exports.
	<i>R</i>	<i>R</i>
Nuts	5,24,889	10,140
Kernels	8,66,120	4,31,008
Coir and rope	2,31,934	2,84,514
Oil	76,648	1,51,843
Shells	5,970	<i>Nil</i>
Cadjans	2,990	<i>Nil</i>
TOTAL	17,08,551	8,77,505

This gives a grand total of R25,86,056; that is to say, less than the foreign imports of last year. To compare with the above statement of **TOTAL TRADE**, the following table of the **FOREIGN TRADE** for 1886-87 (exclusive of all internal and inter-provincial or coasting traffic) may be given :—

Foreign Imports and Exports for 1886-87.

	Imports.	Exports.
	<i>R</i>	<i>R</i>
Nuts	5,98,203	8,462
Copra (or kernels)	11,76,799	79,836
Coir (unmanufactured)	6,839	77,391
Oil (manufactured but exclusive of ropes)	1,50,701	19,14,443
Oil	7,54,515	13,74,589
TOTAL	26,87,057	34,04,726

whereas in 1850 (removing approximately the items of coasting trade) the most striking

knopra (a commercial name for the kernels) have fallen to the past forty years. How far the returns of foreign trade can be accepted as an indication of total trade may be learned from the following statement

The Cocoa nut Palm			COCOS nucifera.
of the values of the coasting trade in cocoa nut products during the year 1886 87 —			TRADE.
Coasting Trade in	Imports	Exports	
	<i>R</i>	<i>R</i>	
Nuts	24 21 941	16 88 773	
Kernels (copra)	35 31 15	23 00 958	
Coir	12 20 749	9 27 302	
Oil	20 60 067	20 74 455	
TOTAL	9 33 872	69 91 483	

The table furnished by Royle for the trade in 1850 practically corre

illustration one item of this internal trade Bengal sent to Assam in 1883 84 cocoa nuts to the number of close upon two millions valued at Rs 69 000 In a like manner Bombay imports cocoa nut products from Madras Ceylon Zanzibar, &c and distributes doubtless a large pro

product Even when this has been done a very imperfect idea will have been obtained of the value of the tree to the people of India. The mere returns of trade cannot give a just conception of the importance of a product which like the cocoa nut to a large population may be said to be the resource of wealth as well as the food drink, and occupation

TRADE IN COIR, MANUFACTURED AND UNMANUFACTURED

In all the returns of this subject care is taken to explain that these do not include ropes—coir ropes and cords being placed under a general heading with all vegetable cords

The exports of Raw Coir are however so insignificant that a false impression is likely to be conveyed. The so-called manufactured coir, which figures extensively in the returns, appears to be largely crude

The Cocoa nut Palm - Coir Rope.

COCOS
nucifera.

COIR ROPES.

It is, however, better suited for running riggings its lightness being taken advantage of. In the *British Manufacturing Industries* (on Fibres and

across the path, some of these were made of coir.

OIL.

OIL
1570

The sliced kernel, dried at ordinary temperatures, either in the sun or artificially, contains from 30 to 50 per cent of oil. The method of extracting this oil in India is as follows. The kernel is squeezed in a press and the oil is found to rise to the surface.

The
climate
odour

in tropical
agreeable

In
soap
and for
rancid

ndles and
hen fresh,
dy when

Regions where Oil is Produced — While in the above sentences a brief abstract has been given of cocoa nut oil it is necessary to deal with this subject in greater detail. Enquiries are frequently addressed to the Government of India by merchants interested in the trade in this substance, so

cocoa nut oil industry is that written by Lieutenant H. P. Hawkes and published in 1857. Gazetteer writers have contented themselves with

COCOS
nucifera.

The Cocoa-nut Palm: Its Oil.

OIL.

treating the subject as too well known to call for any detailed description, and at most only the meagrest accounts have been given. To the merchant dealer of the oil, the oil is a commodity of the

the cocoa-nut are coir-nut, oil, and toddy, or the juice from which sugar and spirits may be prepared. We know that in Bombay the juice is largely extracted from the tree, that in Mysore the fibre is the chief preparation, and that in Madras and Travancore enormous quantities of both fibre and oil are produced.

from the same trees or even prepared by the same cultivators—certain plants or portions of the plantation being periodically set apart for these several industries. Under coir fibre it has been said that the green or unripe cocoa-nut is alone used for that purpose, while most writers seem to agree that the ripe kernel is necessary for the oil. It would be most instructive to know if cultivation had resulted in the production of certain races of cocoa nuts famous for their oil-yielding properties, just as the inhabitants of the Laccadive Islands appear to have developed a small-fruited one with a specially good fibre. In connection with commercial reports on cocoa nut oil it is generally stated that the finest qualities are obtained from 'Cochin' (Spon places Cochin after Ceylon). It

the
oil
Pre
of
gre

peculiar cocoa-nut that would seem to be inferior to the Malabar either as an oil yielding or an edible nut. The imports from the Maldives and Nicobar Islands into Madras are very unimportant as compared with those recorded against Bengal, yet Madras, and not Bengal, appears to control the commerce.

ledge of at present, or that a large proportion of the coast cocoa nuts or those of certain localities only are always or periodically set apart for oil-yielding. It may, of course, be the case that the trees are, so to speak, pruned by the removal for coir of so many green nuts from each tree, the remainder being allowed to ripen for oil purposes or as articles of diet.

This brief review, from want of definite information, may be accepted as indicating the direction that future reports might assume, but it may safely be concluded that, as with coir, so with cocoa nut oil, Madras is the

The Cocoa-nut Palm: Its Oil.	COCOS nucifera.
chief seat of the trade. Certain writers familiar only with Bengal (with	oil
<p>on the Madras Presidency.</p> <p>Mode of Preparation of the Oil —The ripe kernel is cut out of the shell in various ways, and either dried by exposure to the sun or by artificial</p> <p>parts, or split Under to dry</p> <p>them, after which they are exposed to the sun on mats, and when thoroughly dried are subjected to pressure in an oil-press." Balfour remarks: "The</p>	1571
	1572
	1573
	Khobrel.
	1574
<p>seer</p> <p>rior</p> <p>fres</p> <p>scrapings are then put in a copper vessel over a slow fire, and after boiling are squeezed; sometimes instead of boiling them the scrapings are</p>	Avel
	1575
<p>in water. The pieces are then crushed in water and the whole is again boiled over a slow fire, when the oil rises to the surface and is skimmed off." It is worthy of careful observation that practically the difference between <i>avel</i> and <i>muthel</i> oil is, that the former is made from fresh kernel instead of from copra. Dr. Shortt says: "Boiled oil is obtained by bruising the kopra or the fresh cocoa-nut, mixing it with an equal quantity of water and then boiling the mixture. As the steam evaporates the oil</p>	Muthel
	1576
	1577
	1578
<p>and is supposed, for that purpose, to be superior to oil obtained from copra. Hawkes says of the hot expression oil: "When required for edible purposes, the kernel of the fresh nut is taken, rasped and mixed with a little boiling water. This yields by pressure a milky fluid</p>	1579

COCOS nucifera.	The Cocoa-nut Palm • Its Oil.
OIL.	<p>which, on being boiled until all the water has evaporated, produces a clear edible oil. Only just sufficient water to moisten the pulp should be added, as a larger proportion prolongs the operation and deteriorates the product. When fresh prepared, this oil is comparatively free from smell, but speedily acquires an unpleasant odour; many attempts have been made to divest the oil of this smell, which renders it inapplicable for the perfumer's use, but only with partial success." Nearly every writer describes a different mode of preparing the oil obtained by the hot moist process. The reader is referred to a further page where this subject will be found to be dealt with under the head of <i>The Oil as a Medicine</i></p> <p>pose or other</p> <p>In the Jury Reports of the Madras Exhibition interesting information regarding the extraction and yield of cocoa-nut oil has been recorded. "Half a hundredweight of the dried kernel is a charge for a full-sized <i>cheeko</i>" (or country mortar-like oil-mill), "and a pair of stout well-fed bullocks will get through four such charges in a day, so that twenty mills are required to get through two tons in the twenty-four hours. The man who drives has usually a boy to assist him in taking the oil, which is got</p>
1580	<p>the kernel burn brilliantly</p> <p>non- When 1, but n of a ng to rd ng bland ed in with yields re oil broic, nock veral sists ling ls by</p> <p>In <i>Spons' Encyclopædia</i> it is stated that "Its principal fatty acid is lauro- stearic, together with oleic, palmitic, myristic, and some others of less im- portance all combined with glycerine." One of the most remarkable fea-</p>
1581	C 1581

The Cocoa-nut Palm Its Oil.

COCOS
nucifera.

tures of this oil is that it will take up a larger amount of water than any other commercial oil. This makes it eminently suitable for soap-making, and but for the smell which such soap leaves on the skin the oil would be even more extensively employed by the soap-maker than it is.

Industrial and Domestic Uses of the Oil—This oil has now for many years been largely used in the candle trade. Messrs Price & Co introduced in 1840, on the occasion of Her Majesty's marriage (when for illumination a cheap self-snuffing candle was required), a new composite candle, which was a mixture of stearic acid and cocoa-nut stearine. This was subsequently greatly improved until at one time cocoa nut oil was the chief feature of Price's patent candles. The immense improvements which have taken place in the European candle industry have to some extent lessened the demand for the oil, but it is still largely employed. "It is an excellent illuminator, in both candles and lamps, as it emits no smoke."

Of no less importance is cocoa-nut oil to the soap-maker. "It forms a hard and very white soap more soluble in salt-water than any other

OIL

CANDLES
1582SOAP.
1583

water, in combination with sulphate of soda and other substances, and yet

vegetable butter, is capable of taking up a larger percentage of water—and still forming a hard soap—than any other known fatty matter. The soap made from it, moreover, is more soluble in saline or "hard waters," even sea-water, and from this reason it has long been made into soap called *marine soap* for use on board ship." The odour which it imparts to the skin or garment washed with it will last for several hours. The odour resembles that of infants vomit. On this account it should never be added to the ingredients used in the manufacture of a toilet soap. It does not readily saponify with caustic soda leys by itself, but does so readily when mixed with tallow or palm oil.

A large amount of the soap made from this oil

1584

sively used medicinally

Prices and Yield of the Oil—Speaking of the year 1854 Hawkes

1585

January 1855 was £46 10s per ton, the average being from £46 to £48."

C. 1585

COCOS
nucifera.

The Cocoa-nut Palm: Its Oil.

OIL.

... that the "Cocoa-nut" oil is not more than the
 ... of there
 ... necessar-
 ... It may,
 however, be accepted as somewhere between 30 to 50 per cent. Hawkes
 ... No
 ... ocoa-
 ... oil-mill 180lb
 ... required to
 ... d to yield

1586

TRADE IN COCOA-NUT OIL.

Royle remarks that the imports into Great Britain of cocoa-nut oil were in 1850, 98,039 cwt., of which India furnished 85,096 cwt. Hawkes states: "The ... from the Madras Presidency for ... Of this by far the la ... dom and France, the remai ... Bombay, and the French (Indian) ports." In 1850, as in the present day, the cocoa-nut oil trade almost entirely centred in Madras, so that the above passages may be taken as approximately indicating the extent of the foreign demand for the oil forty years ago. In 1880-81 the foreign exports ... Madras alone having in addition by 886 87 the ex-ports 556,562 .. 689,087 gal-1 000,180 gal-

R20,60,067; the exports were 1,942,809 valued at K20,74,455 ... of India, one of

coastwise imports were unimportant. Local production added to these imports would constitute the supply from which the exports could be made, and in the case of Madras it is noteworthy that that presidency imported

C. 1586

The Cocoa-nut Palm • Its Oil.

COCOS
nucifera.

OIL.

Turning to
and to prevail,
allons and in
il production
n these presi-

dencies. Cocoa-nut oil is thus a speciality of Madras trade

COPRA OR DRIED KERNEL

COPRA
1587

A very imperfect idea of the supply and demand for this oil would, however, be conveyed were we to omit to examine in this place the trade in copra or dried kernel, the substance from which the oil is expressed. This is largely exported to foreign countries and sent from one province of India to another to be locally made into oil.

	1884-85		1885-86		1886-87	
	Cwt	R	Cwt	R	Cwt	R
Imports	39,653	3,95,685	105,296	10,20,841	125,222	11,76,799
Exports	64,323	5,34,291	21,755	1,86,800	9,337	79,836

The imports come chiefly from Ceylon and the Straits Settlements, and are almost exclusively delivered in Bengal and Bombay, only very small amounts being received by Madras. The exports, on the other hand, go mainly from Madras (8,135 cwt of last year's exports) Bombay being next in importance. The greater part of these exports (7,149 cwt) go to Portugal, Persia, Ceylon, Russia, and Arabia, each receiving from 300 to 500 cwt. So far for the foreign traffic. The imports and exports coastwise have now to be considered. The total imports by coasting traffic were 347,255 cwt, valued at R35,31,115, and the exports 236,250 cwt, valued at R23,00,958. Of the imports Bombay received 219,201 cwt, Bengal 62,971 cwt, Sind 34,658, Madras 27,025 cwt. Of the exports, Madras sent to other Indian ports 182,509 cwt, Bombay 53,295 cwt, Bengal exporting

OIL-CAKE OR PUVAC—Before passing from the consideration of cocoa-nut oil it is necessary to say something about the oil-cake. This is viewed as an exceedingly valuable manure, especially to cocoa-nut palms grown inland. It is also largely used to fatten fowls, pigs, cows, and other

OIL CAKE.
1588

COCOS nucifera.	The Cocoa-nut Palm as a Medicine.
	animals It is sometimes exported to Europe In Madras it sells for 3 to 4 maunds (of 25lb) per rupee
MEDICINE	MEDICINE.
Fruit	The GREEN FRUIT is given as a refrigerant, the FLOWERS as an as-
1589	tringent, and the OIL employed as a substitute for cod-liver oil The milk
Flowers.	of the nut, the juice from the FLOWERING SPIKE, and the tomentum from
1590	the LEAVES are all used medicinally
Oil.	
1591	
Spike.	
1592	
Leaves.	
1593	
Water.	
1594	scrotum,
Edible Pulp.	THE EDIBLE PULP AND THE MILK PREPARED THEREFROM—The
1595	The milk of the
	s
	l
	y
	has to for it
	,
	s,
	,
	s
	t
	,
	t
	t
	t
	t
	t
	t
	t
	t
	t
	t
	t
	t
	t
	t
	t

The Cocoa-nut Palm as a Medicine.

COCOS
nucifera.

MEDICINE

Shell-Oil.
1599

ous used medicinally, the most convincing statements have been published both as to " and made of no account. The "A very cheap, hard, white pharma- ceutical purposes, of soap liniment" (Dymond, says this oil is inferior vehicle for liniments Sakha prepared for

1600

oleine obtained by pressure refined by being treated with alkalis, and then repeatedly washed and distilled with water" The therapeutic pro- "In as a rable rption oduce

1601

1602

much used as a local fevers and debilitating vermifuge in Jamaica sugar, in flux. An

C. 1602

COCOS
nucifera.

The Cocoa-nut Palm as a Medicine.

MEDICINE

I603

and pulmonary diseases generally. Pound the kernel with water, place it to settle, and skim off the cream. This is preferable to the expressed oil."

"Cocoa nut oil was proposed by the late Dr. Theophilus Thompson (*Proceed of Royal Society, 1854, Pt. III., p. 41*) as a substitute for cod-liver oil; and in this character it has been favourably noticed by Dr. J. H. Warren (*Boston Med and Surg. Journ., Vol. III., p. 377*) and others. The substance used in these cases was not the ordinary commercial oil, but the oleine obtained by pressure from the crude oil (in the solid state it is met with in England), refined by being treated with alkalis, and then repeatedly washed with distilled water. In his Lethsman Lectures Dr. Thompson gives the result of his treatment with this agent in 53 cases of phthisis. Of the first 30, 19 were much benefited, in 5 the disease remained stationary, and in the remaining 6 the disease continued to advance. Of the second 23, 15 were materially benefited, 3 remaining stationary, and 5 became worse. Dr. Garrod (*Brit and For. Med Chir Rev, Jan 1856*) has shown that it exercises a marked influence, almost equal to cod-liver oil, in increasing the weight of the body. The great advantage of its employment experienced by Dr. Thompson, Dr. Garrod, and also by the Editor, who instituted some trials with it, is, that under its prolonged use it is apt to induce disturbance of the digestive organs and diarrhoea. Its use is favourably noticed in the Report of Drs. Van Someren and Oswald, and Mr. J. Wood." (*Pharmacopæia of India*)

Dr. Dymock says cocoa-nut oil has been tried in Europe as a substitute for cod-liver oil.

general use :

with disad-

and induce

formed by some writers regarding

fact that nearly every author describ-

and consequently that it is possible many different substances or a substance in many stages of purity or impurity may have been experimented with? In the Maldives cocoa-nut oil is esteemed a powerful antidote against the bite of poisonous reptiles.

THE JUICE.—The freshly-drawn JUICE is considered refrigerant and diuretic, and is valuable as a preparation known as toddy poultice (see below).

It constitutes one of the most powerful of the

A tumblerful of the

on account of its

(Dymock)

Inside SCRAPINGS OF THE HUSK

use and heal them rapidly if

application was proved by the

case of two bad ulcers occasioned by the bite of a negro's teeth. The young roots boiled with ginger and salt are efficacious in fevers, the same as the bamboo" (*Royle*)

THE COTTON OR TONENTUM.—"This is a soft, downy, light-brown-coloured substance, found on the outside of the lower part of the branches of the cocoa-nut tree, where they spring from the stem, and are partially covered with white hairs. The coco is covered with blood, in cases of hemorrhage, and is admirably fitted to be used with tomentum. It is used under the bark of the tree." (*Royle*)

Tomentum.
I606

C. 1606

The Cocoa-nut Palm as a Medicine.

COCOS
nucifera.

THE FLOWERS —Are sometimes used medicinally, being said to be astringent

MEDICINE.
Flowers

1607

Nuts.

1608

Roots

1609

in sore-throat

Ashes.

1601

THE ASHES —“The ASHES of the leaves contain an amount of potash, they are used medicinally,”

Bud.

1611

THE BUD —The tender buds of this palm, as also of *Borassus* and *Phoenix*, are esteemed as a nourishing, strengthening, and agreeable vegetable.

Special Opinions —§ “The husk of the fruit of the *Cocos nucifera* is used in the treatment of

of male fern when tak

W Nolan M D, Bomba

acidity and gastric irritati

ed as a local application

B A, M B, Monghyr

eczema of the scrotum

ing is a popular dome

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

give is from 20 to 30

daily An ash is prep

is a valuable ant-acid

sweet extract is also

(*Civil Surgeon R L D*)

from this palm is very refreshing and possesses laxative properties

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

(*A Civil Surgeon*) “If the flowers are mixed with sugar, the root of *Alhus-khus*, and white *chandani*, with a little water, the combination will be found good in bilious fever, will check vomiting, and produce a cooling

COCOS
nucifera

The Cocoa-nut Palm - Its Edible Products.

MEDICINE.

1612

sensation" (*Civil Surgeon William Wilson, Bogra*) Useful "in dysentery, diarrhoea, menorrhoea, and stomatitis" (*Native Surgeon T Ruthnam Moodelliar, Chingleput, Madras Presidency*) "*C. mamilaris*, dwarf cocoa-nut tree, Pemba, East Africa fruit large, smooth, distinctly three-cornered pinkish yellow when ripe without the fibrous pericarp of the common cocoa-nut Yields very little oil, but supplies a refreshing drink in fevers and in hot weather, and is said to produce free diuresis used when the nut is full grown, but before it begins to ripen Vern. of East Africa *Muosi C nucifera*, *Muusi-ya Pemba*, *C mamilaris* (*Surgeon-Major John Robb, M D, Surat, Bombay Presidency*).

FOOD

FOOD PRODUCTS.

Under the head of food products obtained from this palm we may

Cocoa-nut
Cabbage

1613

he tree
, obtain

Young Cocoa-
nut

1614

Young Cocoa-nut (VERN dab) — This is the tender fruit, plucked off the tree for the cooling, sweetish, clear water, and the soft, cream-like pulp it contains The water is drunk and the pulp eaten by natives of all classes

Mature
Cocoa-nut.

1615

Mature Cocoa-nut (VERN jhuna narkel) — This is the fruit in its mature state, with its outer, thick, fibrous covering completely dried It contains less water, but has a thicker and harder albuminous layer than the tender fruit, when dried this albuminous substance is known as copra. It is eaten with parched rice, or rasped and put into curries or made into sweetmeats Copra is either allowed to ripen and dry within the shell, when it separates naturally and is removed entire, or the shell is broken and the copra cut out and dried either in the sun or over fires The former exists in large pear-shaped pieces smaller than, but of the same shape as, the interior of the nut, and is known as "natural copra" The latter occurs as the irregularly-cut pieces known as "artificial copra" An oil is extracted from copra which is employed for various culinary purposes, and is also exported to a certain extent (For further parti-

Juice

1616

Root

1617

NUTS

1618

in pan

THE NUTS

The above is a brief abstract of the food products of this palm The extent to which the unripe fruit is cut, the water and unripe kernel being consumed and the husk made into coir, may be partly inferred from what has been already said regarding the fibre To a large population in

TRADE
In nuts

1619

and coasting trade in these nuts, as recorded in Mr J E Cooch's

ripe
1 to
1 dia
107

The Cocoa-nut Palm: Its Edible Products.

COCOS
nucifera.TRADE
IN NUTS

1,434,821, and East Africa 627,346. Of these imports Bengal took 8,430,229, valued at Rs. 1,75,552, Burma 5,618,949, valued at Rs. 3,72,702, Bombay and Madras each received 700,000, and Sind 86,800. Bengal exported no cocoa-nuts to foreign countries, but Bombay and Madras each sent about 150,000 to Egypt, Arabia, and Turkey in Asia. The foreign trade in ripe cocoa-nuts is therefore very unimportant, and but for the Maldives being viewed as foreign territory (while the Laccadives and Nicobar Islands are not), it would be scarcely worthy of notice. It is noteworthy that India at present takes practically no part in meeting the

Of the coastwise exports in 1886-87 Bengal sent to Burma, according to one official table of coastwise trade, 1,676,773, but according to another

into Burma alluded to above

JUICE FROM THE COCOA-NUT.

Dr. Hugh Oleghorn has described as follows the process of tapping the palm for its juice in Madras—a process which is essentially that followed in Bombay and other parts of the country. This palm is not tapped in Bengal. When the spathe is a month old, the flower-bud is considered sufficiently juicy to yield a fair return to the (Sánár)

JUICE
Madras.
1620

the cut end of the spathe to crush the flowers thereby exposed and to determine the sap to the wounded part, that the juice may flow freely. The stump is then bound up with a broad strip of fibre. This process

COCOS
nucifera

The Cocoa-nut Palm Toddy.

JUICE

is repeated morning and evening for a number of days, a thin layer

A single spathe will
 continue to yield toddy for about a month, during which time the Sanár
 mounts the tree twice a day and empties the juice into his *eroppetty* (a

about a quarter of a measure per tree. The length of time a tree

Bombay.
 1621

in 1933, and in Kandra (AV, Part I, 58, Part II, 203) to the returns the writer has had access to, there are some 3½ million trees
 in Bombay, of which about 30,000 to 40,000 are tapped for their juice

The following table shows the number of trees tapped in the various districts of the Bombay Presidency

in Malabar and Deogad 2½d (1 anna 8 pie) a month or 2s 6d (11½) a
 year on each tree tapped. Under the new system a special license is
 granted to tap trees, at a fixed rate for each tree, and under certain con-
 ditions as to the number of trees included in the license. The licensees

to distil,
 keepers
 juice of
 6d (12
 st of fuel
 the tap-

C. 1621

The Cocoa-nut Palm • Toddy.

COCOS
nucifera.

JUICE

ping tax he had paid to Government Government levies from the liquor shop keepers £60 (R600) a year for every hundred trees tapped. Three fourths of this the liquor-shop keeper pays, the remaining fourth he recovers from the Bhandári who supplies the liquor. The Bhandári's share of the tax amounts to £15 (R150) on one hundred trees for one year,—that is, a monthly charge of £1 5s (R12½) on the one hundred trees, or on each tree a monthly tax of 3d (2 annas).

In Ratnágiri the juice is obtained from each tree in (1½ seers) of juice a seldom sold raw but by him to the liquor-shop keeper. With the wages of an assistant the monthly charge for distilling the produce of one tree is about 2d (1½ annas).

Spirit

300 trees he makes a fairly good income.

Of Ratnágiri, it is said, there are ordinarily three kinds of palm spirit, known respectively as *ras*, *phal* or *dharti*, and *phen*, *ras* being the weakest and *phen* the strongest. In some places a still stronger spirit called *duvas* is manufactured. The average wholesale rates at which for the imperial gallon, 7 pie, *phal* is 1½d (8 annas) 4s 9½d (R2 6-4).

The spirits are distilled in private stills, licensed to be kept at certain Bhandáris' houses under fixed conditions as required, in proportion to the number of trees licensed to be tapped in the vicinity. One still is usually allowed for every 100 trees, and the still-pot is limited to a capacity of 20 gallons.

Rasi.
1622
Phul
1623
Pheni.
1624

FERMENTED AND UNFERMENTED BEVERAGE

TARL
1625

This is one of the forms of the so-called palm-wine so much extolled by the early European visitors to India. From what has been said in the preceding pages regarding the juice it may have been inferred that, if left for a short time after removal from the tree it rapidly ferments and becomes intoxicating. This is the *tari* or toddy (or in the case of the cocoa nut more specifically known as the *nira*), a beverage very extensively consumed in India. Fermentation is said to be prevented by the addition of a little lime to the fluid. The earthen vessels into which it drains are generally powdered with lime when the fluid is to be drunk in its fresh unfermented state, or is intended to be boiled down to sugar or jaggery. It is also drawn early in the morning instead of being left on the tree overday. Robinson says that they are still so strict in the abstinen- the manufacture of toddy would not be self-fermented toddy is extensively used of yeast. When fermented the juice may be distilled into spirits or made into vinegar. One hundred gallons of *tari* yields on an average twenty-five of *arak* by distillation.

COCOS
nucifera.

The Cocoa-nut Palm: Sugar.

PALM SUGAR

PALM SUGAR

Instead of being fermented, the liquor may be evaporated down and its sugar thus extracted "Eight gallons of sweet toddy, boiled over a slow fire, yield 2 gallons of a lusciously-sweet liquid, which is called *jaggery* or sugar-water, which quantity being again boiled, the coarse brown sugar called *jaggery* is produced. The lumps of this are separately tied up in dried banana leaves" (Royle). Dr Shortt says "The sap is poured into large pots over an oven, beneath which a strong wood-fire is kept burning, the dead fronds and other refuse of the plants being used as fuel. The sap soon assumes a dark brown semi-viscid mass, well known as

pots or pan
jaggery, it

state it is sold to *adatti* contractors, sugar renners, or melchams. The sugar refined comprises several sorts, known in the market as moist, raw, coarse, and fine sugar. The *jaggery* is placed in baskets and allowed to drain, the watery portion or molasses dropping into a pan placed below. This is repeated, so that the *jaggery* or sugar becomes comparatively white and free from molasses. This sugar—for so it may now be called—is put out to dry, and the lumps broken up, when dry it is termed raw sugar, and weighs about 25 per cent of the whole mass, the rest of it being collected in the form of molasses." Thus cocoa-nut sugar is chiefly met with in the form of *jaggery*. It is well known, however, that it is capable of being refined according to European principles, and a certain amount of cocoa-nut sugar is regularly prepared. "The success of Dr J N Fonseca (author of the *History of Goa*), in converting toddy of the cocoa-nut tree into crystallized sugar, has been hailed with satisfaction by the press at Goa, and flattering calculations are made of the advantages that will accrue to the country from the development of this new industry" (*Bombay Gazette*). A similar sugar is prepared from the date-palm, from the palmyra-palm, and from the Indian sago-palm (*Caryota urens*). The date palm is very largely used for this purpose in Bengal, and the cocoa nut and palmyra palms in Madras while in Bombay, apparently, sugar is only very occasionally made from the juices of these trees, but when extracted it is most generally prepared from the palmyra or *Caryota* palms. Some years ago the Government of Bombay, getting alarmed at the growth of the habit of toddy-drinking, brought Jessore sugar manufacturers to try the experi-

Refined.
1626

cally failed. It is not known whether or not sugar to any appreciable extent is actually prepared from the Bombay palms, nor even whether a license is necessary to tap trees for sap intended to be so used. Of the Thána district it is said "Coarse sugar or *gúl* is also made by boiling the juice in an earthen pot over a slow fire." It is worth recording that, accord-

—Bombay, of
said to
ed Of
tapped,
48,000 of these occur in Kánara, 21,672 in Kolába, and the remainder in Ratnágiri.

In a recent report on the trade in Indian sugar issued by the Revenue and Agricultural Department, no mention is made of palm sugar being

C. 1626

The Cocoa nut Palm* Sugar.

COCOS
nucifera.

prepared in Bombay, so that it may be inferred the trees licensed to be

PALM SUGAR

	Acres
Palmyra	24,900
Cocoa nut	5,700
Date	1,600
	<hr/> 32,200

The writer of that report adds "In 1884-85 and 1885-86 the area under cocoa-nut, date palms, and palmyras was 31,000 acres and 28,000 acres

ment in 1886 it was estimated that there were 7,776⁵ acres under that palm. Taking the customary estimate of 100 trees to the acre, we arrive at the conclusion that out of a total of 7,776,500 trees, 570,000 were tapped, or

There exist
sult the gr
be tapped

made with the view to the preparation of the beverage. It would be instructive to know if the 5,700 acres of cocoa nuts in the above statement of 1886 are all under cultivation, or if some are in the wild state.

re are
sugar

nown*
cocoa

on the
as to
sugar

making, we went fully into the matter, receiving considerable assistance from Mr. D. O. Amesekere, a proctor who, when we last heard of him, was practising at Kur.

crystallized cocoa
by smoke. The
when collected w

would render the enterprise unprofitable. What pays natives on a small scale will not pay Europeans when the matter is entered into on commercial principles. An experiment might be tried, however, labour being economised by the use of ladders, perhaps, and a larger use than the natives make in toddy drawing, of safe passages from tree to tree. (*Tropical Agriculturist*, 1881-82: 563)

COCOS
nucifera

The Cocoa nut Palm Spirit.

CEMENT
1627

CEMENT MADE OF LIME AND COCOA-NUT JAGGERY

makes excellent cement" Drury remarks: "This jaggery is mixed with
heat and
bricklayers
urest castor
which the

seeds are boiled"

In *Spons' Encyclopædia* there occurs the following regarding Ceylon
"Amongst the most important articles which it is put is that of
from burnt coral or
receiving so beauti-
stinguished from the

THIS subject appears to be well worthy of chemical investigation, for
there seems every reason to presume that the property of this ingredient in
combination with lime might, with great advantage, be employed to replace
the whitewashes commonly used, to the injury of the garments of who-
ever may lean against walls so coloured (*Conf with opening sentences
under Domestic Uses, and the account given under Dye, C. 1547*)

SPIRIT
1628

PALM SPIRIT OR ARAK

Instead of being consumed as a fermented beverage the palm wine
separate record
rest satisfied
o be tapped
the method of

taxation and process of distillation generally pursued The present notice
of cocoa nut spirit may therefore be concluded by the following note
kindly furnished for this work —

Dr Lyon, of Bombay, has recorded some interesting details regard-
ing the alcoholic strength of toddy from the cocoa-nut, date and brab In
the following table is shown the average alcoholic strength of six night
collected samples of each of the three kinds of toddy at respectively three
and eight hours after collection and the average maximum alcoholic
strength attained by the samples, as well as the strength of samples
collected during the twelve day-hours, when examined the morning after
collection —

	PROOF SPIRIT PER CENT		
	Cocoa nut	Date palm	Brab (Borassus)
<i>Night samples</i>			
3 hours after collect on	7 15	5 8	3 9
8 " " "	10 0	8 0	4 7
Maximum strength	11 9	11 0	7 9
<i>Day samples</i>			
15 hours after collect on	10 8	11 7	6 5

The Cocoa-nut Palm. Spirit.

COCOS
nucifera.

SPIRIT.

"Dr. Lyon finds that in toddy collected in pots which have previously been used, fermentation commences before the pots are removed from the tree. The toddy possesses the following properties:—

VINEGAR
1629

Vinegar from Palm Wine.—Nearly every writer who has dealt with the subject of the useful products of the cocoa-nut alludes to the vinegar prepared from the juice. "One hundred gallons of toddy produce by distillation, it is said, twenty-five of *arak*. Or it may be allowed to undergo the acetous fermentation and produce very good vinegar. Or instead of being allowed to ferment, the toddy may be made to yield *jaggery* or sugar. For this purpose a supply of sweet toddy is procured mornings and evenings, particular care being taken that the vessels employed have been well cleaned and dried" (*Royle, Fib Pl*)

The vinegars prepared from the juice of the various palms that yield such juices do not appear to have been carefully examined. The natives of India attribute peculiar properties to each.

STRUCTURE OF THE WOOD.

TIMBER
1630

possesses great elasticity, and is for this reason particularly well adapted for temporary stockades which are exposed to cannon-shot." (*Drury.*)

DOMESTIC SACRED USES

DOMESTIC
1631

Hukah
Bowls
1632
Ornamental
Objects
1633
Spoons.
1634
Sugar-pots
1635
Tea-pots
1636

a graphic account of the manner in which the cocoa-nut enters into the every-day life of the people of the tropics:—

Dickens in *Household Words* says: "To a native of Ceylon the

C. 1636

COCOS
nucifera

The Cocoa-nut Palm Domestic Appliances

DOMESTIC

cocoa nut palm calls up a wide range of ideas, it associates itself with nearly every want and convenience of his life. It might tempt him to assert that if he were placed upon the earth with nothing else whatever to minister to his necessities than the cocoa nut tree he could pass his existence in happiness and content. When the Cingalese villager has felled one of these trees after it has ceased bearing (say in its seventeenth year) with its trunk he builds his hut and his bullock stall, which he thatches with its leaves. His bolts and bars are slips of the bark by which he also suspends the small shelf which holds the stock of home-made utensils and vessels. He fences his little plot of chillies tobacco and fine grain with the leaf stalks. The infant is swung to sleep in a rude

is it a yoke or pangs formed of a cocoa nut stalk. When he sits, he drinks of the fresh juice of the young nut, when he is hungry he eats its soft kernel. If he has a drink of arrack, he

cocoa nut oil, which softens it with cocoa nut

chars, the tree jars his Over his cocoa nut course, a

accordance with fact. It is however a true picture of the all important of the Prince of Palms to the inhabitants of the tropical regions.

In order to convey some idea of the numerous uses of the cocoa nut palm the following extract from the Colonial and Indian Exhibition Catalogue may be here reproduced. It is a list of certain articles prepared

(1) state &c y the over

the metallic one of not being corroded

(3) Drainer (*Zara*)—Used for draining food fried in *ghí* (clarified butter) or oil

(4) Ladle (*Doho*)—Used for water

(5) Ladle small (*Buddi*)—Used by natives for taking out oil for daily use from an earthen vessel containing the yearly or quarterly stock. It is not corroded by the oil.

(6) Hubble bubble (*Gudgudi*)—This is the *hukah* of the poorer classes

(7) Beads (*Mani*)

(8) Vinegar (*Sirka Amti*)—Made of the juice (*toddy*) of the cocoa nut palm

COCOS
nucifera

The Cocoa-nut Palm • Domestic Appliances.

DOMESTIC

- (37) Toy parrot in cage (*Pinjaryát Popat*) — Made by children from the leaf
- (38) Leaf
- (39) Root
- (40) Rope (*Kathá, Sumbha*) — This is extensively used
- (41) Oil-bottle (*Doula*). — Hung beneath the labour-cart with castor oil and brush in it for lubricating axles
- (42) Nut, immature (*Ahakota*) — Used medicinally as an astringent, children are fond of it
- (43) Trough (*Panshira*) — Trough made of cocoa-nut tree, used for catching water drawn from a well with a Persian wheel for irrigation purposes (model)
- (44) Conduit (*Panhál*) — A conduit put under the hole of the trough for conveying water for irrigation purposes
- (45) Adapter (*Nala*) — Piece of the adapter used for connecting the native still to the condenser
- (46) (*Tuntuna*) — Native musical instrument, used by the poorer classes
- (47) Beam (*Bahal*) — Piece of beam of the shape used for houses. It is also used for fishing-stakes in the sea, generally two cocoa nut trees make a stake 60 to 70 feet long.
- (48) Rosary box — Made of immature cocoa nuts
- (49) Charcoal Powder (*Kolsá*) — Burnt shell used for preparing black and lead coloured washes for houses
- (50) Broom (*Zadu*) — Made of the ribs of the leaf, used by the Bombay and other municipalities for sweeping roads, streets, yards, &c
- (51) Broom (*Zaddú*) — Made of the stems of the blossom and nuts, used by the cultivators for collecting dry leaves for (*rab*) burning on the fields
- (52) Cask
- (53)
- (54)
- (55) the gosavies (a class of professional beggars)
- (56) Sling (*Shinka*) — Used for keeping sundry articles of food out of the
- (57) F
- (58) Tar with acetic acid (*Kartel*) — Made by burning the shells in a
- (59)
- (60) washing baskets and rice drainers (*Shisbum*)
- (61) Sugar, molasses (*Gúl*) — Made of the juice (*toddy*) in Goa
- (62) (*Band*) — Peeled from the outer part of the stem of the leaf. Is used as a cord by the *toddy* drawers
- (63) Cocoa nut gilded (*Karyacha Narel*) — Offered by the higher classes of Hindus to appease the sea on the cocoa-nut fair day. At weddings the bridegroom and bride carry it in their hands

CODONOPSIS
ovata.

- (64) Husk (*Sāl, Chavād, Sodan*) —Used as fuel. Especially for backing purposes also affords coir fibre
- (65) Scoops.—Made of the shell The round and deep ones are used as drinking cups
- (66) Neck belts (*Pattā*).—Used for yoking bullocks and buffaloes to carts, ploughs, oil-mills, &c
- (67) Sack (*Thaṣṭi Gālī*) —Used for sending out articles; a somewhat similar one is attached to the cart for carrying straw or grass
- (68) Tooth-brushes (*Daton*) —The pedicels of the blossom are used as tooth-brushes
- (69) " " " " " " " " " " " "
- (70) " " " " " " " " " " " "
- (71) " " " " " " " " " " " "
- (72) Soap (*Sabu*) —Made of cocoa-nut oil, has larger percentage of water than any other soap
- (73) Puzzles and toys.—Rings, whips, neckties, rattles, crosses, &c.
- (74) Bats for cricket —Made of the wood (cocoa-nut)
- (75) Oil-cakes (*Pend*) —Oil cake from the native mill
- (76) Patimar (ship) (*Fatemarf*) —Toy made by the boys of the fishermen class
- (77) Boat, fishing (*Hoḍke*) —Toy made by the boys of the fishermen class.
- (78) Kernel (*Khobre*) —Dry kernel
- (79) Stem (*Fhsutar*) —Used as broom
- (80) Charpai, Cot (*Khat, Bāj*) —Used by the natives (model).
- (81) Potash (crude) (*Khar*) —The ash of the stem of the leaves, they produce 20 per cent of ash
- (82) Cocoa-nut, abortive (*Vāṇṣā Nārel, Vāḥṣl*) —Used as floats for begin-
- (83) " " " " " " " " " " " "

DOMESTIC

Codilla —A commercial term for the refuse separated on cleaning hemp or flax fibres.

1637

CODONOPSIS, Wall, *Gen. Pl.*, II, 557

[160, fig. 3; CAMPANULACEÆ.

Codonopsis ovata, Benth ; Fl. Br. Ind., III., 433; Royle, III., 253.

1638

• Vern — *Lúdt*

Habitat.—A herbaceous plant common in the N W Himálaya from Kashmir to Gurhwál at altitudes from 8,000 to 12,000 feet, distributed

MEDICINE.
1630

FOOD.
1640

or cooked.

C. 1640

COFFEA
arabica

Coffee.

COFFEA, Linn.; Gen. Pl., II., 114.

[RUBIACEÆ.

1641

Coffea arabica, Linn.; Fl. Br. Ind., III., 153; Wight, Ic., t. 53;

COFFEE, Eng.; CAFÉ, Fr.; KAFFEE, Germ.

Vern.—*Bun* (the berry), *Kahwa* (the same roasted and ground). *Kawa*,
bun, *bun*, *coffee*, *cofi*, HIND
Kawa, *bun*, *kahwa*, *bund*, c
 MAR; *Bünd*, *tochem-keweh*,
cap, TAM; *Kopi-vittulu*, ca
Kaphi, *Bonda-bia*, *kápi-on*,
 ARAB; *Bun*, *qahva*, *kahwa*,
kaphi-si, BURM, *Kopi att*,
 " "

Shoa, and in Yemen as applied to the berry.

Ref. *Fl. Br. Ind.*, III., 153; *Wight, Ic.*, t. 53; *Fl.*, 276;

ings of the
 soc. Jour-
 dia, 1864,
 11er, 1869;
 idia Office
 spondence
 75; Corre-
 ssociation
 1879; D
 eports-of
 Food, 40;
 Welter-
 lackwood's
 lassal, on
 ietner, on
 re Coffee-
 n Mysore
 on, 1871);
 806), W.
 ? W Las-
 n Loerne,
 'ler, Pros-
 Laborie's

Coffee Cultivation

COFFEA
arabicaCULTIVA-
TION.

Habitat—Most authors seem to agree that the coffee plant is indigenous to Abyssinia, the Soudan, and the coasts of Guinea and Mozambique. "Perhaps in these latter localities, so far removed from the Atlantic, it is able to flourish. No one has yet found it of penetrating into the interior, but it will be hard to deny its ability of germinating, often spring up round the plantations and naturalise the species. This has occurred in Brazil and the West India Islands, where it is certain the coffee plant was never indigenous" (*De Candolle*).

It is a small, much-branched tree or bush 15 to 20 feet in height, with whitish bark and white orange like flowers. The fruit, which is red on ripening, is about the size of a small cherry, and contains two seeds, closely united. These on being separated constitute the coffee berries of commerce, and on being roasted and ground, the coffee of the shops.

1 In India *Coffea arabica*—the coffee plant—is largely cultivated, but other species are also met with.

2 *C. bengalensis* *Roxb.*, occurs from Kumāon to Mishmī, also in Bengal, Assam Sylhet, Chittagong and Tenasserim. Fruit ovoid-oblong (*Harina* in Chittagong see *Agr. Hort. Soc. Ind. Proceedings*, Oct. 1865).

3 *C. fragrans*, *Korth.*, found in Sylhet and Tenasserim. Fruit much like the two last.

4. *C. Jenkinsii*, *Hook f.* Khasi Mountains. Fruit and seeds different from the last being ellipsoid.

5 *C. khasiana*, *Hook f.* Khasi and Jaintia hills. Fruit $\frac{1}{2}$ inch in diameter, smooth, seeds ventrally concave.

6 *C. travancorensis*, *W & A.*, occurs in Travancore. Fruit broader than long.

7 *C. Wightiana*, *W & A.*, the Western Peninsula, in arid places from Coorg to Travancore. Fruit much broader than long, with a deep furrow.

With the exception of the first these species are not of any special economic importance, and very little coffee is grown in the tracts in

HISTORY OF COFFEE CULTIVATION AND OF THE HABIT
OF COFFEE-DRINKING

The regions best suited for coffee cultivation lie between 15° N. and 15° S latitudes, but it is grown as far as the 36° N. to the 30° S. in regions where the temperature does not fall beneath 55° F. (13° C). The area of its cultivation is in fact very nearly the same as that of cotton. Within the tropical region it may be cultivated at the level of the sea or even much further to the north and south of the equator than has been indicated. The plant manifests, in other words, a remarkable power of endurance, but it does not follow that where it may be grown as an ornamental garden bush it may there afford the commercial product. Within the tropics it will yield profitable returns only

COFFEE CUL-
TIVATION.
1642

COFFEA
arabica.

Habit of Coffee-drinking

HISTORY

winds blow away the flowers and make 50 per cent difference in crop. If too hot and dry, the plants require shade, and if strong winds prevail during the flowering season, belts of forest have to be left to protect the plantation. This is regarded an important consideration in clearing land for a coffee plantation. Dr Shortt says "In low countries there is not sufficient moisture in the soil and when shaded and irrigated, it produces a coarse and uneven bean devoid of the peculiar aroma essential to good coffee."

the limits of marine influence On this account the recommendations of the early advisers of the Government of India to prosecute experimental coffee cultivation on the lower Himalaya from Darjiling to Kumson have been abandoned. The occurrence of certain wild species on the

seeds

It has been stated that the coffee plant of commerce is truly wild in Abyssinia and that it is there called *bun* or *boun*. This name

coffee was introduced into Aden by a certain Sheikh Shihabuddin

there arose after some few years, in 1511, a crusade against its use as un-

ed a Greek servant, Pasqua Rossie, for the purpose of preparing a favoured beverage. His friends grew so fond of it that to prevent their

Consumption of Coffee

COFFEA
arabica.

HISTORY.

too frequent visits to his house he recommended Rossie to start a public coffee-shop. This was opened in St Michael's Alley, Cornhill. Coffee-shops rapidly multiplied, but the beverage (although from a very different reason) soon met with as much official opposition in London as it had sustained in Constantinople. Charles II (in 1675) viewed these shops as the meeting-places for disaffected persons, and a royal proclamation was issued for their suppression. Coffee is spoken of as being in use in France in 1630, and the first public café was opened in Paris in 1669. Shortly after, it became general throughout Europe. It may be here added that of the three great dietary beverages Cocoa was the

trade which by 1847 checked the further development of the demand for coffee. There are doubtless many causes that may have contributed to bring this about, chief amongst them may be placed the facility with which coffee can be adulterated, the greater consumption of cocoa, and the ease

relative measures appear to have had much to say to the growth of a greater coffee consumption in continental countries than in England, or rather to the decline of coffee consumption manifested in Great Britain with the

—The consumption of coffee
in 1857, 34,518,555lb, in 1867,
to 31,859,408lb, and slightly

improved in 1880, being in that year 32,480,000lb. These figures must not be confused with the imports of coffee. Great Britain does an immense

BRITAIN.
Decline in
Consumption.
1863

throughout 110, and European Russia 4lb. The United States of America are supposed to use on an average 8lb per head of population per annum. Mr. H. Pasteur, in his report on the coffee shown at the

COFFEA arabica.		Coffee Cultivation Extended
HISTORY	Color of seed, leaf, and fruit. tion of tons and J being tons, Ceylon 10,000 to 12,000 tons, and Jamaica 4,000 to 5,000 tons Altho the quality is finer than that of Mocha,	
EXTENDED CULTIVATION 1644	which at one time stood above all others EXTENDED CULTIVATION.—The cultivation of the coffee plant began to extend towards the end of the seventeenth century, being carried on in produces more coffee than all the other plants in the world. In Brazil coffee is completely acclimatised, and there are said to be 530 million plants under careful cultivation. Coffee is also extensively grown in Costa Rica, Guatemala, Venezuela, Guiana, Peru, and Bolivia with Jamaica, Cuba, Porto Rico and the West Indian Islands generally. Its cultivation has been introduced into and India are the countries where its introduction has assumed an important commercial character continued by the natives of Ceylon. In 1825 the impetus to fresh effort was given by Sir Edward Barnes in the establishment of an upland European plantation. In 1877 it was estimated that the capital invested in Ceylon coffee was nearly 14,000,000. The coffee plant is a fungus	
CEYLON Introduction. 1645		

Introduction of Coffee Cultivation into India.

COFFEA
arabica.

cwt. in 1876, to 312,000 cwt. in 1884, and to 230,000 cwt. in 1885"
(Pasteur)

INTRODUCTION INTO INDIA.—The history of the introduction of coffee

HISTORY.
INDIAN.
Introduction.
1646

COFFEE
arabica

Coffee Cultivation—Locality

HISTORY.

thousands of acres of good suitable land for coffee near navigable rivers where manure and labour are cheap.

Coffee has also been introduced into Burma. For some time the effort to open out plantations seemed to be doubtful, and Mr Pettley, speaking of the garden on the Karen Hills, north-east of Toungoo, reported recently that much damage had been done by a mole cricket. Since then, however, the report is that the plantations are doing well, and the coffee is reported to have been sent to Upper Burma.

that the Arabian variety does best on the Toungoo Hills, while at Tavoy the Liberian variety is alone thought worthy of cultivation. "Local demands, too, are increasing, as land is being taken up along the lines of railway between Rangoon, Prome, and Toungoo, and gardens have been formed whereon small grantees are now cultivating fruit and other useful trees as well as coffee."

METHODS.
I 647

METHODS OF CULTIVATION

Space cannot be afforded to deal with every feature of this subject; the reader is referred to the numerous special publications quoted under the paragraph of references, only the more salient features will be touched upon, and especially those which have a bearing on the future expansion of the industry.

LOCALITIES, CLIMATES, AND SOILS SUITABLE FOR COFFEE CULTIVATION AS AN AGRICULTURAL PRODUCT—Under the heading "History of Coffee," the subject of the region of coffee cultivation and the climate necessary have been discussed. Dr Shortt says of soil, "This should be rich, abounding in moisture and containing much humus or vegetable mould, consequently we find that the plant thrives best on either red or black clay, containing combinations or preparations of iron, and covered over with humus formed by the decay of vegetable matter produced by dense forests. When these points are overlooked, the results are soon seen in the young plantation. The planter, perhaps, instead of choosing forest land, has taken up a poor grassy or stony expanse and however much water he may have access to, his plants are stunted and soon become yellow, unless he resorts to heavy manuring at a very early stage, which materially increases the expense of the concern. In hard rocky soils the plants require to be deeply excavated to permit of the tap roots of the plant striking perpendicularly down, and even when every precaution is taken, it will be found that estates opened out on poor soils will always prove more expensive than those on forest land, and are not so lasting. The berry produced on rich ferruginous clay is found to contain more aroma and the bean is heavier when compared with those of other localities. This fact is so well known to coffee-brokers generally that, in London, a new importation is frequently weighed after being roasted." Some difference of opinion prevails as to the degree of moisture the soil should contain. In *Sprays Encyclopædia* there occurs the following: "The points which determine the value of a plot for coffee culture are—1, elevation, 2, aspect, 3, shelter from winds, 4, shelter from wash, 5, temperature, 6, rainfall, 7, proximity to a river, 8, character and richness of soil. Most of these are necessarily subject to variation according to locality. Shelter from wind is perhaps of paramount importance and should not be sacrificed for richer soil, as the latter can be artificially obtained much quicker than the former. In wooded country the estate may be laid out in blocks of 20 acres, enclosed by

Coffee Cultivation—Seed

COFFEA
arabica.

METHODS.

deadly effects of a damp atmosphere, for, in all probability, he will have to spend his time surrounded by the direst malaria, &c.' Spens', on the other hand, says — "The most suitable climate is precisely that which

Nursery and Seed.—Having selected the site for a plantation, cleared and burned the trees (taking care, where necessary, to have protecting belts against prevalent winds), laid out the roads and carried the water-supply to the coffee-house, it next becomes necessary to select and pre-

Nursery.
1648

water

Seeds.
1649

the morning or after sunset.

The selection of seed is of great importance. The stock should be

12 inches apart from each other, so as to give the plantings plenty of room to grow, and subsequently enable the planter to remove them with facility from the nursery to the plantation, or the seeds may be sown in drills

COFFEA
arabica.

Coffee Cultivation—Planting.

METHODS.

and as the seedlings begin to grow the drills should be thinned out to the

planted, in damp, cloudy weather, from the seed-beds to the nurseries, and placed 9 to 12 inches apart. Care must be taken not to double up the tap-root, and not to leave a

roots. If the tap-root is very

when it soon shoots again

nurseries is not practised, the

have grown larger, but Stainbank and others strongly recommend the former plan, as, by checking the growth, the young wood becomes hardened, and better able, when finally planted out, to resist insects and unfavourable weather. A practical suggestion for preventing young seedlings being eaten off at the surface of the ground by grubs, is to lightly

wrap round a piece of paper about 3 inches broad, where the stem joins the root, on planting." (Spons)

Planting out
1650

LIVING AND PLANTING OUT—Soon after being cleared the estate is "lined out" if

are in vogue (

up and down

this line, stake

upon for the position of the plants. To each stake a rope is fixed, and stretched parallel with the base line and as straight as possible. Small

stakes are provided along these lines. A rope is finally held across them at succeeding stages of equal width, as guided by measuring poles, and

the small stakes are put in where the moveable rope crosses the fixed ones, each stake indicating the site for a plant. (2) A rope is furnished with

bits of scarlet rag at the distance fixed upon between the plants, it is stretched across the plot and stakes are inserted at each rag, the rope is

ring rods. The
it is more labor-
etch of the rope

in their perma-
lected for trans-
plantation, many coffee planters prefer to have two-year old seedlings

part, and the ques-
ree of shade, and

s are not likely to
reverse being the

th. In India the
distance adopted varies between 4 and 8 feet each way—7 feet being

very common, or 6 feet between the plants and 7 feet between the rows

points
th each

ion. If
'anting,

should
ng and

Cultural
operations
1651

C. 1651

effly re-
ition so

Coffee Cultivation—Shade.

COFFEA
arabica.

METHODS.

as to prevent the young seedlings from being choked. Staking, or sup-

porting, may not be enough, and the degree to which on the nature of the trees has deprived the

plantation of the natural protection which belts of trees would have afforded. According to many planters, however, all trees should be removed and shade procured through the cultivation of the charcoal tree (*Spondia Wightii*). In two years this forms an ample shade, but as it

is not so hardy as the trees which it replaces, it requires to be renewed. This is

regretted, in his report, as it is a matter for

regret," he

out break o

forwards of

is more important than a complete system of drains and roads. If the operations in this direction have not been completed up to date, the energies of the planter during the first two years may very appropriately be turned to these considerations. Drift surface-water not only removes the soil, but may altogether wash away the plants. A proper system of drainage becomes essential, not only to remove the water from damp and cold water-logged soils, but to provide against the dangers of sudden

Pruning,
1652

COFFEA
arabica.

Coffee Cultivation—Pruning.

METHODS

postpone the operation till the shrubs have borne their maiden crop, even though extra staking be required to withstand the wind. His plan is to remove the two primaries at the required height, by a sloping outward cut close to the stem, and then to remove the top by an oblique cut, so that the stumps resemble a cross, and a firm natural knot remains to guard against the stem splitting down. Hall (Ceylon) contends that the plants should be topped as soon as they have reached the required height, when the soft wood is easily severed by a pinch between the finger and

topped either at their full height—
ucker to grow up on the weather
atter plan is preferred. There is
the height to 5 feet, not only is
without damage to the tree, but
s are more readily made to cover

Dr. Shortt says "Pruning consists of various operations connected with either arresting the height of the plants to cause them to spread out laterally, or in removing the additional growth of wood, to encourage the plants to push out new

ping he
exposed
s it does
r ladders
there are
—vs, to

first result of topping is to induce the growth of masses of shoots, these are removed by what is technically called *handling*. "The first to appear are vertical suckers or 'gormandisers' from under the primary boughs: these are immediately rubbed off without injuring the bark. From the primaries spring secondary branches, in pairs, and at very short intervals. All such appearing within six inches of the main stem are removed at once, so that a passage of at least a foot is left in the centre of the tree for the admission of air and sun. The object of pruning is to divert the energies of the plant from forming wood and to concentrate them upon forming fruit. The fruit of the coffee tree is borne by young wood, and as the secondaries are reproduced when removed, they are cut off as soon as they have borne, and a constant succession of young wood is thus secured" (Spons). This removal of secondary twigs from the primary boughs is what the planters call "pruning". The practical effect of the treatment briefly indicated above is to cause a plant about 5 feet in height to develop horizontally primary branches or boughs at intervals of about 6 inches throughout the height of the stem, and to form along these boughs a constant supply of secondary fruit-bearing twigs. All ascending or cross-wise branches or twigs are at once removed, so as to force the plant into the arbitrary and unnatural type of horizontal spreading branches which have the advantage of exposing to the sun and light a large surface from which the crop can with ease be removed. When practicable, the bushes should be handled twice before the crop, and all

s begin to form, but
hat a flush of so heavy
e necessary to sacrifice

Coffee Cultivation—Season	COFFEA arabica.
<p>this by pruning the plant down to the extent it may be expected to fruit without injury. The lateral or primary boughs should not be allowed to grow more than 2½ feet, otherwise they will droop and exclude the light from those below. In pruning, it is often recommended to leave the white, both stems and leaves, of its fruiting next year. A continuous crop is nipped off, broken, dis</p>	METHODS.
<p>CATCH CROPS.—Much has been written for and against the growing of other crops along with coffee. In Darjeeling it was tried to grow tea and coffee together, but with little or no success, in spite of the fact that the out door labour and manufacture of these crops so fit into each other that economy might be effected. In Natal and other countries, plantains, ven of ind</p>	Catch-crops 1653
<p>SEASONS FOR COFFEE PLANTING AND MANUFACTURING OPERATIONS.—The industry being chiefly in South India, the seasons for operations very closely correspond with those of Ceylon. The season for</p>	Seasons. 1654

for the collection of the crop and the manufacture of the berries. The fruits commence to ripen in October or early in November and continue till January. Thus from flowering to harvest occupies about eight months. None but fully ripe berries (technically known as 'cherries') should, according to Dr Shortt be collected, the women and children going over the plantation periodically to remove all the bright or blood red ones, while carefully leaving the others to mature, once ripe, the sooner collected the better. Mr Pasteur says "The usual course, however, is to pick the cherry before complete maturity, when it is of a deep red or cherry colour, the berry inside being then found to be of a fine dark green or bluish green, which it is the endeavour of the planter to preserve as carefully as possible the value of his coffee depending chiefly on the depth and brightness of the colour." The more gradually the bloom fades the better

berries to be picked up, also the berries that have fallen to the ground. This forms what is generally known as "jackal coffee." Before the boughs are opened out again, the ground around each plant is *manured* and *forked*.

The preparing or manufacturing of the "cherry" into the "berry" will be found dealt with in a further page.

INDIAN AREA UNDER, AND OUTTURN OF, COFFEE

The cultivation of coffee is practically confined to Southern India. During the three years 1883, 1884, and 1885 the average area under mature

INDIAN.
Area and out-
turn
1655

C. 1655

COFFEA
arabica,

Area of Coffee Cultivation in India.

AREA AND
OUTTURN.

plants was returned at 186 500 acres, and the average yield at 31½ million pounds, which were thus distributed —

	Acres	lb
Mysore	82,100	7,110 000
Madras	55,100	13,160 000
Coorg	42,300	9,330,000
Travancore	4 800	820,000
Cochin	2 200	830,000
TOTAL	186,500	31,250,000

These statistics, which are in all probability defective, have been taken from the Statistical Tables of British India published by the Department of Finance and Commerce up to 1887. These tables include the Native States of Cochin, Travancore, and Mysore, and hence the area given is greater than that returned (to the British Government) by the States of British India.

In British India of the Nilghiris it has been said that there exists 200,000 acres of reserve suitable for coffee. The port of shipment for Nilghiri coffee is Calcutta.

not likely to extend, as the cultivation of coffee in Mysore has been its, 81,543

for coffee-planting progressing much further than at present, except on the sheltered tracts.

"A northern aspect is best, being moist during the dry season, and possessing the most uniform temperature, blowing eastwards or westwards according to the prevailing winds. On the western slope the cultivation is not so successful."

may be found useful —

In Mysore the cultivation is limited almost exclusively to the Kadur District. In Malabar, the cultivation is limited almost exclusively to the

MYSORE
1656

Area of Coffee Cultivation in India.

COFFEE
arabica.AREA AND
OUTTURN.

planters was Mr. Cannon, who formed an estate on the high range immediately to the south of the Baba Budangiri, where the original coffee-plants are still in existence flourishing under the shade of the primeval forest

"The success of Mr. Cannon's experiment led to the occupation of ground near Aigur in South Manjarabad by Mr. Green in 1843. During the last fifteen years, estates have sprung up between these points with such rapidity that European planters are settled in almost a continuous chain of estates from the northern slopes of the Baba Budans to the southern limits of Manjarabad, not to mention Coorg and Wynaad beyond."

The above account of the introduction of coffee into Mysore was first published by Colonel Onslow, from whom all subsequent writers have borrowed their information without materially adding to or correcting any one feature of the original statement.

Madras Presidency—The following extract taken from pages 290 and 291, Vol I of the Madras Manual published in 1885, gives interesting particulars regarding the cultivation of coffee in the Madras Presidency: "The principal coffee tract of Southern India is along the western coast, and coffee estates extend in nearly an unbroken line along the summits and slopes of the Western Ghats, from the northern limits of Mysore down to Cape Comorin. The only portions of the area within the limits of the Madras Government are the Wynaad tract and the Nilgiri Hills, the rest being in Mysore, Coorg, and Travancore."

Of the early plantations the Madras Manual adds. "Nearly all the land taken up at this period was what is known as grass or bamboo land, and in consequence most of the estates proved unprofitable. Of many of them not a trace, except the ruins of bungalows remains at the present day. After the first attempt at coffee cultivation in the Wynaad district, the only other attempt was made in the Coorg district in 1843, when a small estate was formed by Mr. Cannon."

MADRAS.
1657South
Wynaad
1658.

1868, and, according to the returns then made, the acreage was 29,909 08, of which 21,479 54 acres were held by Europeans and 8,429 54 acres were held by natives.

	Cwt
1856-57	37,658
1857-58	16,204
1858-59	36,934
1859-60	49,680
1860-61	48,742
1861-62	91,080
1862-63	43,907
1863-64	91,947
1864-65	110,548
1865-66	125,891
1866-67	66,552
1867-68	128,011

C. 1658

COFFEA
arabica.

Area of Coffee Cultivation in India.

AREA AND
OUTTURN
Nilghiris
1659

"Coffee cultivation on the Nilghiris was reported on in 1872. A large area of land on the Nilghiris has proved to be admirably suited for the cultivation of the coffee shrub. Not less than 22,897 acres are now under coffee plantations besides 12,231 acres taken up for planting. Twenty-five years ago the area under coffee did not much exceed 500 acres. This great increase is entirely the result of private enterprise, and has added much to the prosperity of the Nilghiris, while at the same time benefiting the districts immediately adjoining. In the establishment of these coffee estates a property has been created worth about 5 millions of rupees. Of the total expenditure, about one third is for the payment of

... of labouring people
king, &c., a
previous to
only on the

eastern slopes, but they have now been extended to the southern, northern, and north-western slopes, there are also some extensive plantations in the Ouchterlony Valley and in the neighbourhood of Coonoor. Coffee cultivation is also carried on on the Shevaroy Hills in the Salem District, where nearly 6,000 acres are under the crop, and an area of 4,680 acres has been taken up for planting, on the Pulney and Shiroomullay Hills in Madura, where nearly 4,400 acres have been planted and a considerable area has been taken up for planting, and in the Tinnevely and Combaratore Districts, in the former of which there are about 2,000 acres under coffee and in the latter about 800 acres."

Coorg.
1660

... and for there are but few

-port

uro-
a little

ssment

56,440

lots of

size of

Of the

whole area 40,350 are bearing, producing 6,125 tons of coffee, or on an average

which

acre

of

estates,

cultivation at the rate per acre assumed above comes to nearly 32 lakhs of rupees. Of this not less than 60 per cent on an average may be estimated as having been paid to labourers in wages. Calculating that 26,893 labourers, which is about the average number employed throughout the year, received Rs 6 each per mensem, upwards of 19 lakhs of rupees were expended for labour during the year. The value of the coffee produced, taking the selling price to be, on the average Rs 75 per cwt on the spot, was about 36 lakhs of rupees." (*Madras Weekly Mail*)

Travincore—The area under coffee in the former State in 1885 was 4,013 acres, and in the latter 2,407 acres. The area under coffee in Travincore seems to have declined considerably within the past few

Travincore
1661

C. 1661

Coffee Manufacture

COFFEA
arabica.AREA AND
OUTTURN.

the group known as the Anamallays. The plateaux, by reason of their good climate, rich soil, abundant timber and water-supply, are likely to become better known as the demand for coffee land increases. One plateau alone (Eroovimullay, or Hamilton's Valley) is 6 miles long by 3 wide, and contains about 10,000 acres of excellent tea and coffee land.

In Cochin there were, in 1883, 17 gardens, and these gave the return of 342lb to the acre at a cost of Rs 4.

TECHNICAL TERMS USED BY THE COFFEE PLANTERS.—The ripe coffee

COCHIN
1662Technical
Terms.
1663

chinery necessary for this purpose

PREPARATION OR MANUFACTURE

The preparation of the "berry" from the "cherry" may be said to be accomplished in the following stages: (1) *Pulping*, (2) *Fermenting*, (3) *Drying*, (4) *Peeling, Milling, or Hulling*, and (5) *Sizing and Winnowing*.

A volume might be written on the various systems and mechanical

being tedious, ineffective, and expensive this process does not secure the

MANUFACTURE.

Pulping.
1664

ively accomplished if the collections of ripe cherries made each day are passed through the machinery at once. If unavoidably delayed it may be necessary to ferment the cherries before they can be pulped. The most simple machine in use is that known as the "disc pulper." This consists of rotating discs the surfaces of which are covered with sheet copper roughened by having projections punched forward. A "single pulper" of this description will pulp 20 to 25 bushels an hour and may be worked by three coolies. A "double pulper" of this type has two such discs and is furnished with a feeding roller. It will pulp 40 bushels an hour, and may be worked by from four to six coolies, and double that amount if worked by

C. 1664

COFFEA
arabica.

Coffee Manufacture.

MANUFACTURE.

steam The discs work against smooth iron beds so adjusted that the complete cherry cannot pass between. They are torn upwards against the beds, and the projections on the discs tear off the pulp, allowing the beans to drop into one receiver and the fragmentary pulp to be carried into another. The disc pulper is in fact somewhat like the cotton gin which drags the fibre forward and drops the seed behind. The 'cylinder pulper' is an older invention in its conception, but has been improved and perfected to a much greater extent than the disc, the latter, being

cherries are spread out—the pulper

By constructing this building again

cherries may be carried direct into

raised. A good supply of water has also to be conveyed to the loft so as to descend with the cherries into the pulping machine in a continuous stream.

Space cannot be afforded for a discussion of all the inventions and contrivances

from the bean. The
s are carried
re separated
o pass once
s is carried

from the loft of a tube which dips to the bottom of a basin known as the *hopper*. Stones subside in the hopper, while the continuous stream from above causes the hopper to discharge a uniform supply of cherries and water to feed the pulper.

FERMENTING—The parchment coffee, which may or may not have been assorted by contrivances in the pulper and sieves, has now to be fer-

If this be not accom-
ing advantage of the de-
tanks, and these tanks
platforms on to which
sed. There are gener-

ally four fermenting tanks—two in which the fermentation actually takes place, and two in which the beans are washed. One of each is used for the produce of one day's pulping. All the coffee pulped in one day is allowed to remain in the front or receiving cistern until fermentation has set in. The period necessary for this will depend greatly on the temperature of the atmosphere, but from 12 to 18 hours will generally

into the washing
s pro-
may be
as to

Fermenting
1665

Coffee Manufacture.

COFFEA
arabica.

MANUFACTURE.

Drying
1666

and are accordingly preferred. The tanks should slope towards the dis-

event of an occasional shower, but shed accommodation into which the beans may be rapidly conveyed is essential. During the drying, the beans have to be turned over repeatedly either by rakes or by the coolies'

many cases, however, there are neither appliances, time nor labour, to put the fresh-gathered fruit in the sun the cherry dries quickly, detriment of the colour as well difference between *unwashed* or plantation coffee,—the taste of the *washed* coffee being, as a rule, much more delicate, and free from the earthiness and common rough flavour of the *unwashed*

PEELING OR MILLING—This consists of the removal of the parchment and silver from the beans. As already stated, this operation is now chiefly effected by the dealers, at the port of shipment, and not by the planters. Indeed, much has been written in favour of the beans being sent to use in London for eur's report will be indicating a pos-

Peeling.
1667

"Among the samples of Wynaad coffee, those from the Eva Estate deserve special attention, one half of that crop having been despatched in parchment to be peeled and sized in London. The experiment has proved

COFFEA
arabica

Coffee Manufacture.

MANUFAC-
TURE.

coating almost immediately after being picked The curing requires machinery, motive power, drying grounds, delicate manipulation, and constant supervision, where any of those requisites fail, the coffee suffers in appearance, and consequently in value. Suitable machinery for treating parchment has been erected at two of the London wharves, and there is every reason to hope that this is only the beginning of a new and profitable home industry Growers will not be slow to perceive that the small increase of freight which they have to pay on parchment is more than compensated for by the enhanced price which the improvement in the quality of their coffee will enable them to obtain" In the *Kew Bul-*

Sizing
1668

passed through the mill the beans require to be again heated On the plantation this is generally done by exposure to the sun The extent to which this is necessary depends greatly on the nature of the beans, and long experience is required to determine this point As a practical hint it is generally laid down that they should be dried till they resist the pressure of the thumb nail, but no two samples are alike, and overdrying will

Packing
1669

universally employed

PACKING --Having followed all the precautions and adopted all the most approved methods and appliances, the coffee producer, to secure the success of his labours, has now only to attend to packing The beans must be saved from exposure to the air, or from being packed in cases that would impart a false aroma This is usually done by packing the

ADULTER-
ANTS
1670

ADULTFRANTS AND SUBSTITUTFS FOR COFFEE

ally
uch
as
her
dietary article that is so much and so persistently adulterated as coffee This in a large measure appears to be due to the legislative system which has permitted a mixture to be sold so long as it is declared to be such Criminality consists alone in selling as *pure coffee* an article that contains anything but coffee Legally "chicory" may be the roasted chicory root itself or the root of an allied plant or other vegetable substance applicable for the same purpose as chicory No questions are therefore raised as to the ingredients of a mixture, and indeed, if further protection to the manufacturer be necessary, such mixtures may even be registered as patent medicines This fact, together with the long-established custom of mixing chicory with coffee, has given origin

Adulteration of Coffee

COFFEA
arabica.

to a gigantic system of adulteration. The substances which are most generally employed are—

"1st—Roots such as chicory, dandelion, mangold-wurzel, turnips, parsnips and carrots, &c

"2nd—Seeds such as beans, peas, date-stones, malt, rye, &c

"3rd—Burnt sugar, biscuits, locust-beans, figs, &c" (*Bell, Chemistry of Foods*)

Association formed in examining certain well-actices of adulteration s attention was the use on of the real article that the mixture of the spurious with the true coffee beans might be fearlessly ground in the purchasers' presence and sold as *pure coffee*. This subject has already been alluded to under *Chicory* (see *Cichorium Intybus*, C Nos 1107 & 1108), and need not be elaborately dealt with in this place.

ADULTER-
ANTS.

le without being viewed or a sugar-yielding root becomes a serious adul- used of all adulterants

Caramel

facturing special preparations or mixtures. Roasted flour coloured with ferruginous earth is to some extent used as a coffee adulterant, and even roasted liver and other objectionable animal substances are said to have been found in coffee mixtures. A simple mode of detecting the presence of chicory or other caramel admixtures in ground coffee is to throw a little on the surface of a glass of clear water. The readily solvent nature of the

The seeds of several species of *Cassia* have for centuries and are even now used by the inhabitants of tropical countries in place of coffee. These do, as a matter of fact, afford, when roasted and ground, a decoction which closely resembles coffee. The reader is referred to the account given under *Cassia occidentalis* (C No 784) for particulars of a coffee substitute which would seem to deserve more careful consideration. India could produce, at a nominal price as compared to coffee, immense quantities of the so-called "Negro Coffee," if that article should be found to commend itself as a wholesome and cheap substitute for true coffee.

Negro
Coffee.

COFFEA
arabica.

Trade in Coffee.

ADULTER-
ANTS.

The work article. others in injurious reputation, and to place in the hands of the consumer a cheap and pure coffee.

COMMERCIAL
TERMS.
1671

age, and uniformity within the sample. Form to some extent, though not always, depends upon the source: there are three commercial types as to form—*Mocha*, small round peaberry; *Bourbon*, pointed and medium-sized; and *Martinique*, large and flattened. Colour depends entirely on the degree of roasting which is given to the coffee in the preparation.

PRICES.
1672

PRICES OF INDIAN COFFEE.

valued as high even as those of Ceylon; and, as stated in another paragraph, Mr. Pasteur, one of the highest commercial authorities, gives the first place to the Indian coffee.

of native coffee was sold for the same price as a bushel of rice, viz. Rs. 44 and, about the same time, estate coffee from the Wynaad was selling on the coast for Rs. 100 per bushel. (1869), estate the work- e is insign-

TRADE.
1673

TRADE IN INDIAN COFFEE

"India now stands first and foremost among British possessions, both for the quality and quantity of its production." Disease has, however, "in many places affected the vitality and shaken the strength of the trees, so that they have been less able to resist periods of drought or of heavy monsoon weather, and small and irregular crops have been the consequence. It would seem, however, as if plantations were gradually recovering their former strength, and with good cultivation and manuring

Indian Trade in Coffee.

COFFEA
arabica.

TRADE

and fair seasons India may hope to maintain its position as our largest and best field for the production of fine coffee. A hopeful sign for the future may be gathered from the fact that the average yield of the

from 47,000 to 38,000. This has been accounted for by the fires which destroyed certain gardens, the imperfect returns, and the amalgamation of small gardens. The bulk of the coffee exported from India is washed coffee prepared under European supervision, many of the small native planters selling their produce to neighbouring European planters or to the special firms that do a considerable trade in pulping and peeling coffee. At the same time, there is by no means an inconsiderable trade in unwashed or native coffee,—that is, coffee prepared by the crude native process to which reference has been made. Mr Pasteur, in his report of the coffees shown at the exhibition, remarks on the paucity of the samples which would seem to commend the washing industries.

Exhibition, they are quite suitable for our home consumption, and form an important item of the Indian production." The returns for the coffee districts of India show Madras to have nearly a third of its coffee area owned by natives, Coorg about one half, and Mysore fully four fifths. These facts give some idea of the extent of the probable production of native or unwashed berry in India.

400,000 cwt. The average export of the coffee of India is about 400,000 cwt. howe. Coch sent 1 impo supply, as a rule to it come Ceylon and Aden. Bombay receives most of this coffee, a little goes to the other ports. of the total exports shipped chiefly from India of foreign and Madras.

COFFEE
arabica

Trade in Coffee

TRADE

two largest consumers of Indian coffee. During the past five years the coasting trade which consists chiefly of despatches from Madras to places within the presidency and to Bombay, has averaged in quantity 70,000 cwt and in value Rs 22 lakhs.

Towards the close of the account given, on a preceding page of the History of Coffee, Mr Pasteur's statement regarding the decline of the Ceylon trade has been quoted. With the discontinuance of a large part of the Ceylon cultivation the greatest hopes were entertained of a bright future for the Indian coffee industry. Prices revived from 1883 to 1887 and during that period the exports to foreign countries maintained a higher level than during any previous consecutive period. During the

advantage of the decline of the Ceylon industry. The Indian foreign trade in coffee has chronically fluctuated. It attained its highest recorded point of the trade with the Madras exports (given at page 473) from 1836-57 to 1867.

COST OF CULTIVATION AND YIELD

So much has been written on this subject that it scarcely falls within the scope of the present article to deal with the various conflicting opinions that have been advanced. According to some writers the profits on coffee cultivation in India are problematic, according to others the in-

follows —

	R
1st year	7 160
2nd year	3 300
3rd year	4 450
Instruments	700
Buildings and roads	1,830
TOTAL	17 400

This estimate, he states, is applicable to Coorg and Wynad, more especially the former, but he only allows Rs 125 a month for European supervision. He proceeds to state that the third year is supposed to make a return. The average produce of an acre is estimated at 7 cwt, but we could not do better than keep on the safe side and take the produce of an acre at 5 cwt. The 200 acres will yield 1,000 cwt of coffee.

Cost of Cultivation

COFFEA
arabica.

COST

the erection of a pulping house, and other accessories to the preparation of the bean but Dr Shortt adds with reference to this that 'these will at best form but a small item'. But he has omitted apparently to estimate for the purchase of grass and forest land, and to take into consideration the cost of the labour of preparing the beans.

The author of the valuable article on coffee planting in *Spens' Encyclopedia* gives several estimates both for India and for Ceylon. He states 'The following estimate (in rupees) for coffee cultivation in South India is based on the purchase of 300 acres of forest land at Rs 50 and 200 acres grass land at Rs 25 bringing 200 acres of the former into full bearing, labour, 4 annas a day, exclusive of maistries' wages.' Then follows a balance sheet the main facts of which may be expressed as follows —

The 200 acres by the seventh year are brought under full bearing and have not only cleared off the expense of the purchase and cultivation of the estate up to date but the plantation has given its owner over and above Rs 971. To continue to work it an expenditure of Rs 644 would be entailed but the return from the crop would be about Rs 54,000 a year so that with a portion of this the estate might now be extended to its full limits 300 acres. This estimate has not only been framed to cover the charge of building all the necessary houses, but to furnish those with pulping and other machinery, and to stock the yard with 100 head of cattle and provide a horse for the superintendent. The capital necessary to organise such an estate (without having to obtain loans on crops) would thus be about Rs 75,000 or say £5,000, and during the fifth, sixth, and seventh years that sum would be recovered. Interest on

is however unable to verify these estimates but since they have been framed by high authorities they may be viewed as approximately indicating the possibilities of the Indian coffee industry when, with average seasons and fair prices the speculation is entrusted to careful and skilful supervision.

COFFEA arabica

Diseases of the Coffee Plant

PROFITS.

ready made estates, and pleased their own minds and those of the other shareholders with visions of 50 or 60 per cent of profit. As might have been foreseen, such extravagant hopes have never been realised, the anticipated fortunes having retreated far away into the future, and the 50 or 60 per cent dwindled down to 5 or 6. In many cases, indeed, these adventures have, from various causes, proved complete failures, the balance always being on the wrong side, and, taking them as a whole, the results have been such as to render the public distrustful of coffee culture as a safe or profitable investment, and to lower greatly the value of estates" (*Report on the Ravages of the Borer on Coffee Estates*)

DISEASES. 1675

DISEASES OF THE COFFEE PLANT.

There are, however, certain specific diseases some of which have practically baffled both the planter and the scientists, and have proved so disastrous as to have ruined the plantations in large tracts of country. This has been the case with Ceylon, the leaf blight having there proved so far incurable as to have caused the planters to substitute tea for coffee on their estates. Numerous reports have been published such as those by Marshall Ward, Nietner, Bidie, Harman, Forbes Watson, Morris, Cooke, Balfour, &c. To review even briefly all that has been written on the diseases of the coffee plant would take up far more space than can be afforded in the present outline of the coffee industry. It may be said that the specific diseases are referable to two sections—*Fungoid* and *Insectiform*.

The chief FUNGOID diseases are —(a) *Leaf blight* —This is a fungoid disease which is supposed to have first made its appearance in Ceylon in 1869 and to have appeared in South India two years later. It has since appeared in Java and Sumatra, but does not seem to occur beyond the limits of the Indian Ocean. It is caused by the fungus *Hemuleia vastatrix*, an organism allied to mould. It attacks the underside of the leaves, in the form of spots or blotches, at first yellow, but which ultimately turn black. These spots are covered with a pale yellow powder. They eventually extend over the whole surface of the leaf, which then drops from the plant. The disease is generally developing or ar, but in its in the form of k and leaves al cure, but with little success. If powdered sulphur, alone or mixed with caustic lime, be blown over the plants and scattered on the ground below the boughs, the disease is prevented and the coffee plants seem at the same time to be benefited. This is, however, expensive and is more a preventative than a cure. When once the disease has taken hold of the leaves nothing has yet been discovered that will destroy it without at the same time killing the leaves.

(b) *Leaf rot or Candelillo* is a disease attributed by Dr. Cooke to the fungus *Pellicularia Koleroga*, Cooke. It is prevalent in Mysore plantations in July, the leaves, flowers, and berries becoming covered with a shiny

Diseases of the Coffee Plant

COFFEA
arabica

DISEASES

gelatinous substance which turns black about the time that the affected parts fall from the plant (*Kew Reports 1879, 30 and 1880, 15*)

II Of the INSECTIFORM diseases met with in India the following are those which give most trouble —

(c) *Borer* — This pest used to be known as the “worm” and “coffee fly”. It is most troublesome in South India, especially in Coorg and the Wynaad, where in 1865-66 it destroyed whole estates. It has been determined as the beetle *Xylotrechus quadripes*. It is red or yellow, with black in transverse lines. It damages the trees by boring holes into the stem usually a few inches above the ground. These passages are at first transverse but soon ascend spirally to the growing tip where the larvae are matured. The plant early shows signs of death, and ultimately withers down to the point where the beetle entered. This pest is most prevalent in hot exposed gardens, and may be kept in check by free irrigation.

destruction of the parts to which it adheres the flowers and young fruits fall off freely. The pest does not do much harm however until it has been two or three years on an estate. It prefers cold damp plantations at about 3000 feet in altitude. This bug may be first recognised as brownish wart like bodies. These are the females each of which produces some 700 eggs. Fortunately this pest is freely attacked with parasites which greatly help the planter.

The black bug is known as *Lecanium nigrum*. Like the preceding this attaches itself to the tenderest shoots. It also prefers gardens at high altitudes in damp situations. The female somewhat resembles a scollop-shell. When the eggs are incubated the twigs become covered with an
the young berries
what like a wood-
e. It is flat, oval,
ng across the back.

It seems to prefer hot dry plantations and disappears with the rains, only to return in time to destroy the setting of the fruits. It is found on the roots about a foot below the surface of the soil in the axils of the leaves and among the clusters of flowers and young fruits. It may be easily recognised by the white excretion formed around the larvae.

All these and the other less known coffee bugs have a strong dislike to tobacco juice. They may be prevented from developing to an injurious extent by brushing the twigs with tobacco. Some planters recommend saltpetre and quicklime in equal proportions dusted on to the affected

COFFEA
arabica.

Diseases of the Coffee Plant.

DISEASES.

by hand has been tried, but it can only be attempted upon young trees without crop; and Mr. Nietner, although allowing that an immense

now is " (*Balf Cyclop*)

(e) *Grub*—The larvæ of the moth *Agrostis segetum* are very destructive: this disease is known to the planter as "Black Grub." It appears about August to October. It lives in the ground, but during night comes out to feed and does much harm when very plentiful. It is, however, local, preferring certain parts of the estate, but does not confine its ravages to the coffee plant only, as it eats any cultivated plant—vegetable or fruit tree—but despises weeds. It is very destructive to young plants. Mr. Nietner states that he lost as much as 25 per cent of his seedlings through this pest. The "White Grub," this includes the larvæ of several species of *Melolonthidæ* or Cockchafers. These do much damage by eating the roots of the trees. Mr. Gordon considers them as one of the greatest ene-

it
a
y
c
1

berries form the so called Jackal Coffee.

COFFEE-
LEAF TEA.
1676

COFFEE-LEAF TEA.

It has long been known that coffee leaves, if cured by a process similar to that adopted with tea leaves, afford a beverage which contains sufficient caffeine to entitle it to a position as a cheap substitute for tea or coffee. Indeed, according to some writers, the leaves contain more caffeine than the berries. A decoction from the leaves is said to be regularly used by the inhabitants of Sumatra, especially at Padang. A Mr John Gardener of London even patented a process for manufacturing and partially roasting the leaves, from the belief that they were likely to come into use in Europe. Unfortunately, however, the leaves have an unpleasant senna-like flavour which greatly militates against their chances of European popularity. But perhaps the chief objection to coffee-leaf tea rests on the fact that the plants will not afford both a crop of leaves and fruits, and the latter is therefore never likely to be subordinated to the former as a commercial article. But for this fact coffee-leaf might be sold at 2d a pound as compared with tea at 10d.

The following note has been furnished for this work by Prof. Warden of the College Medical College—

the principle
e, contained
but during

The Uses of Coffee

COFFEA
arabica.

the roasting of the berries a larger amount is developed, to which the aroma is due. Caffeine appears to act as a stimulant to the nervous system. Coffee leaves have been used as a substitute for the berries; they contain caffeine. Mr N M Ward of Padang writes regarding the use of the coffee leaves as follows: 'I was induced, several years ago, from an occasional use of the coffee leaf to adopt it as a daily beverage, and my constant practice has been to take a couple of cups of strong infusion with milk in the evening as a restorative after the business of the day. As a beverage the natives universally prefer the leaf to the berry giving, as a reason, that it contains more of the bitter principle, and is more nutritious.' The best mode of roasting is by holding the leaves over a fire made of dry bamboo or other wood which gives little smoke. When sufficiently roasted the leaves have a buff colour, they are ground to a powder and used in the same way as coffee. (*Hanbury*)

COFFEE PULP

COFFEE
PULP.
1677

It has long been known that the ripe pulp of the coffee cherry contains an amount of sugar which might with advantage be converted into alcohol. At present the washings from the pulping machine are run off and no advantage taken of the sugar they contain. Several writers have urged the planters to utilise this by-product, but as yet no definite steps have been taken in that direction. It is indeed even questionable whether or not it would pay the planter to divert his attention to a perfectly distinct enterprise. The tendency of the present day is to enable the manufacturer in every branch of industry to compete to the last degree by affording him the means of deriving additional revenue from the waste or by-products of his industry. In this light it seems possible that the coffee pulp might be utilised in some way.

The pulp is a very little used and is usually discarded. This husk is employed in the preparation of the infusion known as *kahwe* or *kischer*. Dr Shortt states that according to his experiment 8 oz of dried husk, when steeped in water until fermentation sets in, yielded on distillation 1 oz of spirits. If not employed in this manner, might not the dried husk find a demand as an auxiliary to cattle food?

OIL

OIL
1678

The term 'Coffee-oil' is in the trade given to palm oil in which the kernels have been more or less burnt during the process of extraction.

The aroma of the coffee is lost in the process of roasting. By this means the aroma might be restored to the coffee or employed to flavour liqueurs. This empyreumatic oil is formed during the roasting, and probably at the expense of caffeine and other constituents of the coffee (see under Chemistry).

MEDICINE.

MEDICINE
1679

Coffee while not officinal in the British Pharmacopoeia is so in that of the United States of America. Many medical men, however, recommend its use in England for mild affections. Its dietary property, as a

C 1679

COFFEA
arabica.

The Uses of Coffee.

MEDICINE.

stimulant to the nervous and vascular system, is that upon which its claims to medicinal recognition depend. It produces a feeling of buoyancy and exhilaration resembling the first effects of alcohol, but it is not followed by depression and collapse. It increases the frequencies of the pulse, and stimulates the system to throw off feelings of fatigue, or to sustain prolonged and severe muscular exertion. It has even been contended that caffeine has the power of checking the waste of the tissues. Lehmann found that the distilled oil had this effect in quite as strong a degree as tea. The well-established property of coffee in preserving wakefulness depends upon its stimulating property on the nervous system. When swallowed it produces a warming cordial impression on the stomach, quickly followed by a diffused agreeable nervous excitement which extends itself to the cerebral function giving rise to increased vigour of imagination and intellect without any subsequent stupor such as follows on the use of most other stimulants. Moleschott found that it

cient energy of the brain are manifested without congestion or inflammation. In light nervous headaches, not proceeding from derangements of the stomach it often proves immediately effectual. It has acquired much reputation as a palliative in the paroxysms of spasmodic asthma, and has been recommended in whooping-cough and in hysterical affections. "Hayne informs us that in a case of violent spasmodic disease,

highly recommended in cholera infantum, and it has even been used with asserted advantage in cholera. It is said also to have been used successfully in obstinate chronic diarrhoea" (*United States Dispensatory*)

Coffee is much less astringent than tea, and hence it does not cause constipation so readily.

Wood states that "upon those who use it habitually, its characteristic influence is not fully evinced, as it has either lost its power in a great

tonic to the digestive organs, and more astringent in consequence of the amount of tannic acid it contains. Certain it is that tea, especially black tea, may be deleterious."

Pharmacology, I, 625)

th.
or
wit
bei
the
Jo

Chemical Composition of Coffee

COFFEE
arabica.

MEDICINE.

coffee in France is supposed to have abated the prevalence of gravel in that country. In the French colonies, where coffee is more used than in the English, as well as in Turkey, where it is the principal beverage, not only gravel, but gout, is scarcely known."

Unroasted coffee has been employed in intermittent fever, but it is much inferior to quinine. Roasted coffee is said to have the effect of imposing animal and beneficial application

coffee, burnt in the wards of a hospital early in the morning, is a deodoriser, and a very fragrant one" (*P. Kinsley, Honorary Surgeon, Chicacole, Ganjam, Madras Presidency*). "Is also an antispasmodic, when consumed in large quantities, is supposed by the Arabs to have an anaphrodisiac effect" (*A. S. G. Fayakar, Surgeon Major, I. M. D., Muskat, Arabia*). "Dried coffee roasted in an open vessel is a useful deoderant" (*Henry David Cook, Surgeon-Major, Calicut, Malabar*). "Is an antidote in opium-poisoning" (*G. A. Watson, Allahabad*).

CHEMISTRY.

CHEMISTRY.
1680

The roasting or torrefying of the coffee-beans, combined with the pulverising they are afterwards subjected to, induces certain changes to which in a large measure the flavour and aroma of the coffee are due. The woody tissue becomes friable, and at the same time certain chemical changes take place. The chief organic constituents of raw coffee are caffeine, fat, caffeic acid, gum, saccharine matter, legumin, and cellulose. Payen gives the following analysis —

Cellular tissue	34.000
Hygroscopic moisture	12.000
Fat	13.000
Starch, sugar, dextrin, and vegetable acids	15.500
Legumin	10.000
Chlorogenate of potash and caffeine	3.5 to 5.000
Nitrogenous matter	3.000
Free caffeine	0.800
Thick insoluble ethereal oil	0.001
Aromatic oil	0.002
Mineral constituents	6.697

Bell (in his *Chemistry of Foods*) gives the following table of the analysis of two samples, raw and roasted, of both Mocha and East Indian coffees. We reproduce the table, both because of its allowing of comparison between these two coffees and of indicating some of the chemical changes effected by roasting. —

Constituents	MOCHA		EAST INDIAN	
	Raw	Roasted	Raw	Roasted
Caffeine	1.08	.82	1.11	1.05
Saccharine matter	9.55	.43	8.90	.41
Caffeic acids	8.45	4.74	9.58	4.52
Alcohol extract, containing nitrogenous and colouring matter	6.90	14.14	4.31	12.67
Fat and oil	12.60	13.39	11.81	13.41
Legumin or albumin	9.87	11.23	11.23	13.13
Dextrin	.87	1.24	.84	1.38
Cellulose and Insoluble colouring matter	37.95	48.61	35.60	47.42
Ash	3.74	4.56	3.08	4.88
Moisture	8.98	0.03	9.64	1.00
	100.00	100.00	100.00	100.00

COFFEA arabica.

Chemistry of Coffee.

CHEMISTRY.

Should the whole of the testa of the seed (the silver skin of the plant-
beans) be removed, or is it necessary to leave the process of roasting

roasted together, the coffee will be much inferior to that obtained by
roasting carefully picked and assorted beans. The degree of roasting
required for one class of coffee is not the same as that for another. The
heat should not be greater than is sufficient to impart a light brown
colour to the bean. When roasting is carried too far, a disagreeable
smell and a bitter and acrid taste gradually mingle with the essential
aroma, and thus lessen the merit and value of the coffee. By reducing
the beans to charcoal the aroma and flavour are entirely destroyed. When
the roasting has been effected to the right extent, the volatile oil is
produced at the expense of some of the other constituents. A glance at
the table above will show that nearly the whole of the saccharine matter
has disappeared. This is not the case with the sugar in chicory or other
roots, a large proportion remaining as sugar, and hence the rapid colour-
ation imparted to water by a coffee powder containing chicory or other
cane-sugar-yielding roots, as compared with pure coffee. There is some-
thing altogether peculiar in the behaviour of the sugar of coffee under
the influences of torrefaction. How the volatile oil is formed seems to be a
puzzle. This oil has been termed Caffeone, and it is the aromatic principle
of coffee. It is wholly the product of torrefaction, the materials of which
it is formed being obtained by the destructive influence of heat on the

quantities,
eco-
con-
the

roasting, takes a similar
beverage produces"
principle upon which
not appear to be alter

found in tea. Weight for weight, tea yields about twice as much theme
as the roasted coffee-beans yield caffeine. On this account a greater

of nutri-
ling (as
the full

nutritive property of the bean is secured. Several writers have strongly
advocated the adoption of this practice, but it seems doubtful whether this
is ever likely to be followed more than that the tea leaves should be eaten

with
ins
enor
stung

in stock pursued in England, packets of the ground coffee being sold to
the consumer which may be years old, is far inferior to the continental
system of the consumer roasting and grinding his own coffee in small
quantities as required.

Structure of the Wood.—Wood white, moderately hard, close-grained
Pores very fine and extremely fine, medullary rays very fine, numerous

COIX
lachryma

Job's Tears.

Kurz in his
Kyaip
names C
BALAGH

FODDER.
1685

Fodder — Duthie says that in Balaghât in the Central Provinces, it is said to be used as fodder when in the young state Roxburgh, however, remarks that, owing to its coarse nature, cattle do not eat the grass

1686

Coix lachryma, Linn., Duthie, Fodder Grasses, 18

JOB'S TEARS

Syn — C. ARUNDINACEA, Lamk., LITHAGROSTIS, LACHRYMA JOBI, Garin

Vern — A recent correspondence between the Government of India and the

KAREN *Audnia thia* or *kuahati* (the black form), *su isa* (the white), *ka ke si* or *kasi* (collective or generic name), NAGA HILLS, Mung, MANIPUR, *Kirindi-mana*, SING, *Ee jin, ee-yin*, a name used in China and Malacca

hence according to them *Inula* and not *Coix* would be the true Job's

ever,
lath
ow
the
ther
late
and
the

Pl,

357, Dymock, *Mat Med W Ind*, 2nd Ed, 853; Balfour, *Cycl Ind t*
Hooker & Plim Jour, II, 259

Hah tat — Mat

Job's Tears.

COIX
lachryma.

tea, and appear to occur at higher altitudes. They are also more stunted in growth, and the involucre (or shell around the grain) is looser, softer, and apparently always furrowed—at least this is so with all the cultivated forms.

THE FORMS OF JOB'S TEARS.—There are three or four well marked forms of Job's Tears met with in India, which differ from each other in shape, colour, and degree of hardness, and in the presence or absence of

FORMS OF.
1687

only smooth and polished

The writer has had the pleasure to examine a large collection of samples made in Burma and Assam, and would offer the following remarks regarding these

1st.—The cylindrical form—met with in the Pegu Divisions, Pegu, Hanthawaddy and

this berry." It would appear, therefore, that the cylindrical grain may occur in the Miri country, but up to date (in connection with the present enquiry) no information corroborative of this fact has been received from Assam, and the plant does not appear to occur in any other part of India, so that it may safely be viewed as a native of Burma, and possibly distributed into the mountain tracts of Upper Assam and Cachar. The cylindrical grain is always of a white colour, smooth, polished, not furrowed, but constricted towards both extremities and whether wild or cultivated, is collected for ornamental purposes only, and not as an article of food.

2nd.—Of the pear-shaped form there are numerous sorts, varying in size and colour—some pale and bluish white, others grey, yellow, or brown-black. They are often constricted at the base into a disk-like annulus,

and so thick and hard that it can scarcely be broken. The cultivated

It seems probable these belong to a different plant from the forms described above.

COIX
lachryma

Job's Tears.

FORMS OF.

It is somewhat remarkable that in all the cultivated forms the shell is

the means of recording the vernacular names that are in use with reference to the various wild and cultivated plants.

PEGU DIVISION.

BURMA.
Pegu.
1688

In the Pegu District five forms of a large pear-shaped kind known as *cheik* or *kyeikthi* which grow for food or for ornamental purposes, white, the other brown grey.

A brown edible form is cultivated—a polished grain with the characteristic form of *var stenocarpa*, long, thin, and slightly swollen, with a bluish green tinge.

cylindrical from the pear-shaped forms. The best quality is said to come from the upper valley of the Pegu river.

In *Hanthawaddy District* some seven or eight forms exist in a wild state or are cultivated. One only is grown as an article of food, namely, a slaty brown irregular grain, of a dull colour, furrowed, and with an annulus. This is found only on the plains, is called *Kyeikthi*, and is sold for 8 annas a basket. All the others are wild or cultivated, but collected purely for ornamental purposes. One is a medium-sized steel grey seed, smooth, shining, and pear-shaped. Three are pinkish brown, small, of the flattened spheroidal form, and the most perfect beads in the whole collection of Coix seeds before the writer. These have been lettered B D and C.

These are said to be very command, polished, with impossible imitations of the signment of above under Pegu, the sample marked G, agreeing with the so-called "male," and C with the "female" form.

In the *Prome District* both spherical and cylindrical forms are said to occur, wild and cultivated. Of the samples forwarded along with the writer, must be the longer form. The Deputy

Job's Tears.

COIX
lachryma.

FORMS OF,

Commissioner deals in his report with a much more extensive series than he has furnished samples of. He says the forms of Coix are known collectively by the name *Kyeikthi*. The cylindrical being *Kyeikshe* (literally, long *Kyeik*), of the globular form there are names to distinguish certain recognised types thus — *Kyeikphun*, white *Kyeik*, *Sakyeik*, edible *Kyeik*, *Pyaung*, or maize-like *Kyeik*, and *Kyeikm*, or red *Kyeik*.

In the *Tharrawaddy District* the Deputy Commissioner says that all the forms are known by the Burmese name *Kyeikthi*, but that a large round edible form is known to the Karens as *Bè*, and is cultivated, while another smaller round kind is known as the *Be-ma* (or female *Be*) and is collected for ornamental purposes. He further forwards a sample of the cylindrical grain, and says it is known as the *Be-kwa*.

ARAKAN DIVISION.

Arakan.
1689

In the *Akyab District* the pear-shaped form is both wild and cultivated. From the town of Akyab, the Deputy Commissioner has furnished three samples of the wild plant, the seeds being smooth, polished, and very hard, especially a brown form. He states that these forms grow in the low marshy lands and are not eaten. He, however, furnishes a sample of a cultivated form obtained from Myohaung—the largest Coix grain yet examined—which fully supports all that has been stated above. It is steel grey, deeply grooved, with a loose shell and pronounced basal swelling. The Deputy Commissioner describes this as “the cylindrical form,” but while it is certainly longer than the Akyab grain, it is not the cylindrical form (*var stenocarpa*) described above, but is a monster form of the ordinary cultivated pear-shaped grain.

In the *Kyaukse District* three forms of Coix occur—two wild and one cultivated. The writer has not seen any specimens of these, yet has no reason to doubt but that they would answer very much to the types described under Akyab. One of the wild forms is larger than the other and is known as *jaissee* or *kalinsee*, while the smaller form is the *chitsee*. The edible form is also known as *chitsee*, and is both eaten and made into beer.

TENASSERIM DIVISION

Tenasserim.
1690

In the *Amherst District* both the round and cylindrical forms are grown, the former being eaten, and the latter used for ornamenting ladies' dresses. A wild round form is said also to exist. Samples have not been communicated, but the Deputy Commissioner reports that both are known as *kyeit*.

In the *Shwe-gyan District* no form of Coix is known.

In the *Taung-ngu District* it is stated that the cylindrical form grows wild, while the globular is cultivated; both are known as *kyeit*: the former

wild, while Nos. 3, 5, 6, and 7 are used for ornamental purposes, and No. 4 is extensively eaten. It is worthy of note that of these samples only those cultivated, viz., Nos. 1, 2, and 4 have the shell or involucre furrowed—the others are smooth and shining.

(1) *Kaleik* is a dark brown or bluish black polished grain of the pear-shaped series.

COIX
lachryma

Job's Tears.

FORMS OF.

- (2) *Kaleik Kauk-nyin*, the same as the last so far as the appearance of the grain goes
- (3) *Kaleik S* — as "male"
- (4) *Kaleik F* grain in tation, is
- (5) *Kaleik Yingwe* — This is a very small form of the flattened spheroidal, grain of a dirty milky white colour, a little smaller than the Hanthawaddy sample marked D, but of the same shape. The seeds are less than a $\frac{1}{4}$ inch in diameter and not much more than half that size in thickness through the central perforation
- (6) *Kaleik Yaing*, the form of *stenocarpa* that has been described as
- (7)

of the steel grey whites are quite as large as No 7, but few of the straw

In the S cultivated, t Burmese as *kyettthulon* cylindrical the cylindrical and *Tabuse* the globular Both forms are extensively grown in the Shan States, where the cylindrical is sold for R1 a bushel and the globular from 4 to 6 annas

The following abstract of available information regarding Coix cultivation in Assam may be here given to complete this brief review of the subject —

Sir J D Hooker remarks "A great deal of Coix is cultivated in the Khásia hills, the shell of the cultivated sort is soft, and the kernel is sweet, whereas the wild Coix is so hard that it cannot be broken by the teeth, each plant branches two or three times from the base, and the produce

Six varieties of Job's Tears. The generic name is *Xanthoxylum*, and the varieties are as follows —

"*Sibu*" — The seed is of a bluish grey colour and pear-shaped in

Herengisa-si — Of the same colour as *Sibu*, but more shape. Hardly to be distinguished in fact, from *Sibu*, except in being set brown rain, with ly hard to

admit of its being used for ornamental purposes

"*Saniapre*" — Pear shaped in form resembling *Sibu*, but smaller in size. This dark brown regular grain looks at first sight remarkably like some of the forms of black rice. It is about the same size and is pointed at both extremities. It is considerably like an elongated caraway.

Job's Tears

COIX
lachryma.

FORMS OF

"*Kadatha*"—Almost globular in form, of a mottled brown and grey colour. The most marked peculiarity of this grain is that it is dark brown like the *Sipta* form in the lower half and yellow or straw-coloured in the upper.

"*Kasi*"—Globular in form of a light grey or yellow colour. This is the most common variety.

The Naga hill samples, examined by the writer, fully support the

Naga Hills

purposes. It may also be added that the average elevation of the Naga and Khasia hills may be put down at from 3 000 to 5 000 feet whereas the smooth-shelled forms are met with chiefly in the marshes of the plains of India and Burma. The white forms of the Khasia hills are harder, more polished and less furrowed than the cultivated white forms from any other part of India, but they still preserve the characters assigned collectively to the cultivated forms. From the Khasia and Jaintia hills two samples of Coix have been received both of the milky white kind. A large and a small grain from the latter resembles very much the small white grain obtained from Mergui (No 4 above) only

Khasia Hills,

s
e

The dark coloured forms are said to boil softer than the white and the smaller of the two white forms "is slightly better flavoured than the larger."

Food.—This curious grain might almost be said to be unknown to the natives of India generally, except as a weed of cultivation. To the hill tribes on the eastern frontier, however, it is an important article of food, with the Tankhul Nagas of Manipur it might, indeed, be almost described as the staple article of diet. In several districts of Burma it is also regularly grown as an article of food. Mason says the esculent Coix cultivated by the Red Karens is parched like Indian corn. Of the Bassein district Mr W. T. Hall (*Director of Land Records and Agriculture*) reports that it is sown in gardens, the crop ripening in November. The produce sells for Rs 2 to Rs 3 a bushel. That officer has also forwarded to the writer numerous reports received from the Commissioners of the various Divisions, from which the following account of the method of cultivation may be here reproduced.—The mode of cultivation is as follows—*1st*, before the seeds are put in the ground they are tied in a piece of cloth and watered every day for about 7 to 8 days, when whitish roots appear. They are then placed in the ground. In some cases the roots do not appear till 10 or 15 days. *2nd*, at the place where the plants are to be grown furrows are formed and the seeds are laid on the earth which is first mixed with cow's dung, afterwards the seeds are covered up with a little

FOOD
1692

ed leaves

considered

becomes

This will

cause the plants to yield another crop and thus to last much longer." Speaking of the cultivation pursued in Akjab the Deputy Commissioner writes (of the Myohang township) with reference to the form which he calls "the cylindrical," but which, according to the samples discussed above, is a large loose shelled grain of the pear-shaped series—

COIX
lachryma.

Job's Tears.

FORMS OF.

"The cylindrical is sown by the wild hill tribes on Kaing land or on the slopes of hills. They do not till the land for this purpose, the seeds are thrown broad-cast, and no care is taken of them. In times of scarcity of food the cylindrical are eaten, but now they are only used as ornaments for their dresses." The Deputy Commissioner of Kyaukpypu writes regarding a beautiful hard round form which is collected from the wild plant and used for ornamental purposes. Of the cultivated forms he says this is known as *Chitsee*. "It grows in June and July and dies in November and December. The plant is 4 or 5 feet high and like a reed." But a smaller, more delicate, variety is also cultivated, which he remarks is eaten and also used in the manufacture of the small beer known as *Khanag*. He adds, "The seed has to be cleaned and has the taste of maize." Of the two kinds grown he says, "The plants, however, differ widely in other respects, and I am unable to say if they belong to the same variety or not."

CHARACTER OF THE EDIBLE GRAIN—On breaking the outer shell, a cowry-shaped grain is obtained which, Professor Church says, bears on being cleaned the proportion of 1 to 4 to the total weight of the unhusked article. The Professor gives the following analysis—

Composition of Job's Tears (Husked)

	In 100 parts	In 1 lb
Water	13.2	2.02 49 grs
Albuminoids	18.7	2.43 43 "
Starch	58.3	9.143 "
Oil	5.2	0.364 "
Fibre	1.5	0.105 "
Ash	2.1	0.147 "

"The nutrient-ratio is here 1 : 3.8, the nutrient value 89." From these facts it may be inferred that the grain is not likely to prove of greater economic value than it at present is to the poor hill tribes who

sequence of
e will grow
and coarser
it is sold for
for hospital

patients in China. It is worthy of note, however, that in the extensive series of cultivated forms which exist, and the occurrence of a long list of names for the plant and grain in nearly every vernacular language of India and Burma, an indication is given of an ancient cultivation

and through the Malayan
If this,
possible
abandoned
has
partly
also be
region
is con-
n occur
e culti-

anywhere in India proper
vated anywhere in India at
Hills some five or six forms of the loose-shelled and furrowed and are grown, but the plant is said to be rarely, if ever, met with in the wild state, while the cylindrical is reported as wild in the Naga Hills but never

COLA
acuminata.

Job's Tears; Cola Nut

DOMESTIC

identified as *Polytoca Wallichiana*, but have since been determined as *C. lachryma* var. *stenocarpa*. Subsequently, numerous samples of Job's tears, from every district in Burma, were obtained, and it has transpired

PRICE.
1700

form would afford the manufacturer of laces, &c., a choice of two forms which might be elegantly combined

PRICE OF COIX GRAIN—This has been variously estimated at from 8 annas to R4 a basket, but it seems probable that were a regular de-

established, which would pro-
It would have, however, to be
be cultivated without losing
as decorative articles The

be the produce
of *lachryma*),
il) to cultivate
cultivation of
well, and were
d above, their
and, the price
being cultiva-
the plants
also in lower

Nepal, to such an extent that no fears need be entertained of the demand, for some time to come, exceeding the supply.

Coke, see Coal

COLA, Schott; Gen Pl, I, 218.

1701

Cola acuminata, R Br, STERCULIACEÆ

Syn.—*STERCULIA ACUMINATA*, Beauv

References.—*New Reports*, 1880, p. 14, 1881, p. 10; *Christy New Com-
mercial Plants*, No. 8, p. 5; *Treasury of Botany*, p. 311; *Smith Dict
Econ Pl*, p. 127; *Balfour, Cycl of India*, U S Disp, 15th Ed,
p. 1754; *Pharmaceutical Society Journals*

It has been experimentally intro-
rom
-dily
ma

Cacao) It has been said the beverage made with Cola paste is ten
times more tonic than chocolate made with cacao. The reputation of
is such that it is
ities of the world as

and Schlagden-

There are many tracts of country in India that seem likely to prove
suitable to Cola cultivation, and doubtless this subject will in the future
receive a greater degree of attention than it has as yet obtained from the
Indian planters

C. 1701

Official Colchicum

COLCHICUM
autumnale.

COLCHICUM, Linn., Gen. Pl., III, 821.

Colchicum autumnale, Linn.; LILIACEÆ

OFFICIAL COLCHICUM, MEADOW SAFFRON OR AUTUMN CROCUS.

1702

References.—*Pharm Ind*, 243; *Fleck & Hanb*, *Pharmacog*, 699; *U. S. Dispens*, 15th Ed., 469, 470; *Bentley & Trim*, *Med. Pl.*, 288; *at Ind. Preface*, xxi; *Book of Pharmacy*, 1874, p. 103; *clap*, 808; *Balfour*, *Morton*, *Cyclop. Agri*,

490

Habitat.—The plant grows in the meadows throughout Europe. Attempts have been frequently made to introduce several species into India, but with very little success. Mr. Baden Powell says that in the Panjáb a species of Colchicum is known as *Harantutiya*.

The fresh corms and the seeds of Colchicum are official.

C. sp.

1703

Ve *suringan*, HIND,

M *U. S. Dispens*,

variety and the bitter, but adds a third form or rather substitute which he says is the sliced bulbs of *Narcissus tazetta*, which are imported from

chicum variegatum, Linn., a native of the Levant and not known to be found in Kashmir or Persia. Planchon in his account of *Sérinjan* gives a figure of *C. variegatum*, Linn., in the *Bot. Mag.*, t. 1028

References.—*Royle*, *III. Him. Bot.*, 365; *Baden Powell*, *Pb. Pr.*, 381;

Journal, April 1871

HISTORY.
1704

C. 1704

COLDENIA
procumbens.

The Surinjan; Trailing Coldenia.

HISTORY.

Mir Muhammad Husain tells us in his *Makhzan* that the white is the best, and that it is not bitter, next the yellow, both may be used internally, the

MEDICINE
1705

...na aperient, especially useful in gout, rheumatism, liver, and spleen. In gout they combine it with aloes, with ginger and pepper it is lauded as an aphrodisiac, a paste made of the bitter kind with saffron and eggs is applied to rheumatic and other swellings, the powdered root is sprinkled on wounds to promote cicatrization. Two kinds of *Surinjan* are met with in Indian shops, bitter and sweet. European physicians in India who have tried the drug consider the sweet *Hermodactyl* to be inert or nearly so, and the bitter to have properties similar to *Colchicum*.

Medicine.—§ "Purgative, diuretic, sedative, cholagogue, doses 2 to 8 grains, use—
heart dise
constipation
Lall, 1st c
Fubbulpori
the latter
Assistant-Surgeon, Meerut
hitis,
una
ary,
ter,
dose,

Colchicum luteum, Baker, according to Aitchison, in a note furnished to the writer, "occurs in early Spring in the Panjáb from Campbellpore, across to Abbottabad, the Gullies, at Murree, and in Kashmir extending to Zoja pass.

Probably it is the root of this that is *Haran-tutiya*. But the root of *Merendera Persica, Bois* (*Syn Aitchisonii, Hooker*) may be mixed with it.

SUBSTITUTES.
1705

SUBSTITUTE OF *SURINJAN*—Dr Dymock says that the sliced bulbs of the true *Narcissus* (*N tazetta*) which are imported into India from Persia as a substitute for *Surinjan* are easily recognisable. He remarks this drug may be at once detected by its larger size and tunicated structure. The taste is bitter and acrid the substance amylaceous and very similar to that of the *Hermodactyl*. It is used as an external application and, according to the author of the *Makhzan*, has properties very similar to those of *surinjan-i-talkh*. Value, annas 3 per lb.

COLDENIA, Linn , Gen Pl, II, 841.

1707

Coldenia procumbens, Linn , Fl Br Ind, IV, 144; BORAGINÆÆ

TRAILING COLDENIA

Vern — *T. Surinjan* ... *R. J. SIND, Tru*

Habitat. — A small annual weed, usually quite flat, common throughout tropical India, it generally grows on dry rice-fields during the cold season, disappearing about the beginning of the periodical rains. It is common in the hot dry parts of Ceylon. Distributed to Asia, Africa, Australia, and America.

Colebrookia, Country Borage.

COLEUS
aromaticus.

Medicine.—As a medicine, equal parts of the dry PLANT and fennugreek SEEDS rubbed to a fine powder, and applied warm to boils quickly brings them to suppuration (*Amst*). The fresh leaves, ground up, are applied to rheumatic swellings (*Murray*)

MEDICINE.
Plants
1708

COLEBROOKIA, Sm, Gen Pl, II, 1180

A Himalayan genus, comprising only one species, and that one of the commonest and most abundant plants in the Lower Himalaya and mountains of India, ascending to 4,000 feet in altitude

Leaves
1709
1710
1711

Colebrookia oppositifolia, Sm, Fl Br Ind, IV, 642, LABIATÆ

Vern — *Pansra*, HIND, *Shakarāna*, *phisbekkar*, *duss*, *samprā*, *sādli*,

References — *Roxb*, Fl Ind, Ed C B C, 467, *Voigt Hort Sub Cal*,

Habitat — A shrub with grey bark, common on the outer Himalaya,

Mysore It is now viewed as not even worthy of separate recognition as a variety

Medicine.—The leaves are applied to wounds and bruises (*Stewart*) "The down is used by the Paharias to extract worms from bad sores on the legs (*Gamble*) A preparation from the root is used by the Santāls in epilepsy (*Campbell*)

Fodder—The leaves are used as fodder for cattle (*Balfour*)

Structure of the Wood—Greyish white, moderately hard, close-grained Weight 46lb per cubic foot It is used for gunpowder charcoal

MEDICINE.
1712

FODDER
1713
TIMBER
1714

COLESEED or COLLARD, see *Brassica campestris*, Linn, var. *Napus*, B No 810

COLEUS, Lour, Gen Pl, II, 1176

Coleus aromaticus, Benth, Fl Br Ind, IV, 625, LABIATÆ

COUNTRY BORAGE

Syn — *C. AROCHNIDICUS* Lour, *Voigt Hort Sub Cal*, 450; *PLECTRANTHUS AROMATICUS*, *Roxb*, Fl Ind, Ed C B C, 466

Vern — *Pathor chur*, HIND, *Pâtér chur*, BENG, *Páthor chur pathár chár*, *owa*, BOMB, *Pathur chár*, MAR, *Páshana dhedi*, SANS In *Flora Andhrica*, *karpáru-vallu* is applied to this plant, but Dr Moodeen Sheriff is of opinion, that the name is more in use for *Arisochilus carnosus*, than any other name

References — *Dalé & Gibz*, Bomb Fl Supp, 66, *Pharm Ind*, 168; *Moodeen Sheriff*, Supp Pharm Ind, 114 51, *U C. Dutt, Stat Med Hind*, 313, *Dymock Nat Med W Ind*, 505 *Drury*, U Pl, 153; *Lisbon*, U Pl Bomb, 169, *Royle*, Ill Him Bot, I, 303, *Balfour*, Cyclop

1715

COLLOCALIA.

Country Borage; Birds' Nests.

MEDICINE.
Plant
1716

Habitat.—A native of the Moluccas, cultivated in gardens throughout India; has a pleasant aromatic odour and pungent taste

Medicine.—The PLANT "is employed in Cochinchina, according to

other suitable vehicle In his own practice he observed it produce so

who
ough
'pri-
pro-
and
has
the
in a

Juice.
1717

much larger quantity than is usual in Bombay.

Special Opinions.—§ "Expressed JUICE of the LEAVES is considered as an anodyne and astringent, and applied over and around the eyelids, in cases of conjunctivitis" (Anund Chunder Mookerjee, Assistant Surgeon, Noakhally) "Said by Sanskrit writers to have a specific action on the bladder and to be useful in urinary diseases, vaginal discharges, &c." (U C N S C, Calcutta, 1866, p. 100) "leaf in chronic dys-

FOOD.
Plant.
1718

pep
able
say
fragrant, they are frequently eaten with bread and butter, also bruised and put into country beer, cool tankards, &c., being an excellent substitute for Borage"

ns an agree-
Roxburgh
delightfully

1719

Coleus barbatus, Benth, *Fl Br Ind*, IV., 625; Wight, *It*, t. 1432.

Vern.—*Garmal*, BOMB

References.—Voigt, *Hort. Sub Cal.*, 449; Thwaites, *En Ceylon Fl*, 238, Dalz & Gibs, *Bomb Fl*, 205, O'Shaughnessy, *Beng Dispens*, 401, Drury, *U. Pl*, 154, Lisboa, *U. Pl. Bomb*, 168, Royle, *Ill Him Bot.*, I, 101, 103, Balfour, *Cyclop.*

Habitat.—A native of the Deccan in C. at Bombay and of the sub-
3,000 feet.
it is also
was intro-
curiously

FOOD.
1721

at Bombay for the roots, which are pickled (J. Graham) (Drury).
Lisboa says that the pickled root is much used by the Gujaratis.

1721

COLLOCALIA.

It would appear that there are two or three species of Swiftlet which form edible nests. Dr. Jerdon is of opinion that the best nests are obtained from

COLLOCALIA

nidifica.

Eagle's Bird's Nests.

is explored, many more nest-yielding caves will be found. At our present knowledge is derived from the Malays, who, through fear of the Andamanese, did not dare to search the interior. The explorations should be confined to hilly country, where the crystalline limestone formation predominates."

NICOBAR
ISLANDS
1724

NICOBAR ISLANDS.—Mr. deRapstorff, in his official report of the Nicobar Edible Bird's Nests, remarks: "The best nests I found at Katchall. They were entirely snow-white, and of the best quality. The next best quality I have got were from the Island of Bombaina. This island I have not personally visited," but he adds, the nests from it "are quite free from foreign matter, and have not the same snow-white beautiful colour as the ones from Katchall." The nests from Katchall are round and egg-formed, while those from Bombaina are long, like the section of an orange."

"The third quality I have is from Sambelong. This is white enough, but intermixed with little weeds or graminaceous stalks. These nests are of good quality, but need cleaning to separate the stalks. The fourth quality I got from the Car Nicobar from a cave in 'David's Bay' in deRapstorff's time is the north end of this island. These nests were entirely worthless for purposes of trade, consisting of the little weeds which are mentioned in the nests from Sambelong. These nests are, however, fastened together by exactly the same glutinous matter which forms the nests first mentioned."

"The Island of Katchall is mostly formed of coral limestone, and sandstone in all different stages, old, flinty, and yet forming. The island has gone through a series of volcanic revolutions and convulsions, and presents a very pretty landscape, many reefs and terraces, ravines and craters extending far under the earth. In these craters dwell the bats and the little swallows. The light of the sun never shines there. The ground is soft to tread on. If you lift it up and inspect it under the torch-light it is seen to contain the wings of the nesters, that have taken a prey to the bats, glimmering like a to wood like rubble; the soil is moss, spread in a hole, and you see the little long-nosed extremities of the swallows together with the feathers fallen from the roosting birds. This is the ground. The swallows' nests are not easily seen, but if you lift the torch up to the arched roof by the side of the albatross-like transparent calcareous white like these, the black head of the little mother appears out of her white hole nest."

ETNA
1725

IN ETNA.—Mason says of *C. fasciata* (*C. fasciata*): "This particular species occurs abundantly on parts of the coast of the Malayan Peninsula in the Nicobar Islands, and the Margin Archipelago, and so high as on certain rocky isles off the southern portion of the coast of America, where the nests are annually gathered, and exported to China. From all this range of coast we have seen no other species than *fasciata*, nor does it appear that any other has been observed; and I have examined a multitude both of the adults and of the young taken from the nests, collected in the Nicobars and preserved in spirits, all of which were of the same species. Still, what appears to be *C. nidifica* inhabits the mountains like in the interior of India, though hitherto unobserved upon the coast; and it is worthy to notice that *C. fasciata* does not appear to have been hitherto remarked inland in this country." (See note quoted by Mason.)

"It may be here added that *C. fasciata* is constantly seen inland in these provinces. The Karens in the valley of the Tenasserim in the latitude of Tavoy are well acquainted with the bird, and they say it crosses the mountains to and from the interior every year. That it is the same species there can be no doubt, for the Karen name of the bird is 'the white swallow,' from its white belly."

C. 1725

Edible Birds' Nests.

COLLOCALIA
nidifica.

In the Burma Gazetteer a list of the birds found in the province is given, and among these are included three species of Collocalia, viz, *C. innominata*, *Hume*, *C. spodiopygia*, *Peale*, and *C. linchi*, *Horsf*

MALABAR COAST—Very little of a definite nature can be learned regarding the edible swallows' nests collected on the western coast. They are said to be found in Dainger, North Kanara, and elsewhere.

MALABAR
COAST
1726

COLLECTION,
1727

tend that they were made of a sea weed which the bird collected for the purpose and chemically changed in some mysterious way. *Ure (Arts, Manufactures, and Mines)* says "The nests are made of a particular species of sea weed which the bird macerates and bruises before it employs the material in layers so as to form the whitish gelatinous cup-shaped nests so much prized as restoratives and delicacies by the Chinese." On the other hand, many recent writers discredit this theory and believe that the gelatinous material is either the natural saliva of the bird or a substance brought up from the stomach for the purpose and derived from the natural food of the swift, viz, insects. In support of this opinion they point out that the better qualities of the nests are found in caves far removed from the sea. Some of the nesting caves of Borneo are 140 miles from the sea. Mr deRoepstorff points out that there are no edible nests in the Nicobar settlement, but a few miles off in a richer tract of country where insect life abounds they are plentiful. "It is thus," he says, "in places

fresh, but when old brownish

Mr Portman remarks "The this matter, which resembles isinus) resembling Carrageen, an a weed, but have never seen the

Another theory is that the bird excretes this matter from his own throat during the breeding season"

"I am unable to

takes about a month and the so, the collectors should wait go out again, taking care to

observe exactly the same order in their rounds. The nests may be col-

C. 1727

COLLOCALIA
nidifica.

Edible Birds' Nests

COLLECTION

lected until the commencement of the rains, when the collection should cease, and the birds be left to breed. Although the great demand is for the white nests, still it may be remarked that the *fucus* attachments of the grass nests, and the old nests gathered in the November cleaning, may be sold locally at Rs per seer, and should, therefore, be collected. Each collection averages about 52lb of nests." He then proceeds to state

and
and
and
with
hem

carefully in their bag, from which, at the end of the work, they are transferred to a box provided with a lock

"The greatest care is necessary in detaching the nests from the caves, that they should not be broken or soiled. After being brought into the
om the

Cooking
Nests.
1728

to pieces and cleaned. After this they are boiled in clear chicken-broth until dissolved, a process occupying about two hours longer. The usual allowance is one nest (value Rs1) to a teacupful of soup. Any clear soup
ard ng
namen

TRADE.
1729

who reside in Rangoon. They recognise three classes:—

'No 1, large, pure, white nests, averaging from Rs110—115 per viss=

3¹/₂lb,
No 2, clean, but slightly coloured nests averaging from Rs100—140

a viss

Canton,
ise of the
e second
the more
Japanese

do not use the nests but they prepare from a sea-weed an artificial nest called *Dischin-schan*, which they export to China. Of the Ratnagiri district it is stated the right to collect nests is farmed out to Goanese, and fetches about Rs28½ a year. The Andaman contractor used to pay Rs3,000, but last year, owing to the contractor having thrown up his contract, the Government worked the nesting and realized Rs4,900.

GUANO IN THE SWALLOW CAVES

GUANO.
1730

An inquiry was instituted into this subject, and Mr deRoepstorff
I am certain that
least one lakh of
rupees per annum. This opinion was expressed regarding the Nicobar
islands only, so that if to this be added the possible supply from the
Andaman Islands, there would appear to be no reason why India might

Kachú or Taro

COLOCASIA
antiquorum.

not at least meet all its own demands for guano manure if not open up
an export trade in the article.

Collodion, see under Gossypium

COLOCASIA, *Schott, Gen Pl, III, 974*

1731

[*Wight, Ic, t 786, AROIDEÆ*

Colocasia antiquorum, Schott, DC, Mono Phanerog, II, 491,

1732

TARO, EDDOFS, SCRATCH COCO, EGYPTIAN ARUM, COCO, KOPEH
Sometimes but incorrectly called YAM

Syn — ARUM COLOCASIA *Willd, Roxb, Fl Ind, Ed C B C, 624*

Vc " " " "

207

Habitat — Wild over the greater part of tropical India, and also culti-
vated throughout India on account of its corms, which are used as an
It is grown at places

COLOCASIA
antiquorum.

The Kachu or Taro.

floras of the South of Asia, we cannot doubt that this plant is wild in India, as Roxburgh formerly, and Wight and others have more recently asserted likewise in Ceylon, Sumatra, and several islands of the Malay Archipelago."

Engler (in *DC, Mono Phanerogm, vol II*) describes some seven varieties of this plant, three of which are apparently met with in India —

α typica, Wight, *l.c.*, t 786. *Arum colocasia*, Roxb *Fl Ind*, Ed

f. 1, cultivated form

4. *nymphæifolia* (Arum nymphæifolium, Roxb, Fl Ind, El. C.B.C.,
for IV col + Ia + 286 Ph. Ind. Fl. I VI + 1 Th. 1 + the *Sorakichu* of

larger than any of the varieties of *Colocasia* "var. *typica* above), "yet the leaves are narrow in proportion to their breadth". The only good character by which to know this form "is the shortness of the club of the spadix." "Every part of this plant is eaten by the Hindus."

A good deal has been written regarding the cultivated species of *Colocasia*, but it has been found impossible to discover what species, still less which varieties are alluded to. On this account it has been deemed desirable to compile the economic information here given from such authors as could be depended on for the accuracy of their general information, and to thus leave for future research a more detailed description than will be found here.

The following facts seem to refer to *var* typica

The following facts seem to refer to *var typica* Medicine.—The pressed juice of the petioles is styptic, and may be used to arrest arterial hemorrhage. Dr Bholanath Bose reports very highly in favour of this property, and states that the wound heals by first intention after its application (*Pharm Ina*). It is sometimes used in emache and otorrhœa, and also as an external stimulant and rubefacient by the natives.

Special Opinions.—“The juice expressed from the leaf stalks of the
of inflamed glands
of alo-
es and
wasps
e seen
o fresh
than a

C. 1734

MEDICINE
I733

FOOD
I734

The Bish Kachú.

COLOCASIA
virosa.

spinach, but, like the root, they require to be well cooked in order to destroy the acidity peculiar to Aroids. A considerable number of

FOOD

carrot-shaped, often weighing several pounds, and forms an important article of food among the lower classes, where quantity and not quality is a desideratum. It is usually served fried in *ghí* or boiled and pounded into a paste, and also in curries. There are varieties that are very small, hardly weighing more than a quarter of a pound." In the Manual of Coimbatore it is stated that the corms (apparently of *var. nymphaeifolia*) often weigh as much as 70 to 80 lb each, and that an acre will yield 250 maunds (of 25 lb), worth 12 annas a maund. The tubers are used by the natives of Bombay in curries, &c. They form the common food of the inhabitants of Travancore. The Malays hold it in high estimation (*Balfour*).

§ "Is considered very nutritious by the natives, who use it in their curries" (*Honorary Surgeon P. Kinsley, Chicacole, Madras*).

Colocasia cucullata, Schott

1735

Syn. for *ALOCASIA CUCULLATA*, Schott

C. indica, Engl., DC, Mono Phanerog., II, 494.

1736

Syn. for *ALOCASIA INDICA*, Schott, which see, A. 809

This plant is said to be specially cultivated in Brazil for its esculent stems and small pendulous tubers. It is known as *Man saru* in Orissa, and is there used in the treatment of piles.

C. macrorrhiza, Schott

1737

Syn. for *ALOCASIA MACRORRHIZA*, Schott

A species met with in Eastern Bengal and Siam, and also in Ceylon.

kappé. Ainslie (*Nat. Ind., II, 463*) gives its Chinese name as *dea-view*

and the name of the root as *da-wei*.

krit

poes

prac

rubbed on the head, sometimes cures intermittent fevers after every other remedy has failed." The active principle is very volatile, so much so that by the application of heat or by simple drying, the roots become innocuous.

C. virosa, Kunth, DC Mono Phanerog., II, 495, Roxb., Fl. Ind., Ed. C.B.C., 632 (under calla)

1738

Vern.—Bish Kachú

This plant, which is a native of the Lower Provinces, is the only member of the genus which the natives of India regard as poisonous. It is sometimes used medicinally, but is never eaten.

C. 1738

COLOCASIA
VIROSA,

Poisonous Properties of Aroids

CHEMISTRY
1739

Chemistry—Through the kindness of Messrs Pedler and Warden (*Professors of Chemistry in the Calcutta University*), the writer has had the pleasure to receive an advance copy of their paper* on the chemical properties and medicinal uses of the species which, by the early botanists, were all treated as belonging been thrown into some half a paper was to investigate the and the enquiry was suggested on receiving from the Civil Surgeon Dibrugarh "some portions of raw *Bish Kachu* tubers and leaves with the following statement 'A cooly woman administered some of the fried *kachu* to another sick cooly on the same garden, but the man, experiencing a burning sensation in his mouth, instantly spat it out. A pig ate what was so thrown away and died in an hour. A second pig was experimented on with some of the same stuff, and fatal results also supervened.' During the course of the same year a second case of poisoning by *kachu* was referred to the Chemical Examiner's Department, in this case slices of *kachu* tubers were introduced into a jar containing 'goor'. The

writers on economic botany say that the *bish kachu* is *Colocasia virosa*, and accepting this to have been, in all probability, the plant Pedler and

holic extract was prepared and found to have no poisonous effect. The same result followed on the administration of a distillate which was found to have no acid taste, and, as with many other vegetable substances distilled with water, it was found to contain a trace of hydrocyanic acid. "It is possible, however that certain varieties of *ARUM* may contain a larger amount of prussic acid, as, for example, the *A. seguinum* of the West Indies, which is stated to furnish a juice, two drachms of which has proved fatal in a few hours. The tubers left in the retort after distillation with water were still physiologically active, indicating that the active principle was not dissipated by mere boiling with water. Natives, in using *ARUM* for culinary purposes, frequently add an acid vegetable or fruit such as tamarind. We tried the action of certain acids on the fresh tubers and ascertained that boiling with water acidulated with hydrochloric acid when a fragment ted in a similar l, was very much uce any decided idicated the presence of a large amount of potassium and magnesium, calcium was also present, but we failed to obtain indications of sodium. The acids consisted of carbonic, phosphoric, hydrochloric, with traces of sulphuric acid. We also obtained from the dried tubers very marked quantities of potassic nitrate, so that when they had been incinerated they behaved very like

*See *Journal Asiatic Soc. Beng.*, LVII, Pt. II, No. 1 for 1883.

Poisonous Properties of Aroids.

COLOCASIA
VIROSA.

tinder, containing saltpetre. The examination of the ash thus failed to afford us any clue to the physiological action of the fresh tubers."

"It now occurred to us that possibly the painful effects produced by ARUM when in contact with the tongue, &c., might be due to mechanical

CHEMISTRY.

in cold diluted nitric or hydrochloric acid. There appears to us to be no reason to doubt the fact, that the whole of the physiological symptoms caused by ARUMS are due to these needle-shaped crystals of oxalate of lime, and that the symptoms are thus due to purely mechanical causes.

crystals on microscopic examination of dried ARUMS as we had found in the fresh tubers. We explain this apparent anomaly in the following simple manner. In the fresh condition of the tubers, the bundles of crystals of oxalate of lime are so closely packed more or less, the sharp points covering a in the drying of the tubers, the more or less parallel to one another a smaller area. And thus, instead of each crystal acting as a separate source of irritation and penetrating the tissues, the bundles act as a whole."

The poisonous effects of certain aroid tubers are therefore the result of mechanical irritation, similar to that produced by cowage (*Mucuna pruriens*) or to chopped hairs criminally mixed with food. It would be interesting to have this line of enquiry carried to its final issue in a systematic examination of all the plants, like rhubarb, which contain raphides. It is just possible that the crystals of oxalate of lime may

chemically analysed, but it may be said we have not advanced much nearer a full understanding of the chemistry of rhubarb connected with its physiological action than we were before. It is thus probable that the results of Pedler and Warden's analysis of the aroid tubers may have a more extended influence on therapeutic science than they seem to have realized.

Colocynth, see *Citrullus Colocynthis*, Schrad.; CUCURBITACEÆ.

Colombo (or Calumba) Root, see *Jateorhiza Calumba*.

Bladder Senna

[103. LEGUMINOSÆ

Colutea arborescens, Linn, var *nepalensis*, Fl Br Ind, II, [183, Legendre]

Syn —C NEPALENSIS, Sims, *Bot Mag*, t 2622

Vern.—*Braa* LADAK, AFGHANISTAN

References — Brandis *F*
Pb Pl, 64 O Shaug
Pharmacog, 221, U
and Drugs, Sind, 13
of Botany

Habitat — A shrub of the temperate west Himalaya, Kunawar, Tibet,

irgative, and are used to Europe as a substitute for tion They are admin s

tered in infusion or decoction in the dose of about half a pint (*U. S. Dispens., 1617*)

Colza Oil, see *Brassica campestris*, Linn var *Napus*, B No 810

COMBRETUM, *Linn*, *Gen Pl* 1, 688

[COMBRETACEÆ

Combretum decandrum, Roxb, Fl Br Ind, II, 452.

Vern — *Dhobela* CHINDWARA *Pink* GONDA, OUDH, *Ariketa* TEL.
Kalilara, NEPAL, *Pindik* LEPCHA

References—*Roxb Fl Ind Ed C B C*, Brandis *For Fl*, III.
Gaible, *List of Darjeeling Climbers &c*

Habitat.—Abundant in Bengal at altitudes up to 3000 feet. Very common in the North Deccan plateau in the North Western Provinces Tenasserim and the Andamans.

Is said to be used medicinally, but very little is known regarding the uses of the plant. The Santals, who call it *atena*, make baskets from its long thin stems (Campbell)

C. nanum, Ham , Fl Br Ind , II, 457

Vern — *Dant jalthi pharsta* N W P and Pa

Vern — *Dant ydhis pharsia* N W P and Pa
References — *Brandis For Fl*, 221, *Baden Powell Pb Pr* 350, *Royle*,
Ill Him Bot, 1, 209

Habitat.—A decumbent, low shrub of the Himálayan terai, from Sikkim to the Panjáb

Medicine—Mr. Baden Powell mentions this plant among his medicinal plants of the Panjáb

C. ovalifolium, *Roxb*

Vern.—Banda kattu tige yādala chettu, banda kōta, TEL. (the buffalo-calf tree)

A common climber throughout the Deccan Peninsula, probably eaten by buffalos

C. 1746

The Spider-words

COMMELINA communis.

COMBS, fans, brush-backs, and other smaller articles—Woods used for —

Adina cordifolia (combs)
Alangium Lamarckii (cattle-bells)
Albizia stipulata (cattle bells)
Artocarpus integrifolia (brush-backs)
Bauhinia Vahlii (umbrellas, rain-caps)
Buxus sempervirens (instruments, combs, small boxes).
Carissa diffusa (combs)
Casearia tomentosa (combs)
Chloroxylon Swietenia (picture-frames, brush backs).
Cordia allamanda (combs)

Crataeva religiosa (combs)
Elæodendron glaucum (combs, picture-frames)
Gardenia costata (combs)
G. latifolia (combs)
G. lucida (combs)
Gmelina arborea (picture frames)
Olea ferruginea (combs)
Platanus orientalis (pen cases)
Psidium Guava (instruments)
Pyrus Pashia (combs, tobacco-pipes)
Schrebera swietenoides (combs and weavers' beams)
Stephegyne parvifolia (combs)
Sterculia urens (guitars)

WOODS FOR COMBS, &c.
1747

COMMELINA, Linn, Gen Pl, III, 847.

1747

The genus of the Spider words is named in honour of the Dutch botanist Commelin.
Commelina benghalensis, Linn; DC, Mono, 159, Clarke, Comm et Cyrt, 14 Pl IV; Wight, Ic, t 2065, COMMELINACEÆ

1748

Vern — *Kanshura*, HIND *Kanchura kanuraka, kanshira, kashradm, kanchara, BENG, Kai arak, S'ANTAL, Chura, kanna, Pb, Khanna, SIND Kanchata* SANS, *Deya mainaireya* or *diya mentriya*, SING, *Ho tan tu*, CHINESE

References — Roxb, Fl Ind, Ed C B C, 57; Voigt, Hort Sub Cal, 676, Thwaites En Ceylon Pl, 321, Dals & Gibs Bomb Fl, 253, Stewart Pb Pl, 236, Atchison Cat Pb and Sind Pl, 148, Trimen Syst Cat, 95, DeCandolle, Mono Phanerogam, III, 159, Rev A Campbell, Descript Cat of the Pl Chutia Nagpur, U C Dutt, Mat Med Hind, 303, Murray Pl and Drugs, Sind, 22

Habitat — in the penins Deccan Dal Bombay Distributed to Butina, Malay, and China It also occurs large, and the everywhere in

Food — LEAVES eaten by the poor people as a pot-herb, especially in times of scarcity The fleshy rhizomes of some of the species of this genus contain much starch, mixed with mucilage, and are therefore wholesome food when cool Balfour says *C. polygama* (a name which would appear to be a synonym for *C. benghalensis*) is cultivated in China as a pot herb eaten in spring "The juice of the flower is used as a bluish pigment in painting upon transparencies" (Smith).

FOOD
Leaves
1749
Starch
1750
Pigment.
1751

C. communis, Linn, DC, Mono Phanerogam, III, 170.

1752

Vern — *Anna Boma, W'ek kyip* BURM Stewart says that this, as also *C. benghalensis* are in the Panjáb known as *Chura kanna* Balfour gives the following names *Kanang krai, Lunnu kaiti pillu, Tam, Venna devu kura niru kassutu, venna mudra, venna vedara, TEL, Vatsa priam, SANS*

It may be here recorded of the vernacular names given to this and, in fact, to all the species of *Commelina* that they require to be verified and assorted under the modern scientific names for the species of this genus.

COMMELINA
suffruticosa
The Spider worts

References — *Voigt, Hort Sub Cal 677 Dals & Gibs, Bomb Fl 251; Stewart Pb Pl 236, Aitchison Cat Pb and Sind Pl, 148, Balfour's Cyclopædia of India*

Habitat — A native of the hot damp regions of China and Japan. From Chittagong, plants are said to have been sent to the Botan

present been left in the present position

FOOD
Seeds.
1753
Leaves
1754

Food. (There are several other plants which were largely

calves when they wish to wean them from their milk. The leaves are eaten by the natives mixed with other greens"

[Com and Cirt Table I

1755

Commelina nudiflora, Linn, DC Mono, III, 144, C B Clarke's

Syn — *C. CAESPITOSA* Roxb, Fl Ind, Ed C B C, 58 *C. NUDIFLORA*, Linn, as described in Roxb, Fl Ind Ed C B C is *ANEILEMA NUDI FLORUM*, Linn, the *Kundali* of Bengal

Habitat — Frequent in Bengal, and distributed to Burma, Ceylon and the Malay, also to Africa, Madagascar, Mauritius, Sandwich Islands, and Australia, &c

Compare this with the remarks under *C. communis*, Linn., and *C. obliqua*, Ham

1756

C. obliqua, Ham., Clarke, p 19 pl IX

Syn — *C. COMMUNIS* Roxb, Fl Ind, Ed C B C, 57

Vern. — *Kanjurá kána* HIND *Jata kanchura*, *jata kanshira* BENG, *Korna kana* BIJNOR, *Kanjura* KUMAON

Habitat. — This species is common over the low moist parts of India and Bengal, and occurs on the lower parts of the hills. It is distributed to

MEDICINE
Root

1757

FOOD

Root

1758

1759

C. salicifolia, Roxb, Fl Ind, Ed C B C, p 58

Vern. — *Galap ppals languli*, SANS, *Pani kanchira*, BENG; *Galp pari*, HIND; *Bir kana arak*, SANTAL

References — *DeCandolle, Mono Phanerog, III, 157; U C Dutt Mat Med Hind, 300*

Habitat — Common in wet places in the peninsula of India, especially in Bengal, Coromandel and Bombay. Distributed to Burma

Fodder — Cattle are said to be fond of this plant

FODDER

1760

1761

C. scapiflora, Roxb, see *Anellema scapiflorum*, Wight A 1122

C. suffruticosa, Bl, DC, Mono Phanerog, III, 183

Vern. — *Dare orsa* SANTAL

Habitat — A native of Bengal

Medicine — The root is by the Santals applied to sores (Campbell)

MEDICINE

Root

1762

C 1762

Spotted Fern. CONNATUS

CONNAPUS
MONOCARPUS

Conch Shell, a species of *Turbonella*, see Shells, and Bands 2. 382.

Condiments, see Spices

Conessi Bark, see *Holarrhena andysentica*, Wall., APOTHECARIÆ.

CONGEA, Roxb., *Gen Pl*, II, 1159

Congea tomentosa, Roxb., *Fl Br Ind*, IV, 603, Wight, Ic., 1763

Vern.—*Tamarkanwe kajan* BURM

References.—*Kurz For Fl Burm*, II 256 Roscoe in *Poxb Fl Ind*, Ed C B C, 477

Habitat.—A large climber in Chittagong and Burma, distributed to Siam Roxburgh says it is found also in Coromandel where it flowers in the cold season the Chittagong plant flowering in March The *Flora of British India* describes a variety—*Azurea*—as cultivated in North India All the species of this elegant genus are characterised by their purple bracts

C. villosa, Wight, Ic., t 1479, fig B, *Fl Br Ind*, IV, 603 1764

A large climber of Pegu and Mergui, the leaves of which are used medicinally (*Mason, O'Shaughnessy, &c*)

CONIUM, Linn., *Gen Pl*, I, 883

Conium maculatum, Linn., *DC, Prodr*, IV 242; UMBELLIFERÆ
SPOTTED HEMLOCK, HEMLOCK, Eng, CIGUË, Fr, SCHIERLINGS,
Germ 1765

Vern.—*Shomkran*, ARAB, *Kirdamāna*, BOMB

References.—*Pharm Ind* 104 *Ainslie, Mat Ind, Preface p XII*, *O'Shaughnessy Beng Disp* 359 *Dymock Mat Med W Ind*, 2nd Ed 363 *Fluck & Ha b Pl irmacog*, 299, 301 *U S Dispens*, 15th Ed 194 484 *Bent & Trim, Mat Pl*, 118

Habitat.—Met with in Europe and temperate Asia, common in England

Medicine.—Although this drug is commonly used in Indian pharmacy, and largely imported no effort seems to have been made to cultivate the plant in the East. MEDICINE. 1766

CONNARUS, Linn., *Gen Pl*, I, 432, 1001

1767

1768

C. 1768

CONVOLVULUS
arvensis

Connarus, Deer's foot Bind weed.

References.—*Biddome, Fl Sylv App LXVII Wight and Arr* ;
Prod Fl Pen Ind Or, 143, *Thun, En Cey Pl*, 83 *Kurz, For*
Report . *Bomb G*., *AV*, 330. *Da's and G's, Bomb Fl*,
 53. *Rheede, Mal*, *VI*, t 24

Habitat.—A small tree or shrub of the Western Peninsula, from the
 Concan to Travancore, common on the Southern Ghats, very abundant
 in Ceylon. Flowers yellow, fruit long, bright red, the tree becoming very
 ornamental when in fruit

Oil.—The seeds yield an Oil.

Structure of the Wood.—The timber of this, as of most other species
 of the genus, is much valued for ornamental purposes.

L
 1700
 TIMBER
 1770
 1771

Connarus nitidus, Roxb, in Hort Beng, 49

References.—*Loigt, Hort Sub Cal* 15 *Gamble, Man Timb*, 114

Habitat.—Said to be found in Sylhet and British Burma.

Oil.—Dr McLelland says that in Rangoon the seeds of this plant yield
 a quantity of sweet oil. The name *C. nitidus* is not referred to by the
Flora of British India, but it may be presumed that the plant which
 yields the oil in question is *C. paniculatus*

OIL
 1772

1773 C paniculatus, Roxb, Fl Ind, Ed C B C., 505, Fl Br Ind, II, 52.

References.—*Kurz, For Fl Burm*, I, 32, *Gamble, Man Timb*, 114
Wight, Ill, t 64

Habitat.—Roxburgh, followed by Voigt and Kurz, describes this as "a
 large timber tree," but Hooker in the *Flora of British India* says it is "a
 large climber" met with in Sylhet and the Khasia hills, to Chittagong "

1774

C speciosus, McLell

Vern.—*Gwe-loak & don-kzet*, BURM

Habitat.—Said to be a large tree of Rangoon, Pegu, and Tenasserim.

Oil.—McLelland says that the seeds yield an abundance of sweet oil.

The above has been extracted from Dr Cooke's *Report on Oil*
Seeds. The name *C speciosus* McLell was taken apparently from *Bal*
four's Cyclophasia. It seems probable that the tree here alluded to is
C. gibbosus Hall—a large tree met with near Rangoon and in Tenas-
 serim, Penang and Singapore. The Burmese name *Gwe* (*Spondias*
mangifera) seems very near to the above

OIL
 1775

Structure of the Wood.—Balfour says of *C. speciosus* "It has a
 large, heavy, and strong timber, white coloured, adapted to every purpose
 of house-building "

TIMBER
 1776

Conocarpus acuminata, Roxb, see Anogeissus acuminata, III, 1,
 COMBRETACEÆ A. 1145

C. latifolia, Roxb, see Anogeissus latifolia, III, A. 1149

Construction and Railway purposes.—Timbers suitable for,
 see Cart and Carriage Building, C. 632.

CONVOLVULUS, Linn, Gen Pl., II., 874

1777

Convolvulus arvensis, Linn, Fl Br. Ind, IV., 219 CONVOLVULACEÆ

DEER'S FOOT BIND-WEED

Syn.—*C MALCOLMI*, *Roxb, Fl Ind, Ed C B C.*, 102

C. 1777

Scammony

CONVOLVULUS
Scammonia.

Vern — *Veri* (?) *harin-padi*, or by some writers *hiran paddi*, Pb, HIND, *Hirn-pug*, SIND

References — *Voigt, Hort Sub Cal*, 362, *Dals & Gibs, Bomb Fl* 163, *Stewart, Pb Pl*, 150, *Aitchison Cat Pb and Sind Pl*, 98, *O'Shaughnessy, Beng Dispens* 502, *Murray, Pl and Drugs, Sind*, 164, *Year Book Pharm*, 1879, 467, *Medical Top of Ajmir*, 150, *Baden Powell Pb Pr*, 367

Habitat — An abundant weed of cultivation all over the plains of the Panjab and Western India, from Kashmir to the Deccan, ascending to 10 000 feet in the Himalaya. Flowers large, deep rose coloured, sweetly scented. They appear in the cold season, very common on the black soil of Gujarat and the Deccan.

Medicine — The official *hiran paddi* (or *harin padi*) appears to be this plant. The roots possess cathartic properties. Murray says the roots are sometimes used by the Sindis as jalap.

Fodder — *Veri* is a dark green weed usually found in wheat fields. It is said to be greedily eaten by goats and cattle, and is gathered by village children as a fodder.

MEDICINE.
Root
1778
FODDER
1779

Convolvulus Batatas, Linn, see *Ipomœa Batatas*, Lamk

C. parviflorus, Vahl, *Fl Br Ind*, IV, 220

Vern — *Alaranyi*, TEL

A native of Assam, the Deccan Peninsula, and Ceylon, but largely cultivated throughout India.

1780

C. pentaphylla, Linn, see *Ipomœa pentaphylla*, Jacq.

C. pluricaulis, Choisy, *Fl Br Ind*, IV, 218

Vern — *Porprang*, *gorakh panw*, *baphalli dodak* Pb

References — *Stewart, Pb Pl*, 150, *Aitchison, Cat Pb and Sind Pl*, 99

Habitat — A common plant in many places throughout the plains of Panjab, Hindustan, and Behar.

Food and Fodder — "It is eaten by cattle and is reckoned cooling, and used as a vegetable or given in sherbet" (*Stewart*).

FOOD and
FODDER.
1782

C. reptans, Linn, see *Ipomœa aquatica*, Forsk.

C. Scammonia, Linn, *DC. Prodr*, IX, 412.

SCAMMONY

Vern — *Mahmudah* (?), *sakmunia*, Pb, *Sugmonia*, *sak mania*, HIND, SIND, ARAB, PERS

References — *Kurz, For Fl Burm*, II, 212, *DC Origin Cult Pharm Ind*, 153, *O'Shaughnessy, Beng Dispens*, 500, *Dymock, Mat Med*, Res, pens, drugs, 151,

Irvine Mar Med India, 14.

Habitat — A climbing perennial, native of Syria, Asia Minor, and Greece. Cultivated in some parts of India.

Gum resin — A gum resin imported into India. It is obtained by incision from the living root. It occurs in irregular pieces of an ash grey colour and rough exterior. When broken, it presents a resinous surface, and of a shining black colour when dry. Thin pieces are translucent and

GUM-RESIN.
1784

C. 1784

COPPICE or COPSE.

Plants for Coppicing

greenish It has a cheesy odour and flavour. The bazar *Scammony* in Bombay, Dr. Dymock states, is all false, and is made at Surat

[DC ; COMPOSITÆ.

Conyza alopecuroides, Lam. ; see *Pterocaulon alopecuroides*,

C. anthelmintica, Linn. ; see *Vernonia anthelmintica*, Willd.

C. balsamifera, Linn. ; see *Blumea balsamifera*, DC.

1785 Cooawanoo Oil.

This oil is said to be prepared from the Chelonian reptile *Casua olivacea*, Gray—see Turtles.

Cookia punctata, Hask. , see *Micromelum pubescens*, Blume, Var 1st, RUTACEÆ

1786 Copal Gum, or Gum Anime.

A hard transparent resinous substance obtained from a natural

much superior to that obtained from living trees. It occurs in immense masses, found buried in the sand, far away from any living trees, and chiefly in the coast sands There are other Copals sometimes met with Brazilian Copal is obtained from *Hymenæa Courbaril*. Madagascar Copal from *Trachylobium verrucosa*. West African Copal is furnished by *Guibourtia copalifera*, and Indian Copal from *Vateria indica*, which see The Australian and New Zealand Copal is the produce of *Dammara australis* (CONIFERÆ) This forms large solid masses, often found in places where the trees do not now occur, and in New Zealand is known as *Kauri*; and in European Commerce as DAMMAR or COWDIE PINE.

Copper, see Cuprum.

1787 Coppice or Copse—Plants suitable for—

The following, among many others, are plants specially mentioned as suitable for this purpose, but those given under Hedges and under Pollard may also be added. —

Acacia arabica
Acer Campbellii
Albizzia Lebbek
Anogeissus pendula.
Bauhinia Vahlil.
Carissa diffusa.
Castanopsis indica,
C. tribuloides
Casuarina equisetifolia.
Cedrela serrata.
C. Toona
Celtis australis
Dalbergia latifolia

Helicteres Isora
Hentiera littoralis
Lagerstroemia parviflora.
Lebdiereopsis orbicularis.
Morus montana.
Odina Wodier.
Pithecolobium dulce.
Populus euphratica.
Prosopis spicigera.
Quercus acuminata.
Q. semecarpifolia.
Streblus asper
Teucrium macrostachyum.

Coptis or Mishmi Teeta.

COPTIS
Teeta.

Copra or Khopra—The dried kernels of the cocoa-nut, see *Cocos nucifera*.

COPTIS, *Salisb.; Gen Pl, I, 8, 953*

1788

The name **COPTIS** has been given in allusion to the much cut leaves of the plants which have been referred to this genus

Coptis Teeta, *Wall, Fl Br Ind, I, 23, RANUNCULACEÆ*

1789

COPTIS OF GOLD THREAD, COPTIDIS RADIX, OR MISHMI TITA

VERN—*Tita*, Ass., *Mamira*, or *Mamiran* (DYMOCK) HIND, *Mahmira*, SIND, *Pita karosana* SING RICE says that *tita* is a corruption of *tikta*, SANS, "bitter"

References—*Voigt, Hort Sub Cal, 3 MacIsaac, Trans Med and*
DL C C III O III T R D L D A B J

Habitat—A small, stemless herb, with perennial root stock, met with in the temperate regions of the Mishmi Hills, east of Assam Cooper says that the plants grow on the ground among the moss around the stems of trees "From each root," he remarks, "springs a single stem, about four inches high, bearing three serrated leaves, attached to the head of the

to suggest that
early European
on the fact that

HISTORY.
1790

mahmira is the name of a drug used in Sind in the treatment of eye diseases, a purpose identical with that for which the *Majupás* was em

rium clears the sight, and as a snuff the brain, and that it relieves tooth-ache Internally it is given in jaundice, flatulence, and visceral obstructions' (*Mat Med West. Ind, 2nd Ed, 18*)

Dymock further remarks that two kinds of the drug are at the present day met with in Bombay The best quality is only about the thickness of a crow-quill or a little thicker, it is a yellowish rhizome, hav-

1791

C. 1791

Coptis or Mishmi Teeta.	COPTIS Teeta.
drugs imported into India, or may have been conveyed overland from the Indo-Chinese frontier to Chinese ports. Hence, as far as our present in-	HISTORY.

1794

sulting
almost
original
Pereira

may have been mistaken in referring the *Maupás* of the ancients to *Coptis Teeta*, since it is this imported Chinese drug that is the *mamíran* of Upper India. Further, it seems even probable that the knotty, yellow, often ramified rhizomes of *Picrothiza*—according to modern writers the spurious *mamíran* of the Indian bazars—may have been the drug originally so called, or at least been the Indian drug which most closely resembled the

of the wildest of our tribes. But there is nothing in all this to justify the inference that, in ancient times, there may have existed a much larger export

possible, however, that in later times the Chinese supply may have been

ne extent,
iters came
on, in his
year, col-
natives in
a yellow

1795

1796

watery juice, as every plant with a yellow juice seems to be by them considered a sovereign medicine, and all are called indiscriminately *mamíran*. He further states that the roots of *Geranium Wallichianum* were shown to him as a medicine called "*mamí-ran*."

It has been pointed out by chemists that both *Coptis* and *Berberis*

alter the same fashion as the *Maupás* of the ancients. But *berberine* is present in a great many other yellow and bitter substances, and it may therefore have been a mere coincidence (suggested by external appearances) that the root now called *mamíran* and the *Maupás* came to be used for the same purpose. Indeed, *Picrothiza*, on being chemically examined, may also be found to possess that alkaloid, since *berberine* is one of the most frequently met with of all the alkaloids present in vegetable substances. But even should it not possess *berberine*, that could scarcely

1797

C. 1797

Coptis or Mishmi Teeta.

be viewed as militating against its having been adopted as a substitute for a drug for which *Cephus* would have proved more suitable. At the same time the fact that *Cephus* is the treatment of eye affections is but a minor matter. The *Materna Medica* while the drug *Picrochuza* was known to the earliest Sanskrit writers. The late Dr. U. C.

Sanskrit writers, but it seems conclusively established that even the drug *Coptis Teeta* is but of modern introduction into India. The Muhammadans were so little familiar with *Picrohiza* that they frequently confused it with Hellebore, and may thus be readily believed to have given to *Picrohiza* or to *Coptis*, when separately presented to them, the name of *mamuran*—the name of a drug which either or both may possibly have closely resembled. The Hindus are uniformly precise and accurate in their information regarding *Picrohiza*, but say nothing of *Coptis*. The earliest writers on Indian *Materia Medica* who allude to *Coptis* attribute to the indigenous and imported Chinese drugs tonic properties of remedial value in the treatment of nervous diseases and in debility after fever, they rarely make any mention of its use as a collyrium in eye affections. The tonic properties of *Coptis* are possessed in a scarcely less degree by *Picrohiza*, and it may be concluded that Mir Muhammad Hussain's designation of *Coptis* as *mirzabadi* is a very probable one.

we neared the highest elevation, scattered trees and shrubs seemed to grow from a thick bed of dry moss and here, for the first time, I saw the *find* plant growing abundantly. The roots (from which, when brewed and steeped in hot water, the famous febrifuge is made) are embedded in moss. From each root springs a single stem, about four inches high, bearing three serrated leaves, attached to the head of the stalk-like elongated trifol. The Mishmees gather the roots towards the end of the rainy season, and carry them packed in tiny wicker work bamboo baskets to Sadya, where they are eagerly bought by Assamese and Bengali merchants. The Secretary of the Government of India, Mr. Dymock, has been to Sadya, and has estimated that a maund of a half is brought down in small open bamboo baskets, weighing about 1/2 a chattrak each, and is sold at the price at which the Government would buy it, but the small dealers are out of all retail price which the drug fetches. Dr. Dymock says of the Bombay supply: "Both

Coptis or Mishnu Teeta

COPTIS
Teeta.

kinds of the drug come from China *via* Singapore, in bulk. The first is worth Rs 3½ per lb, the second Rs 2. O'Shaughnessy says "Coptis Teeta has found its way through the drug-shops of Bengal, and is even occasionally exposed for sale in the Upper Provinces."

MEDICINE.

MEDICINE.
1799

Therapeutic and Chemical Properties—Coptis trifolia, a creeping

1800

its influence several patients, recovering from acute diseases, manifestly, and very rapidly, improved in strength. The dose was 5 to 10 grs of the powder, or an ounce of the infusion thrice daily." Dr K. L. De, CIE, says "In this indigenous article, though a costly one, we have an adequate substitute for Columba root, which it resembles not only in its medical effects but also in its physical properties. An essence of this drug has been recently brought forward for use by Messrs Bathgate and Co., of Calcutta."

1801

and Coptis d. the colouring matter in which the rhizome of Coptis

less than 8½ per cent, which is more than has been met with in any other

note
a b

1802

The Barberry.
Columba root
Hydrastis canadensis
Xanthorrhiza apifolia

Xanthoxylon fraxoneum.
Menispermum canadense.
Coscium fenestratum
Coptis trifolia

C. 1802

CORAL.

Teeta: Coral.

MEDICINE

"*Thalictrum foliolosum*, DC, common at Mussooree and throughout the temperate Himalaya at 5,000 to 8,000 feet, as well as on the Khásia hills, also affords a yellow root, which is exported from Kumáon under the

-*Coptis*
up in
emble
) See
stated
called

CULTIVA-
TION.

1803

mamian.

CULTIVATION OF TÍTA.—In concluding this brief account of *títa* it may be remarked that little or no difficulty would be experienced in cultivating the plant in many parts of India, but that up to the present date no attempt appears to have been made to do so, although the retail price paid for the drug would apparently justify the suggestion that it would be found a remunerative crop.

1804

CORAL.

1805

1806

Coral

CORAL.

may be described as covered externally by the outer fleshy wall and terminated

1806

or *sclerobasic coral*. Such a coral can therefore alone be produced in a compound organism. In the *sclerodermic coral* each polype has a complete skeleton of its own and may hence exist independently or be combined into a colony

1807

as a source of manure

Coral.

1808

CORAIL, *Fr*, KORALLEN, *Germ*, KORAALEN, *Dutch*, CORALLO, *It*, CORAL, *Port & Sp*, KORALLU, *Rus*, CORALLIUM, *Lat*, Κοραλλιον, *GREEK*

Vern — *Murjan munga* HIND, *Bekhi marjan* (fragments of red coral used medicinally) sang : marjan, *PS*, *Galli*, DEC, *Pavalam*, *nurai kai* (foam stone), TAM, *Pdguilam*, TEL, *Yidruma*, *prabala*, *pravdla*, SANS, *Biséd*, ARAB, *Murjan* or *merjan*, PERS, *Babalo*, SING,

R

CORAL.

Coral.

Habitat.—The Coral zone extends on either side of the Equator for about 1,800 miles. Mr. J. Murray, of the *Challenger* Expedition, has pointed out, however, that within this area the corals abound most on the

coral luxuriates requires to have a surface-water temperature of 70°F., and to never vary from this more than a limit of 12°F. There are a few

fluences that confines the coral regions but fixes each species

in which alone it is found to gr

corals, the ornamental corals occur, and luxuriating, under lower temperatures, they are found in tropical seas at much greater depths than the reef-forming. The latter class of corals grow between 5 and 30 fathoms of water. They are killed by exposure to the sun, and must therefore be below low-water level. On a land subsiding they will accordingly build vertically so as to preserve their favourite depth, and on a land ascending they will extend horizontally, advancing into the requisite depth of water as the older landward and exposed portions are killed by being carried above the level of the water. This was the theory established by Darwin, and universally accepted for a quarter of a century, the atolls being viewed as monuments erected by the Actinozoa to a vast Pacific continent which had gradually sunk beneath the ocean. While this may take place, a new school has advanced the theory that it is by no means essentially necessary that to construct an atoll, the island which it encircles need be subsiding. Growth is attributed to the food materials being most abundant along the face of the reef, the approaching water being richer than that within the lagoon. It is even further explained that the chemical action of the sea on the dead coral, thus excavating the shallow

face of the reef and the

explain the fact that o

the present face of the r-

which we have no evidence of its having the power to live, or then pit

... from a peduncle

und in
st. m. 1

it may
evidence

...identical
...power

...considerably above the

tionably rising, and reefs of various ages are now considerably above the level of the surrounding sea. The coral from these dead reefs, atolls

level of the sea, whereas a few miles seaward from these dead reefs, shoals are being formed around the islands of the Indian Ocean.

are being formed around the islands of the Indian Ocean.

A.—CORAL REEFS.

1. (b) Mr. J. Edgar Hoover testified that the coral reefs of

to the Agent at Calcutta, who has stated that the coral reefs are for Calcutta.

he past twenty

from the neces-

... solved, and the

... a source of

C 1800

C. 1809

REEFS.
1800

C. 1809

Coral Reefs.	CORAL.
lime, abundant fuel, and labour at command, there can be little doubt that Calcutta might be supplied with excellent lime at a comparatively small cost, and a useful and profitable occupation would be thus afforded for the convicts."	CORAL REEFS Andamans. 1810
In the Nicobar Islands upraised coral reefs are found on the coast of	Nicobar. 1811
(dam) across the Habb river, a thin bed composed of corals appears a few feet above the base of the Gáj group. This bed can be traced for many miles to the south. All the species of coral (five or six) are encrusting forms or small branching kinds. A <i>Pachysens</i> , or some closely allied form, and two or three species of <i>Hydnophora</i> , are specially common." So again near Nari he writes of coral beds: "The marly shales pass up into light yellow and brown limestone, with a coral zone abounding in	Sind. 1812
fessor P. Martin Duncan and W. Percy Sladen (see <i>Palæontologia</i> -91). But Mr. Fedden contigs the Gulf of Cutch from Nao off the coast, is fringed with much exposed at low spring up to high tide level. The coral has very substitute for stone	Bombay. 1813
that the Mr Fe that he	Cutch. 1814
that the Mr Fe that he	Madura. 1815 Tinnevely. 1816 C. 1816

CORAL.	Coral Reefs.
<p>CORAL REEFS</p>	<p>Chattiram, the thickness of the coral reef exposed above the surface of the water is at least 10 feet, and probably much more." Further on he remarks: "At the Pamban end of the raised reef it shows a slight northerly dip, and masses of dead coral, apparently <i>in situ</i>, protrude through the sand below high water mark. Reefs of living coral fringe the present coast, but these I was unable to examine, so cannot say whether the corals now growing there are specifically allied to those which</p>
<p>Trichinopoly. 1817</p>	<p>the South Indian coast — the great coral reefs of South India — y to oral s of fore me, rose the land of that remote age, worn and wasted, it may be, in the sequence of the myriad centuries that have since rolled over it, but in the present of fact — a submerged ination, I might have coral descri unalte gathered surface. a e sh and bbour- d hard espect from a modern beach. "But though, to an uncritical eye, the shells of that old sea might seem very like the volutes, olives, cowries, and ark-shells now thrown upon the Madras sands (and perhaps, indeed, they were their remote ancestors), it needed but to look on the great coiled ammonites scattered here and there in the broken ground, to know of a surety that around me lay the relics of a cretaceous sea. When these enclosed and protected living organisms, the coarse sandy deposits that underlie our English chalk, were slowly accumulating in a shallow ocean, where now spread the corn-fields and hop gardens of Surrey and Kent, and on the sinking sea bottom of South-Eastern England, Northern France, and Belgium, the</p>

Ornamental Corals.

CORAL.

thousands of feet of white calcareous mud that, long since upheaved and hardened into chalk, greets the homeward bound Indian in the Dover Cliffs, had yet to be slowly extracted through long ages from the sea water by minute organisms long since extinct."

B.—ORNAMENTAL CORALS.

ORNAMENTAL
CORALS
1818

Very little can be learned for certain of the indigenous living ornamental corals. Indeed, it seems probable that in some of the passages

already
of the
series
ing
indige

since, for ornamental purposes, it is only the sclerobasic polypes that form a calcareous substance of sufficient consistence to admit of being cut

White

1823

1824

1825

1826

1827

1828

1829

1830

1831

1832

1833

1834

1835

1836

1837

1838

1839

1840

1841

1842

1843

1844

1845

1846

1847

1848

1849

1850

1851

1852

1853

1854

1855

calls "club-shaped *Porites*" He also says — "I have noticed in the bazars, though I have never gathered it on the coast, a curious species of coral resembling the horse-tail *Isis*. It is branched like a tree with white striated stony joints and black horny smaller joints between, which render the whole flexible" It may be here remarked that

man of the sclerobasic polype has a alternat or part one of a calcareous

long moss, also occurs, and 'black coral,' of which beads are made, is brought from the Mergui Archipelago" Of Tenasserim Mason further says — "A tree coral two feet long, of a deep scarlet, is found on the coast, which the residents often call 'red coral,' but it is not the red coral of commerce, it does not grow like that, and the red colour is confined to the epidermis, the substance of the coral within being grey"

In concluding this brief review of the literature of the Indian ornamental corals, it must be admitted that we are grossly ignorant of the subject. There are no coral fisheries in India, and we do not know whether or not this is due to the absence of corals of commercial value, nor do we possess any knowledge as to the likelihood of the more

1823

CORAL.

Trade in Corals.

valuable corals succeeding, if introduced into Indian waters. No effort has as yet been made to propagate new species or improve the existing Indian corals.

TRADE.
1829

TRADE IN CORAL.

Some conception may be arrived at of the magnitude of the trade in Coral when it is recollected how many races of people in India regularly wear necklaces of coral. How far the prized ornaments may be derived from India, we are not able to say.

Prepared,
1830

Beads.
1831

Imitation,
1832

... and ... d coral
... ne years
... his may
... ent. in
... 126 and
... 18 Of
... some-
... again
... of

MEDICINE.
1833

real.
Medicine.—In addition to being used for adornment ornamental
... time and are
... ed by being
... nd corals are
... onsumption,
... ns "Ainslie
... hen calcined

CORCHORUS
acutangulus.

The Angular Fruited Corchorus.

JUTE.
1839

CORCHORUS, Linn.; Gen. Pl., I., 235.

The generic name for this group of annual plants is derived from the property of the leaves (*κορη* the pupil of the eye, and *κορηω* to purge or clear).

1840

Corchorus acutangulus, Lam.; Fl. Br. Ind., I., 398; Wight,
[TILIACEÆ.]

Syn.—C. FUSCUS, Roxb., Fl. Ind., Ed. C.B.C., 429, Ic t. 739

Vern.—Tistápat, BENG.

References.—Dals and Gibs, Bomb. Fl., 25; Kura, Contrib. Burmese Fl., 130; F. von Mueller, Sel. Extra-Trop. Pl., 88.

the nodes
g woolly
e upper
t paral-
most);
base of
groove.
1 patch

wild species in India,

the hotter parts of India and Ceylon. Roxburgh remarks that it flowers during the rainy and cold seasons, is never cultivated, and differs from *C. tridens*, L., in having only one style; and from *C. trilocularis*, L., in having only one row of seeds in each cell. Dalzell and Gibson say that in Bombay it is a common weed, and Roxburgh that it is a native of various parts of India.

all over Burma
angulus as
lar for the
nly is the
all is the
trilocularis.

C. 1840

The Round Fruited Corchorus.

CORCHORUS
capsularis.

JUTE.

the tips spreading somewhat as in *C. acutangulus*. Duthie's 7,121 has the foliage, capsules, and hairs of *C. trilobularis* with the seeds of *C. olitorius*."

Fibre.—A coarse fibre is sometimes extracted from this species and Müller alludes to this plant as an occasional source of jute.

FIBRE.
1841

1842

Corchorus Antichorus, Roemerh ; *Fl. Br. Ind.*, I., 398; *Wight, Ic.*, [t. 1073.

Syn.—CORCHORUS HUMILIS, Munro, ANTICORCHORUS DEPRESSUS, Linn.

Vern.—Baphull, HIND, Baphalli, kurand, bophalli, baháphalli, babuna, PB, Mudhíri, SIND.

References.—Dals. & Gibs., *Bomb. Fl.*, 25; Murray, *Pl. & Drugs, Sind.*, 65.

FIBRE.
1843
MEDICINE.
1844
FODDER.
1845
1846

by camels.

C. capsularis, Linn ; *Fl. Br. Ind.*, I., 397; *Wight, Ic.*, t. 311.

Vern.—*Ghr-nalid-pat* (according to Roxburgh); *Narchá* according to U. C. Dutt, BENG. The last mentioned author in the Glossary to his Mat. Med. of the Hindus gives this plant the Sanskrit name *kálasáka*.

1847

C. 1847

CORCHORUS
capsularis.

The Round Fruited Corchorus.

JUTE

References.—Roxb., *Fl. Ind.*, Ed. C.B.C., 429; *Loureiro, Fl. Coch. Ch.*, VI, 408; *Rumph.*, v. 1, 78, f. 1; *Voigt, Hort. Sub. Cal.*, 127; *Brandis*,

Botanic Diagnosis.—Alone distinguishable from *C. oltorius* by the short rounded capsule—a very unimportant character. Gamble's No. 15,912 has one capsule nearly round, while the others are distinctly those of *oltorius*, but some are 4-valved, others 5-valved. Kurz's No. 1231 of *C. acutangulus* has both 4- and 5-valved capsules, and Clarke's No. 24,899 has a 3-valved capsule. Clarke's No. 31,637 of *C. trilocularis*, has a 4-valved capsule, and Hooker and Thomson's sample of that species, from the Panjab, has a 3-valved capsule. The capsule is thus variable.

Habitat.—A common plant "throughout the hotter parts of India" This statement, originally made by Roxburgh, is current in the literature of jute. While it need not necessarily be implied that a plant is wild (e.g., indigenous) in the area where it is common, still that is the opinion popular writers have derived from the above carefully worded botanical description. The major portion of all we have learned regarding

favoured the writer with a note to the effect that he found *C. capsularis* in

condition.

and which

of Corcho-

list of the

that it is

the other

dens) has

sheet of Corchorus

tivated" has not

should have had

cept perhaps one

ever, may be an

knowledge Burmese

uently seen in de-

ected towns along the borders of forests and villages, &c." It is

h represents the

Wallich, Griffith,

ot be a specimen

a Mr. Talbot

n's specimens of

rium, have not

marked as cultivated.

nd fruited

the only

ered from

on in pro-

deceman-

duced to

of Bengal.

deceman-

non as to

1848

1849

C. 1849

The Round Fruited Corchorus.

CORCHORUS
capsularis.

its nativity. Edgeworth says of the Banda district, N.W. Provinces,

JUTE.

that *C. capsularis* does not occur in Madras. DeCandolle, after enumerating all the countries where the plant is cultivated (*vis*, the Sunda Islands, Ceylon, India, Southern China, the Philippine Islands, and Southern Asia generally) says: "I am not convinced that the species exists in a truly wild state north of Calcutta, although it may perhaps have spread from cultivation and have sown itself here and there."

The writer speaks of a greater portion of that e across either *C. capsul* wild or rather indigeno in some

parts of Western India, but grave doubts may be entertained as to either being natives of Bengal,—the province where they are now mainly cultivated, and where they exist frequently enough as weeds around the cultivated jute fields. The suggestion is offered, that, by experimental cultivation, it might be found possible to produce forms of *Corchorus* from some of the truly wild species which would closely approximate to *C. capsularis* and *C. olitorius*. With the imperfect knowledge we possess of this subject, the writer would be much more willing to admit the possibility of some such theory, to account for the cultivated jutes, rather than believe that manifest escapes from recent cultivation are the sole survivals of the wild forms of these plants. The scientific distinction based on the length of the fruit vessel (round in *C. capsularis* and elongated in *C. olitorius*) is, to say the least, scarcely worthy of as much consideration as the peculiarities recognised by the cultivators in distinguished forms that yield the distinction in the shape to give origin to certain

species of *Distichlis*, and which can be produced from the seeds of any one by careful cultivation

It is noteworthy that definite Sanskrit names should not exist for these most useful plants, while other plants of far less value have assigned to them names so precise as to distinguish their varieties, to separate their wild from their cultivated forms, and to indicate every possible structural peculiarity. There are neither Arabic nor Persian names for the

1850

1851

CORCHORUS
capsularis.

The Round Fruited Corchorus

JUTE.

urged that when Roxburgh was told that the plant grown in the Botanic Garden was jute, there were in all probability no such dealings in the fibre between Calcutta and Eastern Bengal. Besides, Mr Kerr rejects this derivation of the word, on the ground that jute is in no way a waste, rejected, by-product or remnant, as would be implied by the word *uch-chista*. At the same time Mr. Sen's idea would simply be that it was in

1852

Roxburgh were most probably, as at the present day natives of Orissa, and that, therefore, the name jute given by Roxburgh, the first European writer who used that name, was in all probability a softened form of *jhot*, a word which may be admitted to have come from the Sanskrit *jhuta*, unless we presume Mr. Sen's derivation of the word to have prevailed all over Orissa prior to Dr. Roxburgh's discovery of the plant.

1853

The Sanskrit word *Nadika* is said by Dutt to have been given to *C. olitorius* and *kalasaka* to *C. capsularis*, but while Dr. Dutt's work is devoted to the *Materna Medica* of the Hindus and is compiled from Sanskrit medical works, he only gives the above names in a Glossary at the end, and does not attribute to the plants, to which he says they refer, any properties as known to the Sanskrit writers, while the modern Hindus use the leaves of jute and the species of *Corchorus* generally, both as food and medicine. Dr. Moodeen Sheriff, a high authority on vernacular

names, does not give Sanskrit, Arabic or Persian names to the species in use in
ribited as
to jute
or other,
usage it
lly given
mention-
roduced),
irs in the
that it was then
the root from
derived, no-
e, nor indeed
operty of jute
patta may be
hata patra,ka,
would suggest

1854

a later introduction than *Crotalaria juncea* to which *patta* is compared. This idea receives further support from the fact that while *sana* occurs in the most ancient Sanskrit works, *patta* appears in the comparatively recent. In one of the references to *patta*, it is spoken of as the *chimi* (probably a misspelling for China) *pât*, a fact which would point to the cultivated jute plant having come to India from China. Mr. Hem

The Tufted Corchorus

CORCHORUS
fascicularis.

Chunder Kerr rev
to fibre or to rope
cations does there
several works *pat*
form of hemp but which by the home authors was pronounced to be more
nearly allied to flax By the beginning of the present century the word

JUTE.

the cultivation of the plant has been introduced from some other country
and most probably subsequent to the date of even the most recent Sans
krit works If a modern development we can scarcely admit that the
stock from which it was derived could have disappeared while numerous
wild plants closely allied to *Corchorus capsularis* and *C. olitorius* are

fibres only inferior

The seat of the

district through

ing lands of the

olitorius on the other hand, occurs

tern side of the Hooghly river,

and in Western and Southern

Although there are numerous references to *Patta Jutā* &c. in early
Ind an writings enough has been said to show that the greatest caution

British rule and in a fourth it is put down at 400 years ago In all
districts it is spoken of, however, as a crop regarding which some period
could be fixed, while no such language is used with regard to rice, cotton,
sunn hemp, or any other crop of an importance at all comparable with
Jute (*Conf with C. olitorius in a further page*)

Fibre — See a further page, and also Jute

Medicine — The leaves dried are used medicinally being eaten at
breakfast time with rice in cases of dysentery The cold infusion is also
administered as a tonic in dysenteric complaints, fever, and dyspepsia

Oil — 'The seed when fried over the fire yields an oil chiefly used for
lighting purposes' (*Ramshunker Sen Agr. Gaz.*, 1863)

FIBRE.

1855

MEDICINE

1856

OIL

1857

1858

Corchorus fascicularis, Lam ; *Fl Br Ind*, I 398

Vern — *Hirankhori*, *bhauphali*, *Boms*, *Tangla* or *ban ddt*, *bil*
nalita BENG

bdphull is also given to *C. Antichorus*

References — *Roxb Fl Ind*, Ed C B C, 429 *Dymock, Mat Med*
W Ind, 2nd Ed 115

Botanic Diagnosis — Capsules small ($\frac{1}{2}$ – $\frac{3}{4}$ inch) almost cylindrical, very
hard beak 3–4 splitting with the dehiscence of the capsule Seeds trian
gular or diamond shaped, more pointed at the lower end and very similar
to those of *C. olitorius* but smaller

C. 1858

CORCHORUS
olitorius.

Jew's Mallow

FIBRE.

1850
MEDICINE.
1860

Habitat—A common wild plant throughout the hotter parts of India from the Panjab to Bengal, and westward to Bombay (common, for example, at Surat). Distributed to Ceylon

Fibre—The fibre extracted from this plant is employed in Sind in the manufacture of ropes.

Medicine—Sakharam Arjun mentions the fact that the whole of this watery extract mixed
It is also given in
by the "whole plant
somewhat astringent

and is valued as a restorative." The name *hirankhorī* given to it, means deer's hoof

1861

Corchorus olitorius, Linn ; *Fl Br Ind*, 1, 397

JEW'S MALLOW

Vern—*Pat*, *koshta* (*bhunyi pat*, according to Drury, and *bhunyi*, in *ji pat*, *bhunyi* or *ban ascha*, *koshta* (according to eddy, *Tam Pannia* in N-W P (Atkinson to Dutt), *patla* (according to HUXU, and *Sing giru* (according to Ainslie), *SANS*

.. speak
so used
.. *li nalis*
.. *ists by*
.. *ins, the*
.. *ig, and*
.. *narchā*
.. *itorius*

Sir Walter Elliot alludes to this species but makes no mention of *C capsularis*, and neither assigns *Jutā* nor *Patta* to Jute

1862

Ainslie was perhaps the first European writer who assigned to this plant the Hind name *singgin-ganascha*, and while this has been reproduced by several subsequent authors the word does not appear to be in use in India at the present day, at least not in Hindustan proper. The Sanskrit names given above have already been commented on under *C. capsularis*. Mr Hem Chunder Kerr points out that the word *bhunyi* (given by various authors

.. *cent day* It is
.. *alls in a*
.. *t gunny*
.. *In ever*
.. *with that*
.. *reduced*
.. *n hemp*

References —

387 *Roxb* .. *Mat Ind*, 11,
& Gibs, *Ban* .. *ds*
333, *Athinsc* .. *d*,
Mooden Sheriff, *Supp Pharm Ind*, 114, *Murray* .. *1*,
Sind 64, *Benson Saidapet Exper Farm Man*, 63, *DeCandolle*,
Origin Cult Pl, 132 .. *es*

Botanic Diagnosis—Glabrous except the upper half of the petiole, and the primary veins on the under surface, where woolly hairs occur, nervules transverse, nearly parallel, pellucid, and anastomosing. Capsule very long and glabrous, beak straight, remains of the flower forming a thick scar. Seeds somewhat triangular, pointed at both extremities, but much more so to the hilum, surface often roughened, so as to appear as if minutely hairy

C. 1862

or Edible Corchorus

CORCHORUS
olitorius.

JUTE

Bombay, and Talbot (a botanical observer whose opinion must carry considerable weight) remarks "Abundantly wild about Yellapur" Dr Gibson has left a specimen of this species in the Calcutta Herbarium

brous the capsules are hairy along the angles and have a few of the peculiar tufted hairs of *C. trilocularis*, as well as the long narrow capsules of that species. It has also the thick and somewhat linear, coarsely serrated, leaves peculiar to that plant, but the leaves are not only hairy but have a few of the tufted glandular hairs on the under surface as well as on the fruit. Kurz gives the habitat of *C. olitorius*, as far as Burma is concerned, as "Ava, Pegu, cultivated and wild in rubbishy places and agrarian lands." Atkinson says that it is found in "Dehra Dun," but in this connection it may be added that in the Saharunpur Herbarium while there are specimens of the allied species, *C. acutangulus*, from various localities in the North Western Provinces and the Panjáb, there are none of *C. olitorius*. One specimen of *C. acutangulus* is marked as collected at Dehra Dun, and it is probable this may be the *C. olitorius* alluded to by Atkinson, Stewart, and other writers on the Flora of Northern India. In the report (to which reference has been made under *C. capsularis*) on jute cultivation in Madras, it is stated that a considerable amount of *C. olitorius* is grown in Ganjam, Godavery, Kistna, and Nellore but not for its fibre. The Collectors of Ganjam and Godavery say it is wild in their districts. The only district in the southern parts of the Madras Presidency where the plant was discovered was Salem, the Collector having found a specimen on the margin of a field, which Dr Bidie identified as *C. olitorius*. A sample of *C. trilocularis* is, however, in the Saharunpur Herbarium named *C. olitorius*, and this was apparently collected by Mr J S Gamble in the Kistna District, it bears the number 12662. The merest possibility of such a mistake existing regarding the Kistna samples reported on above may be admitted as sufficient to throw a doubt on the indigenous character of *C. olitorius* in even the northern districts of Madras. At all

plant that yielded the so-called jute of their former communication was a species of *Crotalaria* and not of *Corchorus*. Roxburgh points out in the *Flora Indica* that there is a wild form of the plant known in Bengal as *ban-pat* or wild *pat* which has reddish stems. In his *Hortus Bengalensis*, he speaks of two varieties of *C. olitorius*, a green form (the *pát*) and a reddish (the *ban pát*). This opinion is accepted by Ainslie and by

C. 1864

1863

1864

CORCHORUS
olitorius.

Jew's Mallow

JUTE.

O Shaughnessy, both of reddish *C. capsularis* the present day, applied from either of the above wild in the Panjáb, but it does not give its Panjab name, while he says it is the *bah-pat* of Bengal, a circumstance that would seem to justify the inference that Stewart's wild *C. olitorius* should be corrected into *C. fascicularis*, the more so since that species is undoubtedly wild in the Panjab, although not alluded to by Stewart. (For another error committed by Stewart see the remarks under *C. acutangulus*.) At the same time the writer, on looking over the Saharanpur Herbarium collections found one specimen, apparently correctly named *C. olitorius*, which was discovered by Dr. Aitchison (No. 476), and on which the note occurs, "occasional from Thul to Kuram." The Saharanpur Herbarium, as already remarked, does not, however, possess a sample of *Corchorus olitorius* as found in the Panjáb proper.

1865

If, after carefully considering these somewhat conflicting opinions, we still believe that *C. olitorius* is indigenous to India; if, indeed, we accept

being viewed as indigenous rests at present on doubtful evidence, but it may at least be confidently asserted that it is not wild in the districts where it is now or ever has been known to be cultivated for its fibre. Indeed there is some doubt as to whether it is really indigenous to any land.

olitorius than for capsularis. The latter would appear to have been cultivated in China before it was introduced to the people of India. It is now common in the neighborhood of Canton for

Oi moa Mr Hem

This name to the Sanskrit *au-ma* signifying 'flaxen' call *C. capsularis*, *Rami tsjima* or Chinese hemp. But in the same way *C. olitorius* has been known to the Egyptians and Syrians for a very long time, their acquaintance with it being possibly prior to the date of the evidence of a positive character, that a knowledge of the properties of the plant was possessed by the inhabitants of India. The Greek *κορχορος* was applied to a pot-herb, but in all probability the plant alluded to was not the *Corchorus* of the present day. Accepting the derivation of the Greek word as implying a drug useful in the treatment of eye diseases, it may be pointed out that no such property is claimed for the species of *Corchorus*. It is perhaps only a fanciful idea, but this property of a collyrium associated with *μαυρα* and *μολερυα*

1866

C. 1866

or Edible Corchorus

CORCHORUS
olitorius.

Mallow. It began apparently to be cultivated in Egypt about the beginning of the Christian era. It is there known by an Arabic name *melokych*, a word which seems in Crete to pass into *maulchia* (Conf. DeCandolle). It will at once be seen that these Arabic names (if indeed they be Arabic) bear no relation to the vernacular synonyms given the Hindus) to any form of the plant. The plant is known to the Muhammadans not having been introduced by their successive invasions, but as existing in the country for thousands of years from the 7th century.

JUTE.

1867

And added that it is a native of the country of the Ganges.

Proteralis is used—the name is given in order to indicate the degree of its

asserted that both forms of the jute plant are natives of Bengal, because they are plentiful weeds in cultivated situations. (Conf. with *C. capsularis*)

Fibre. Some of the most used is to

FIBRE.
1868
MEDICINE.
1869

are emollient and used in infusion as refrigerant in fevers and special diseases. The dried plant toasted and powdered is used in visceral obstructions."

Dr. K. L. Dé, O.E., says: "The dried leaves of this plant are sold in the market. A cold infusion is used as a bitter tonic, and is devoid of any stimulating property. Mr. Simon of Assam informs me that it

1870

FOOD.
1871

C. 1871

The Tuftedly hairy Corchorus.

peculiar form which may prove an undescribed species, it is known to them as a useful pot herb under the name of *bir-narcha* (Rev. A. Campbell), a name most probably derived from the Bengali *narcha* (*C. capsularis*), hence of some importance historically, since it would indicate that the knowledge of the plant was derived from the East and was anciently possessed by this primitive people. *Economic Products* gives (Part V

of baskets, &c

Corchorus tridens, Linn ; Fl Br. Ind, I, 398

Pattern - 7 sample. If 2 more used related to the next species more distinct

distributed" : "Generally

Fibre—Murray specially mentions this species as affording a cordage fibre in Sind.

C. trilocularis, Linn ; *Fl Br Ind*, I, 397.

Vern—*Kuruchuntis*, BOMB, the seeds are in the bazars sold under the name of *Rujajira*, *kaunti*, SANS, *Tandassir*, KAN (according to Lisboa), the seeds are known as *Isbund* in Sind (according to Murray)

Reference.—*Dymock, Mat Med W Ind*, 2nd Ed, 115

Botanic Diagnosis —Stems, petioles, and under-surfaces of the leaves
 fruit short
 which is often
 square on

section, obliquely and sharply truncate at both extremities, hilum large with a raphe-like cord thrown from it to the top of the seed crossing one of the angles. The writer would be disposed to unite *C. trideas* and *C. trilocularis*, and bring with these, into a section characterised by the seeds, the species *C. urticæfolius*. He can put no reliance on the presence or absence of a short style or of a spreading stigma, as he has found both these conditions on the same plant. The fruits of the species of *Corchorus* are more variable than any other part of these plants.

Habitat—The *Flora of British India* states that this species is met with in the N-W Provinces, the Panjab, Sind, and south to the Nilgiri hills. Roxburgh, however, says that it is a native of Bengal, and flowers about the end of the rains, and Lisboa that it is found in Gujarāt, Sholih

...and other

Greeks. Theophrastus says *παρορμιζόμενα, διὰ τὴν πικροτῆτα κάρ-
χυρος* (H. P., 77) Pliny (21, 32, and 25, 13) also mentions it as a poor
kind of pulse growing wild." Murray states that "the plant macerated
in water for a few hours yields a mucilage which is prescribed as a

The Commercial Fibre.

CORCHORUS.

demulcent, and the seeds as a specific in rheumatism" (*Pl and Drugs, Sind, 65*)

The *Ulfas Udwyeh*, by Noured-din Mahomed Abdulla Sherazi, uses the name of *isbund* for a species of what appears to be mustard seed.

JUTE.

JUTE.

JUTE.
1879

In connection with the reports of the Calcutta International Exhibition the writer published the greater portion of the facts which will be found in the present account of the fibre obtained from the species of *Corchorus*. In a further volume the commercial aspects of jute will be given (see **JUTE**), while in the following pages an effort is made to present a general and historic sketch of the subject together with certain facts of economic interest connected with the species of *Corchorus*. It may here be stated that the

a trade in *Malachra capitata*. The reader is, therefore, referred to the ac-

Comm. and Vern. Names.—Jute, or Jew's Mallow, **ENG**; *Jute, mauve des juifs, corde textile*, **FR**; *Jute*, **GERM**, *Pat*, **BENG**. Roxburgh says that "the Bengalis call it jute," but Royle enters into an explanation of

References.—*Hem Chunder Kerr's Report on Jute and other Fibres in*

Corchorus

HISTORY OF THE JUTE INDUSTRY.

HISTORY.
1880

The history of the modern Jute industry is exceedingly interesting and intimately associated with the British rule in India. There can be no doubt that jute was known to the people of India from compa-

CORCHORUS.

The Jute Fibre

HISTORY.

... ..
 :
 :
 f
 1

sunni, *patla*, and *blangi* were synonymous and generic terms for fibre and coarse cloth, without much regard to the plant from which the fibre was obtained. If so, about the beginning of the present century, the word *pat* became fixed and associated with the fibre of *Corchorus olitorius* and *C. capsularis*. Prior to that date the Government returns of exports from India mention *hemp fibre*; this must have been either *sunni* or *jute*, since the true hemp fibre has not been cultivated for centuries at least, and modern export...

1881

largely clad in jute cloth of home manufacture, such as, at the present day, is used by the aboriginal tribes. The increased facilities for the importa-

bags were required for this tra-
 greedily bought up. The hig
 tive to increased activity, and
 a recognised part of the Bengal peasant's work. By and by, however,
 European machinery began to compete with manual labour, and in due
 time it gained the day. Jute was exported to Europe for cordage, and
 ultimately for the manufacture of the bags required in the grain trade.
 The first commercial mention of the word "jute" is in the customs
 returns of the exports for 1828, when 364 cwt. were sent to Europe. Soon
 the agriculturist found that his time would be more profitably spent in
 preparing an extra quar-
 compete with steam and r
 speedily outstripped the

... ..
 trade took a new start in
 no effort was made to im-
 prove the British manufacture
 In that year, however, the "I
 ed at Ishera near Serampore
 coffee plantations in Ceylon,
 Council of that Island; these
 Company, Limited," and are
 Three years later...

jute factory came into existence following these...
 rapidly in every direction around Calcutta. In the Trade Returns for

C. 1881

of European Commerce

CORCHORUS

was 6,441,863 gunny bags brought into competition steadily, and in 1879-80, exported from India. The relative importance of the export trade in raw jute, as compared with the exports in manufactured jute of all kinds, may be seen by a careful

HISTORY.

This is of course a comparison between the total exports of raw jute and a portion of the Indian manufactures. In a further page the relative amount of Indian manufactured jute exported as such and the amount used up locally or devoted to the export trade in grain, will be found. But

which the jute manufactures have passed out of the hands of the Indian peasants, who alone, little more than 40 years ago, met the demand for gunny bags. This is seen very clearly when the above figures are compared with the exports of 1850-51. At that time the value of the gunnies exported was greater than that of the raw jute,—the former being £215,978, the latter, £197,071. There were no European factories in India in 1850, so that the market was supplied by the Indian peasant's hand loom. Steadily the exports increased, the demand for gunnies calling into existence the Dundee mills, and soon after the Indian factories. Nothing could demonstrate the development of the jute trade more than a careful examination of the exports of raw jute and manufactured jute from 1854 to 1887. During that period 24 factories, larger than the entire number of European factories, came into existence, and

1882

Dundee and other foreign manufactures.

CULTIVATION AND PREPARATION OF THE FIBRE.

CULTIVATION.
Area.
1883

ed
ten
Go
hae
20
wit
that more than half the annual yield of fibre is exported to foreign countries and mainly to Great Britain and the United States of America, the

CORCHORUS

The Jute Fibre

CULTIVATION

Tipperah 117,000, Furreedpore 85,000, Rajshahye 45,000, 24 Parganas 44,000, Dinagepore 40,000, Bogra 34,000, Nuddea 30,000, Jessore 30,000, Khoolna 30,000, Purneah 24,000, Hooghly 19,000, Goalpara 15,000

In other provinces, jute, though occasionally cultivated, is rarely so on

Impossible
in Madras
1884

to Government on certain samples of jute produced in "Tausa" but

Madras Manual (Vol. I, 361), it is stated that a portion of the jute used by Messrs Arbuthnot & Co. is produced locally, "but it is hoped that before long the supply will be drawn entirely from the district." Recent experiments have, however, been made in order to discover whether the true jute plant could be profitably grown in Southern India. Mr Benson (in his *Saidapet Experimental Farm Manual and Guide*, page 63) gives the result, arriving at the conclusion that, unless some parts of the Northern Division be more suitable, jute cannot be grown in Madras. So in a like manner it has been tried in Bombay and Burma, with apparently the final verdict that, in these provinces, it cannot be produced at a price to compete with Bengal. The plant can be grown most successfully in Burma, but the cost of labour has proved fatal to any idea of an extensive commercial industry. In 1872-73 Mr Hem Chunder Kerr estimated that there were one million acres under jute in Bengal and Assam distributed over 37 million acres of country, and that should the

Actual area
1885

per acre.
in 1884
maunds
assumption
8 maunds
Upon
to 1884
to 1880,
is of jute

into Calcutta were carefully recorded and the above figures may therefore be accepted as indicating the expansion of the area under jute in Bengal. As confirmatory of this general conclusion, based on the pub

*An effort has been made to correct returns in maunds into cwt. as being more likely to be understood by European readers, but where this has not been done, the result may be arrived at by the following simple rule: maunds $\times \frac{1}{4}$ = cwt

C. 1885

of European Commerce

CORCHORUS

lished figures of imports into Calcutta and Chittagong, it may be here added that Mr. Finucane (Director of Land Records and Agriculture in Bengal) in his report of 1886 to the Government furnished him by Naraingunge Hejute of 400lb each. Wilson adds the mills in Bengal,

CULTIVATION.

but the
tion of
Mr. Finucane
on the
tration

1886

afforded responsible for the italics in the above quotation. It is desirable to draw attention to the fact that the record of the jute trade preserved by merchants bears a close approximation to that tabulated by Government from the very extensive and complicated returns of road, river, and railway traffic, the concentration in the ultimate centre thus being seen to preserve a distinct relation to the far-reaching ramifications of the stream of supply. But Mr. Finucane concludes his review of Mr. Wilson's figures as follows:—"If the annual average of the eight years ending 1884-85 be taken into consideration, the difference between the two sets of figures is not considerable, the estimate worked out in this office from the data above described being only 3.97 per cent. less than that of Mr. Wilson."

Soil.—Jute seems to be capable of cultivation on almost any kind of

Soil
1887Climate
1888

down. The cultivator did not regard as worth cutting

Preparation
of Soil
1889

Preparation of Soil—It may be stated that, when the crop is to be raised on low lands, where there is danger of early flooding, ploughing commences earlier than upon the higher lands. The more clay in the soil, the more frequently it is ploughed before sowing. The preparation thus commences in November or December, or not till February or March, the soil is generally ploughed from four to six times, the clods are broken and pulverised, and at the final ploughing the weeds are collected, dried, and burned.

Seed
1890

Seed—No special attention is paid to the selection of good seeds, nor do the cultivators buy and sell their seeds. In the corner of the field a few plants are left to ripen into seed, and these are, next year, sown broadcast. The sowings, according to the position and nature of the soil, commence about the middle of March and extend to the end of June.

Harvest.
1891

Harvest—The time for reaping the crop depends entirely upon the date of sowing; the season commences, with the earliest crop, about the end of June, and extends to the beginning of October.

C. 1891

CORCHORUS

The Jute Fibre

CULTIVATION

The crop is considered to be in season whenever the flowers appear, and past season, with the fruits. The fibre from plants that have not flowered is weaker than from those in fruit, the latter is coarser and wanting in gloss, though stronger. It is late reaping that is chiefly accountable for the coarse fibre found in the market.

Crop
1892

Crop—The average crop of fibre per acre is a little over 15 maunds, but the yield varies considerably, being as high as 30 to 36 in some places, and is also very dependent on the soil, the season, the management of the farm, the ground, the weather, and the average yield.

Retting
1893

Retting—The process of retting is carried out in two or three days, to give time for the decay of the leaves, to discolour the fibre in the retting process, in others the bundles are carried off and at once thrown into the water. There is some ground for thinking that, if the drying of the leaves by stacking does not prevent the discoloration of the fibre, the fibre itself is likely to be benefited by the process, since it is found to separate more readily from the stems, and is thereby saved from the danger of rotting from over maceration. In some districts the bundles of jute stems are submerged in rivers, but the common practice seems to be in favour of tanks or road-side stagnant pools.

It varies from two to twenty days, and the bundles are turned over the surface of the water, the kind of soil, the season, the management of the farm, the ground, the weather, and the average yield. It varies from two to twenty days, and the bundles are turned over the surface of the water, the kind of soil, the season, the management of the farm, the ground, the weather, and the average yield.

It varies from two to twenty days, and the bundles are turned over the surface of the water, the kind of soil, the season, the management of the farm, the ground, the weather, and the average yield.

It varies from two to twenty days, and the bundles are turned over the surface of the water, the kind of soil, the season, the management of the farm, the ground, the weather, and the average yield.

It varies from two to twenty days, and the bundles are turned over the surface of the water, the kind of soil, the season, the management of the farm, the ground, the weather, and the average yield.

It varies from two to twenty days, and the bundles are turned over the surface of the water, the kind of soil, the season, the management of the farm, the ground, the weather, and the average yield.

It varies from two to twenty days, and the bundles are turned over the surface of the water, the kind of soil, the season, the management of the farm, the ground, the weather, and the average yield.

It varies from two to twenty days, and the bundles are turned over the surface of the water, the kind of soil, the season, the management of the farm, the ground, the weather, and the average yield.

It varies from two to twenty days, and the bundles are turned over the surface of the water, the kind of soil, the season, the management of the farm, the ground, the weather, and the average yield.

It varies from two to twenty days, and the bundles are turned over the surface of the water, the kind of soil, the season, the management of the farm, the ground, the weather, and the average yield.

Extraction by
Machinery
1894

C. 1894

Mr W Cogswell, however, who is an undoubted authority on all questions connected with jute, expressed in December 1881 his opinion that a softer fibre was obtained by the old process (*vide A H Society's Proceedings, December 1881*)

PROPERTIES OF JUTE FIBRE.

Chemical and Microscopic—"The fibre, as found in commerce, consists of the fibre bundles separated from the cortical parenchyma. The bundles contain 6 to 20 fibres. The fibres are firmly coherent in the bundle, the cohesion taking the form of fusion of contiguous walls, the line of fusion being very apparent. The ultimate fibres are of the normal fusiform type, 1.5–3 mm in length. In section they are seen to be thick walled and polygonal. *Reactions*, characteristic of the jute-allied group of fibres, are brown with iodine, deep yellow with aniline sulphate, purple with phloroglucin and hydrochloric acid, a strong affinity for the basic colouring matters. *Mercerised fibre*—*Microscopic features*. Concentrated solutions of the alkalis have a remarkable action on fibres of this group. They resolve the bundles more or less completely, and cause the fibre wall to swell so as to almost obliterate the cavity. The filaments, in addition to being made finer, are much softened in texture, and develop a wavy outline, giving the fibre very much the appearance of wool' (*Cross, Beavan, King, and Watt, Report on Indian Fibres, p 36*). The chemical analysis, as given in the report just quoted, may be here briefly reviewed. Jute, in point of percentage of cellulose (perhaps the best criterion for judging of the value of a fibre), is about equal with Urena 77.7, Calotropis 76.5, Abutilon 75.0 and Agave 75.8, and follows after Abroma 80.0, Rhea 80.3, Flax 81.9, Sida 83.1, Crotalaria 83.0, Marsdenia 88.3 and Girardinia (Nilgiri nettle) 89.6. Jute possesses 76.0 per cent, and is thus in point of cellulose about the eighth most valuable fibre in India. It is noteworthy that of the fibres enumerated—Abutilon, Urena, Abroma, Sida, and Jute are obtained from closely allied plants and yield very similar fibres. But of these jute is the next to the last in point of chemical merit, Sida being the first of the series. This is a fact of the greatest importance, when it is added that the experts who examined these fibres at the Colonial and Indian Exhibition pronounced

PROPERTIES
OF JUTE
1895

Mercerised.
1896

Cellulose.
1897

Ash,
1898

the cell cavity completely, thus causing the filaments to become much finer and softer in texture. By nitration jute gains in weight, becoming 128, being in this respect inferior to any of its allied fibres, but it is found to contain 47 per cent of carbon having the highest amount of any recorded Indian fibre, Sida, for example, possesses 45.2, flax 43.0, and Bauhinia fibre only 40.7

CORCHORUS

The Jute Fibre

PROPERTIES
OF JUTE.

The results of the chemical and microscopic investigation of jute, instituted by Messrs Cross, Beavan, and King, may be briefly stated to

Strength
1899

wool

Strength and Industrial Properties.—Royle remarks "Jute is certainly characterised by fineness, silkiness, and facility of spinning, but it is less strong than many other Indian fibres, which are possessed of similar

1900

to the cultivator's necessities and the manufacturer's wants. The question is not, therefore, one as to whether jute or *Sida* is more easily cultivated and gives the better result in point of yield of fibre, but whether the intrinsic superiority of *Sida* fibre would justify its experimental and systematic cultivation until a stock was produced that could be grown as readily and admit of as rapid decortication as is the case with jute. The plant is wild to-day, and it is unfair to compare the yield of fibre from such a plant with results obtained from jute. After careful cultivation for 10 or 20 years it would be fair to compare the ease of cultivation and yield of fibre in *Sida* with that of jute and during this experimental stage remunerative returns might easily be obtained since there can scarcely be two opinions as to the superiority of *Sida* over jute for the finer textile purposes. Roxburgh found in his comparative tests of the fibres of India that a "dry line" of *Corchorus capsularis* broke with a weight of 164lb and a "wet line" with the same weight, whereas *Corchorus olitorius* gave way with 113 and 125lb respectively, the wet line gaining 11lb in weight. This fact of the superiority of the fibre of *capsularis* over *olitorius* is well known in modern commerce. To compare with these results it may be mentioned that, under the same test, a "dry" and a "wet" line of sunn-hemp broke with 160lb and 209lb, respectively, the latter gave way for water for *Corchorus* give way observed in the tanned ropes, but the tarred seemed to preserve strength considerably, the line fresh and tarred broke with 61lb, and after maceration for 116 days bore a weight of 60lb.

1901

The defect of jute is the difficulty to spin the higher counts so being about the finest made, commercially, and when manufactured the fabric lasts well, so long as it is not submitted to a damp influence, but rots rapidly when damp and exposed to the atmosphere.

C. 1901

of European Commerce

CORCHORUS

PRICE OF CULTIVATION

No trustworthy figures are available of the prime cost to the cultivators of raising and extracting a maund of jute fibre. But the following figures which have been kindly furnished by a mercantile firm lead to the rates paid to the growers. Jute landed in Calcutta cost as follows per maund in the four years ending 1883 —

PRICE OF
CULTIVATION
1902

Qualities		1879-80	1880-81	1881-82	1882-83
		R a p	R a p	R a p	R a p
Narayanj	Fine	5 2 9	5 0 3	4 15 0	3 7 6
	Medium	4 9 6	4 6 9	4 3 4	2 15 2
	Common	4 0 9	3 13 7	3 10 4	2 7 6
Seraganj	Fine	5 4 0	5 2 0	5 1 0	3 9 0
	Medium	4 1 0	4 8 0	4 4 0	3 1 0
	Common	4 2 0	3 15 0	3 12 0	2 9 0

The average prices for the last four years were as follows —

	Bengal	Assam
	R a p	R a p
1883-84	3 12 0	4 0 0
1884-85	3 4 0	2 13 0
1885-86	3 4 0	3 1 0
1886-87	3 10 0	3 2 0

The charges per maund incurred from the time the jute is purchased from the producer to the time it is landed in Calcutta are approximately as follows —

	Narayanj	Seraganj
	R a p	R a p
Freight to Calcutta	0 8 0	0 8 0
Unloading shipping &c	0 2 0	0 2 0
Attda	0 2 0	0 2 0
Bepasport	0 5 0	0 5 0
TOTAL	1 1 0	1 1 0

Deducting the charges just shown from the cost of the jute landed in Calcutta will give the rates paid to the grower thus —

Qualities		1879-80	1880-81	1881-82	1882-83
		R a p	R a p	R a p	R a p
Narayanj	Fine	4 1 9	3 15 3	3 14 10	2 6 6
	Medium	3 8 6	3 5 9	3 2 4	1 14 2
	Common	2 15 9	2 12 9	2 9 4	1 6 6
Seraganj	Fine	4 3 0	4 1 0	4 0 0	3 0 0
	Medium	3 0 0	3 7 0	3 3 0	2 0 0
	Common	3 1 0	2 14 0	2 1 0	1 8 0

The prime cost to the cultivators must be something lower than the figures shown in this last statement, and assuming that the data fur

CORCHORUS.

The Jute Fibre

PRICE OF CULTIVATION.

nished are near the truth, if not correct, they lead to the following important inferences, *vis*, (a) that the price during the past few years, and (b) that men have not varied, those of the with the fall of prices in Calcutta. siderably; a good year induces an indiscriminate extension of the area which must of course be attended the following year by a fall in price, and from the

from all sources was practically the same as in the previous year; while the value of the exports from Chittagong was twenty-seven lakhs more

1903

May, when the young plants were seriously damaged by floods which accompanied the cyclone, especially in the districts of Rungpore, Rajshahye, Dinagepore, Bogra, Julpigoree, and parts of Hooghly. These localities, however, excepting Rungpore, are not of first-rate importance

said that
ar; and,
normal,
s will be
ted that
id on the
August."
which
rice of
to the

	Average whole-sale price in 12 selected districts in Bengal.			Average declared value as per Custom House Returns		
	R	a.	p.	R	a.	p.
1876-77	3	0	0	4	4	0
1877-78	3	0	0	4	12	0
1878-79	4	0	0	4	10	0
1879-80	4	10	6	4	13	0
1880-81	4	8	0	4	14	0
1881-82	4	8	0	4	14	0
1882-83	3	8	0	4	1	0
1883-84	3	12	0	4	12	0
1884-85	3	4	0	4	1	0

C. 1903

COMMERCIAL VARIETIES

COMMERCIAL
VARIETIES.
1904

There are several well known commercial VARIETIES of jute fibre, of

order those of importance being marked *

I. Bakrabadi—A beautiful soft fibre, one of the finest qualities from the

2 r rope
south

3. ' chiefly
e near

Faridpur where there was formerly a large mart for this variety of jute. The name is given to all the jute from Backerganj and Faridpur.

4. * Desi (in commerce *Dassees*)—This is a useful and good fibre, largely used for gunnies, it is long, soft, and fine but it has a bad colour and is pronounced 'fuzzy'. It is produced in the

5. *

ganj and is said to consist of two kinds or sub-varieties —

(a) Bilan Deswal, or fibre from the crop grown over *bhals* or marshes

(b) Charna Deswál, or fibre from the crop grown on *churs*

6 Jangiputi—A poor fibre short weak, and more suited for paper manufacture than for spinning. It comes from the Pubna district.

7 Karimganj—A fairly good fibre very long and of good colour. It comes from the Mymensingh district, taking its name from a

88

9

10 "

II "

These 11 qualities and others of minor importance, are in commerce generally grouped under four leading classes represented by the *Seraganj*, *Naranganj*, *Desi*, and *Deora*, and these, again, are classed as "Fine," "Medium" and "Common," according to the qualities of the fibres. Mr James Duffus, in a letter addressed to the writer, says of this

1905

C. 1905

CORCHORUS.

The Jute Fibre

COMMERCIAL
VARIETIES.

subject. "Every small mart in Eastern Bengal has a jute of its own, quite as worthy of mention as many of the minor forms alluded to above." This remark has an interest beyond that of commerce, for we must either

FOREIGN
TRADE
1906

FOREIGN TRADE IN JUTE AND JUTE MANUFACTURES.

F
The
dust
tion of the plant, and of the Indian manufactures.

INTERNAL
TRADE
1907

INTERNAL AND COASTING TRADE

.. .. . discussed under the
me Consumption
to indicate very
various existing
modes of conveyance. In a special Report on this subject Colonel L.
.. .. . but he refers
by him he
and Chitta-
r, the latter
the foreign
e of British

India for that year the foreign exports were put down at 8,369,686 cwt and the coasting trade at 1,267,034 cwt, making a total of jute shipments from Indian ports of 9,635,720 cwt. Colonel Conway-Gordon gives the total imports into Calcutta as 9,392,813 cwt, of which 3,579,062 cwt were conveyed by native boats, 1,969,237 cwt by steamers, 3,482,522 cwt by the Eastern Bengal Railway, 148 cwt by the South Eastern State Railway, 356,496 cwt by road, and 5,348 cwt by sea. Thus the COUNTRY BOATS head the list, carrying to the sea-board 38 1

1908

to the mills it would be seen that jute is of importance to a far greater number of persons than to the 50,000 who find daily employment in the

* For the purpose of allowing of comparison with the returns of foreign trade, Colonel Conway-Gordon's figures of maunds have been converted into cwt

C. 1908

of European Commerce

CORCHORUS.

European factories But even this estimate would leave out of all consideration the indigenous hand looms that are still able to compete with steam in the production of jute cloth, bags, and cordage

HOME
MARKET

RAW JUTE

EXPORTATION AND HOME CONSUMPTION

EXPORTS.
1909

The following abstract of the EXPORTS OF RAW JUTE FROM CALCUTTA will be found interesting, as showing the steady and constant increase and development of the jute trade The mean exportations for

Up to	Average of five years in cwt
1832-33	11 800
1837-38	67 483
1841-43	117 047
1847-48	234 055
1852-53	439 830
1857-58	710 826
1861-63	969 724
1867-68	2 628 10
1871-73	4 858 162
1877-78	5 362 267
1882-83	7 274 000

The foreign exports of raw jute were, in 1882-83 10 348 909 cwt

1910

senting an increase in value from Rs 20 to Rs 84 69 259 in the short period of 55 years (e.g., from £62 to £5 846 925 for exported raw jute alone) speaks volumes for the noble fleet of merchant vessels trading with our Indian ports Mr Hem Chunder Kerr, in his valuable *Report on*

the held of European commerce

The figures of Indian trade show that the exportation of jute steadily increased from 1 092 668 cwt in 1860-61, to 3 754 083 cwt in 1870-71, that in 1871-72 it suddenly rose to 6 133 813 cwt, and during the past 5 years has preserved an average of about 7 274 000 cwt

In 1882-83 Indian commercial men calculated that on an average Scotland consumed over 18 400 bales (73 600 cwt) a week Of these Messrs Cox Brothers take 2 200, Messrs Gilroy & Sons 750, Messrs Malcolm, Ogilvie & Co, 650, Mr John Sharp 700 In England the weekly consumption is over 1 860 bales, the largest consumers being the Barrow Company, 600 In Ireland the total weekly

1911

C. 1911

CORCHORUS.

The Jute Fibre

EXPORTS.

consumption is about 730 bales, the largest firm consuming under 300 bales a week. Thus Great Britain requires over 21,000 bales or 84,000 cwt a week, or 4,200,000 cwt a year to keep her existing jute factories

1912

consumption of 195,000 cwt. The Scotch power-loom alone consume 73,600 cwt. a week, or 3,710,000 cwt. a year. Although in some respects this estimate has been disturbed, it is relatively correct for the present year 1887-88.

France requires 4,000 bales a week, its largest consumer, Saint Freres, requiring 700 bales; Germany requires 2,170 a week, of which the Brunswick Jute Spinning Company consume 770 bales; Belgium requires 845 bales a week; Austria, 580, Spain, 250; Holland, 400; Norway, 100. Taking annual figures for the whole of Europe it is found that Great Britain and the Continent of Europe require 1,800,000 bales a year, or 6,428,580 cwt. It may be here stated that as merchants adopt the calendar year, and Government the financial, e.g., from April to March, considerable difficulty has been experienced in comparing the Government Statistical Tables of Exports with those kindly supplied by one or two well known jute firms in Calcutta.

1913

Comparing with the above figures the 22 Indian factories at work in India in 1882-83, each 700 bales per week, or 600,000 bales a year, that to keep these jute-consumers were required, as summed by America, Australia, and other foreign countries, viz., 600,000 bales, or 2,142,493 cwt, not included in the above calculation, the annual These

Annual
Capital.
1914

Looking at the exportation of raw jute, of manufactured jute, and the home (Indian) consumption known to our commercial men, the statement that the jute trade is at least represented at the present date by an annual consumption of over 15,000,000 cwt. of raw jute does not seem to be far from correct. This is roughly equivalent to an annual turn over of capital equal to about 12-14 millions of pounds sterling as compared with the exports in 1823 of £62.

MANUFACTURES.
1915

THE MANUFACTURES OF JUTE AND THEIR EXPORTATION FROM INDIA.

... factories have sprung and an- started
... to Cal- k Com-
... The
... o to 40
... 3 spin-

C. 1915

of European Commerce.

CORCHORUS.

dies, and they give employment to 29,660 men, 11,198 women, 5,113 young persons, and 3,044 children. The Madras private jute company employs about 878 persons. Thus, up to the present date, there are in all India 24 Jute factories, which give employment to 49,015 persons and use up 2,869,088 cwt of jute. They are almost exclusively employed in the gunny bag or cloth trade, three only doing a small business in cordage, floor cloth, or other manufactures.

In 1870 there were in England 14,000 spindles, in Scotland 99, in Ireland 10, and in India there

MANUFACTURES.

every individual factory. Judging from the published statistics of jute factories in Scotland during the year 1879, and comparing a fixed number of these with the Indian factories for the same year, we may, however, conclude that the Indian mill workman was inferior to the Scotch workman in the ratio of 3 to 7. That is to say, it requires 7 persons to work one loom in an Indian factory, against 3 workmen in a Scotch factory. This conclusion is arrived at by dividing the total number of persons employed in a factory by the number of looms. For all Scotch factories and the average for all Indian factories, of course this calculation is open to the objection that the Indian factories not manufacturing the same goods may be accepted as giving some sort of comparison.

1916

FOREIGN TRADE IN MANUFACTURES.

Prior to 1857 the exports of Jute manufactures from India represented hand-loom fabrics. In 1850 these were valued at £215,978, whereas the trade in raw jute was only £197,071. Fifteen years later the manufactured jute, exported to foreign countries, was valued at £18,27,983 (£182,798) and the raw jute at £75,06,690 (£750,669). In 1870-71 the exports were of manufactured jute £34,24,249 (£342,424) worth and of raw jute £2,57,75,526 (£257,755). But the revival in the exports of manufactured jute indicated by these figures, as also the partial decline of the foreign raw jute trade, was at once the death of the old hand-loom industry and the birth of the new power-loom. Ten years later (1880-81) the total exports of manufactured jute were valued at £1,13,06,716 (£1,130,671), of which the hand-loom produced £2,69,553 (£26,955), and last year they were valued at £1,15,18,577 (£1,151,857), of which the hand-loom produced £89,220 (£8,922). These figures indicate unmistakably the growth of the Indian power-loom foreign trade and the decline of the hand-loom. In a further page some idea will be given of the extent of the home market for jute goods.

Foreign Trade in Manufactures.

1917

LOCAL OR HOME CONSUMPTION

Local Consumption.

1918

CORCHORUS

The Jute Fibre

MANUFACTURES
Home Consumption

third of the number actually manufactured. The following table will show the relations of the home consumption to the exports more clearly —

Statement of Home Consumption and Exports of GUNNIES from 1st January to 31st December 1882

1919

Burma	13 312 306	
Straits	9,153,233	
Bombay and Persian Gulf	20 001 303	
Madras and Malabar	1,064 843	
Coromandel Coast	3 609 950	
Ceylon	177,777	
Up-country by rail	11,351,000	
Used in the export trade of Calcutta	11,848,742	
Total of Home Consumption		77,519,164
Australia	11,372,387	
New Zealand	5 060 60	
Cape of Good Hope	706 308	
Mauritius	119 078	
Egypt	691 078	
America	20,554 51	
Hongkong (not Hessans)	413 700	
Great Britain	516 417	
Europe	90, 31	
Total of Foreign Exports		41,523 607
Grand Total of Home Consumption and Foreign Exports		119 047 771

The total number of gunny bags brought to and carried from Calcutta during the past three years may be here given and alongside of these the foreign exports —

	1884-85	1885-86	1886-87
Imports	18 196 002	20 6 6 541	23 596 472
Total Exports (to other provinces of India and to foreign countries)	137,870,318	127 084 964	124 957,225
Foreign exports only	82,779 207	63 760 546	64 572 157

1920

total production of gunny bags in Bengal was perhaps little more than 150 millions, of which 64½ millions were sent to foreign countries and 85½ millions used up in India. This may be accepted as representing the bags employed in the home, cotton, oil seed, rice, and wheat trade, and

225,374
interprovincial
quantity
5,267,418
sent to these
per borne

C. 1920

Of European Commerce

CORCHORUS

Traffic of Bengal for 1887 states that 605,846 pieces were sent up-country by river "direct from the jute mills without passing the Port Commissioner's wharves" A piece of power-loom gunny is equal to 80 yards, of hand-loom, to 22 yards, so that this power-loom trade alone re-

MANUFACTURES.
Home Consumption

industry is conducted in Dinagepore, Purneah, Rungpore, Julpaiguri, and Tipperah, Julpaiguri turned out last year 2,336,660 and Rungpore 1,222,410 hand-loom made bags

CLASSIFICATION OF THE JUTE MANUFACTURES

CLASSIFICATION OF MANUFACTURE.
1921

The manufactures from *jute* or *pat* may be referred to three primary sections —

These three sections may each be referred to a number of sub-divisions, which for convenience may be arranged in two leading groups, *vis*, native and indigenous manufactures, "hand loom," and European or "power loom" manufactures, whether made in Europe or in India. We shall first enumerate the indigenous manufactures, since these bear on the history of the industry.

INDIGENOUS MANUFACTURES

Indigenous

Ind
beam
talking
of gun
poses
or gun
Hortic
on

ra, the
is said
tended

1st, *Thick cloth* used for making gunny bags. Of this there are three qualities, the best being known as *amrabati*. These correspond to the three qualities of hand-loom gunnies in commerce

CORDIA
fragrantissima

The Jute Fibre.

CLASSIFICA-
TION OF MA-
NUFACTURES

2nd, *Fine cloth*—This is generally known by the name of *mekls dhokra*, and is chiefly used as a cloth to sleep on, it is often beautifully striped blue or red

3rd *Coarse cloth*—This is largely used for making the sails of country boats (*gun*), and also for bags to hold large seeds or fruits

The following are the principal districts in Bengal where indigenous jute manufactures (hand-loom) may be said to exist to any considerable extent—Hugh, consuming about 1,20,000 maunds of jute a year, Dacca, 90,000, Rungpore, 50,000, Moorshedabad, 38,000; Malda, 25,000; Julpaiguri, Pubna, &c, smaller quantities

European
Manufac-
tures,
1923

EUROPEAN MANUFACTURES

Cloth made in Factories—Jute is now largely used in the manufacture of carpets, curtains, shirtings, and is also mixed with silk or used for imitating silk fabrics. It has been applied extensively as a substitute for hemp for this purpose the fibres are rendered soft and flexible by being sprinkled with water and oil, in the proportion of 20 tons of water and 2½ tons of train oil to 100 tons of jute. Sprinkled with this the jute is left for from 24 to 48 hours, when after being squeezed by rollers and heckled, the fibres become beautifully soft and minutely isolated, and thereby suited for a number of uses.

the
and

and other fibres were not adulterated with jute. In 1832 an enterprising Dundee manufacturer experimented once more on the fibre, and the result was that he was able to show that it might be used as a substitute for hemp. From that date jute gained rapidly in public favour. It is

and is now used for a number of purposes, and is one of the most valuable of our sub-

1. — 2. 106

op-
587.
ues

are almost exclusively the various forms of gunnies

JUTE
WHISKEY.
1924

JUTE WHISKEY.

In concluding this account of jute it may be mentioned as a curiosity that it has been proposed to utilize the jute ends in the preparation of a spirit which somewhat resembles the whiskey made from grain. The waste fibre is by means of sulphuric acid converted into sugar and the resulting product thereafter fermented and distilled

CORDIA, Linn, Gen Pl, II, 838

1925

Cordia fragrantissima, Kurz, Fl Br Ind, IV, 139, BORAGINACE

Vern—*Kalamet toungkalamet* BURM

References—Kurz, For Fl Burm, 27; Gamble, Man Timb, 71

C. 1925

The Sebesten Fruit.	CORDIA Rothii.
cultural implements, in Bengal for canoes. It might be tried for tea-boxes. It makes an excellent fuel. In a report of Chanduka in Sind (1847), it is stated that "the wood is used for sword sheaths." The Santals regard the wood as specially useful for yokes, as it does not rot into the wood of the handle.	DOMESTIC. 1939
North-Western Provinces that the leaves are used as plates, and that the viscid pulp of the fruit is used as bird-lime.	1940
Cordia oblqua, Willd.	
This is the larger SEBESTEN according to Stocks, Dymock, Birdwood, &c., <i>C. Myxa</i> being the lesser, but the vernacular names given would imply the reverse to be the case.	
The text does not give this plant the elegant name of <i>Alma viri chetiv</i> , and remarks that its synonym <i>Sleshmataka</i> is correctly translated "phlegm-speller."	
References—Roxb, <i>Fl Ind</i> , Ed C B C, 198, Brandis, <i>For Fl</i> , 336, (in part), <i>Thwaites, En Ceylon Pl</i> , 213; <i>Dalé & Gids Bomb Fl</i> , 173, <i>Sind Gas</i> , 603, <i>Bomb Gas</i> , V, 27, Dymock, <i>Mat Med W Ind</i> , 2nd Ed, p 579; Atkinson, <i>Him Dist</i> , 733, Birdwood, <i>Bomb Pr</i> , 58, 169, Smith, <i>Dic</i> , 374.	
Habitat—Found in Western India (especially Guzerat), from the	
"regarded as a demulcent."	MEDICINE. 1941
Special Opinion—"The fruit in its raw state contains a gum used hencefully as a demulcent."	
demand	FOOD 1942
C. Rothii, Rom & Schult; Fl Br Ind, IV, 138	TIMBER. 1943 1944

CORDIA
vestita.

Cordage and Ropes

ones of North-West, Central, and
Stocks says that it is sometimes

GUM
1945

fields a gum which is reported to be
prepared at Coimbatore In the Bombay Gazetteer of Baroda District,
it is stated "fruit eaten by the poor and pickled, as is the gum which
exudes from it"

FIBRE.
1946

MEDICINE
1917
FOOD.
1948
TIMBER.
1949

Medicine—The decoction of the bark possesses astringent properties,
and is also pickled

Used for fuel,
elements Baden
Powell remarks that the wood is tough and is employed for making car-
riage poles Stocks says the wood of the *lyar* is much used in Sind.

1950

Cordia vestita, Hook f & Th, Fl Br Ind, IV, 139

Syn—*GYNALION VESTITUM*, DC

Vern—Kumbli, kari k PB, Kum paiman, pin, indak, chinta, ajaria
bairula, lerula, HIND

References—Brandis, For Fl 338; Gamble, Man Timb, 271,
Atkinson, Econ Prod, N W P, V, 81, Baden Powell, Fb Pr, 575

Habitat.—A small deciduous tree of the sub Himalayan tract, from
the Jhelum to the Sarda River and Oudh

MEDICINE.
1951

Medicine—Fruit used similarly to the other species, and when ripe
is an article of food, it is considered better than that of *C Myxa* Mr
Atkinson states the flowers appear in spring and the fruit ripens in
the rains He remarks that the fruit is full of a gelatinous pulp which is
commonly eaten and considered refreshing

TIMBER.
1952

Structure of the Wood—The wood is very similar in appearance to
that of *C Macleodii*, except that the concentric lines are occasionally
interrupted, it is strong and is used for wheel and well-work.

1953

CORDAGE AND ROPES

Many fibres are used for this purpose, in fact, the natives of India are
never at a loss when in the forests to find a plant the bark of which will
serve the purpose of a string or rope The majority of such plants are
more or less used locally in the preparation of ropes or cords, a con-
siderable number are of commercial importance. Against the names in
the following list have been placed one or in some cases two * to indicate
the fibre-yielding plants frequently used for cordage, or the fibres which
hold a position of commercial importance (* * indicating greater
importance than *) —

* *Abroma augusta*
Abutilon asiaticum.
A *Avicennae*
** *Agave americana*
Alnus nitida (bridge ropes)
Artocarpus Lakoocha.
Arundo K

Bauhinia angulata.
B racemosa
* *B Vahl*
Bixa Orellana
Boehmeria macrophylla (fishing
nets).
** *B river*
Bombax malabaricum.

C. 1953

Coriander

Borassus flabelliformis.
 Broussonetia papyracea.
 Butea frondosa.
 Calamus Rotang
 * Calotropis gigantea (milkweed)
 ** Cannabis sativa.
 Careya arborea.
 Caryota urens
 Chamocrops Ritchiana
 ** Cocos nucifera (coconut)
 * Corchorus sp (jute)
 Cordia Myxa
 C Rothu
 Crotalaria Burhia.
 ** C juncea (Sunn-hemp)
 Daphne papyracea.
 Debregeasia bicolor (fire tree)
 D leucophylla
 D longifolia
 * Desmodium tiliaefolium.
 Dombeya umbellata
 Edgeworthia Gardneri.
 Eriolæna spectabilis
 Ficus bengalensis
 * Gerardia heterophylla.
 Gnetum scandens (fish tail tree)
 ** Gossypium sp (cotton)
 Grewia asatica
 G oppositifolia
 * Hardwickia binata
 Helicteres Isora
 ** Hibiscus cannabinus
 H esculentus
 H tiliaecus
 Holostemma Rheedei
 * Ischaemum angustifolium (grass)
 Laportea crenulata

INE.
56

CORIANDRUM

The name of this genus is peculiar
 peculiar smell of the plant
 plant to be viewed as a weed
 (popularly called seeds) are used
 vegetable accord only used as a spice
 races as a drug from almost
 in Britain prior to the Norman

OOD
1957

Coriandrum sativum, L.

CORIANDER

Vern — *Dhanya* or *dhan* (seed)
Kothamra (seed)
Mar Dhano (seed)
Dutt *dhanyaka*
Ko amall *TAM*
Na nau *BUM*

ays it
MIR
TARA
TARA

1958

Coriander.

CORIARIA
nepalensis.

been indicated by the formula $C_{10}H_{18}O$, and is therefore isomeric with *borneol*. By abstraction of the elements of water (by means of phosphoric anhydride) this is converted into an oil having an offensive odour

being submitted to distillation" (*Professor Warden, Calcutta*).

Medicine.—The medicinal properties attributed to this plant are many,—namely, carminative, refrigerant, diuretic, tonic, and aphrodisiac. The dried fruit and the volatile oil are used as an aromatic stimulant in

MEDICINE.
1956

Greeks."

with good results (*Bhagwan Dass (2nd), Assistant Surgeon, General Hospital, Rawal Pindi, Panjab*) "The roasted fruit is generally used" (*Dr. Bensley, Civil Surgeon, Rajshahye*). "A strong decoction of the seeds with milk an

(*D R Thomson, M*
aromatic, stimulant
Moorshedabad) "

Assistant Surgeon,
useful in colics of children, powder of fried seeds" (*Shib Chunder Bhat-
tacharya, Assistant Surgeon, In Civil Medical Charge Chanda, Central
Provinces*).

Food.—Eaten by the natives as a vegetable. The seeds are univer-
sally used as a condiment, and form one of the ingredients in curry.

They are also employed in confectionery, and for flavouring spirits.

FOOD
1957

CORIARIA, Linn ; Gen. Pl, I., 429

Coriaria nepalensis, Wall. ; Fl. Br. Ind., II., 44 ; CORIARIA.

1958

CORIARIA
nepalensis.

Coriaria.

Sind Pl., 36; O'Shaughnessy, Beng. Dispens., 270; Flück. & Handb. Pharmacog., 221; U. S. Dispens., 15th Ed., 1622; Baden Powell, Fl. Pr., 336, 575; Atkinson, Him. Dist., 749; Balfour, Cyclop., 813; Treasury of Bot., 331.

Habitat.—A deciduous shrub or small tree of the outer Himalaya

West and

Yunan

rich, but in

have been

tern Prov-

near this station, Amnold, the capital of Kumaon, being in a like manner the vernacular name for *Rumex acetosa*.

Tan.—All parts of the plant are rich in astringent acids which might be used for tanning or for dyeing.

Food and Fodder.—“The branches are browsed by sheep. The fruit is first

is a powerful poison when given in large doses. The seeds are stated to sometimes produce symptoms like tetanus.

species in French gardens, and its leaves are often employed as a black dye, and were at one time extensively used as an adulterant in Senna. Much has been written of the poisonous properties of the New Zealand species, the Toot-poison—*Coriaria ruscifolia*. Mr. Lander Lindsay gives an elaborate account of the properties of that plant in the British and Foreign Medico-Chirurgical Review (1865, p. 153, and 1868 p. 463). M. Riban attributes the poison of the fruit to an active principle, which he has called coriamyrtin, the composition of which is represented by the formula $C_{30}H_{36}O_{10}$ a substance ranked with the glucosides.

The inhabitants of New Zealand extract an intoxicating beverage from the pulp of the fruit.

— brief note
resinous
cat, after
wever, the
by cattle

artwood
t be used
ood, but

* References to the Mediterranean or New Zealand species

TAN.
1959
FOOD and
FODDER
1960
MEDICINE.
1961

TIMBER.
1962

The Cornus

CORNUS
macrophylla.

orn—a term often specifically applied to *Avena sativa*, but generically given to all cultivated grasses which yield farinaceous grains, such as Wheat, Maize, Barley, Oats, &c. When ground, Corn is designated flour or meal. See *Avena* Vol. I, 1631.

orn-flag, see *Ins*

orn-Indian, see *Zea Mays*.

orn-silk—the silky stigmata of *Zea Mays*, from which a medicinal preparation is made. See *Zea*

CORNUS, Linn.; *Gen Pl.*, I, 950

[t 122; CORNACEÆ

Cornus capitata, Wall; *Fl Br Ind*, Vol. II, 745, Wight, Ill,

Syn.—*BENTHAMIA FRAGIFERA*, Lindl

Vern — *Thammial*, *tharbal*, *tharwar*, *thesi*, *bamaur*, *bamora*, HIND.,
Tumbuk, LEPCHA, *Tharwar*, *thesi*, PB, *Bamaura*, KUMAON

References.—*Brandis*, *For Fl*, 253, *Gamble*, *Man Timb*, 212, *Stewart*,
Pb Pl, 111; *Ainslie*, *Mat Ind*, II, 454, * *O Shaughnessy*, *Beng*
Dispens, 375 *O Shaughnessy*, *Beng Pharm*, 40, *Atkinson*, *Econ*
Prod, V, 75, *Treasury of Bot*, 333

Habitat.—A small deciduous tree of the Himalaya, from the Beas to Bhutan, between 3,500 and 8,000 feet met with also in Khasia hills, where it is glabrous or nearly so

The Himalaya, in April and May, often becomes almost yellow from the conspicuous cream coloured bracts which surround the flower-heads of this plant. In the North-West Himalaya, it is particularly abundant in the lower hot valleys growing along with the berberry

Food—Dr Stewart says that the ripe fruit is sweetish, and is apparently made into a preserve and eaten by the natives. It resembles a strawberry somewhat in external appearance, and ripens in October.

Structure of the Wood—Whitish, with reddish-brown heartwood, warps in seasoning, very hard, close-grained, used only for firewood.

C. macrophylla, Wall, *Fl Br Ind*, Vol II, 744

Econ Prod, V, 75

Habitat—A tree, 40 to 50 feet high, frequent in the Himalaya, from the Indus to Bhutan, between 3,000 and 8,000 feet, found by the writer in Manipur. It flowers in May and June

Oil—A species closely allied to the *C sanguinea*, and may, like that species, be found to afford an oil from its fruits

Food and Fodder—Goats feed on its leaves, and the natives eat the fruit

Structure of the Wood—Pinkish-white, hard, close-grained, warps badly, and has an unpleasant scent, yields good gunpowder charcoal

1963

1964

1965

1966

FOOD.
1967WOOD.
1968

1969

OIL.
1970
FODDER
1971
WOOD
1972

* *Cornus florida*, alluded to as having a medicinal bark, very similar in its properties to the bark of *Melia Azadirachta*

C. 1972

CORUNDUM.

Emery Stone

1973 *Cornus oblonga*, Wall; *Fl. Br. Ind.*, II, 744

Vern.—Kagsht, SUTLEJ; Dab, KUNAWAR, Kasimol, batar, ban-jakir, hald, HIND.

References—Brandis, *For Fl.*, 253; Kurz, *For Fl.*, I, 545; Gamble, *Man Timb.*, 212; Stewart, *Pb Pl.*, 111; O'Shaughnessy, *Beng Dispens.*, 375; O'Shaughnessy, *Beng. Pharm.*, 39; Baden Powell, *Pb. Fr.*, 576.

Habitat—A small tree of the outer Himalaya, from the Indus to Bhutan, between 3,000 and 6,000 feet, met with also in the Martaban Hills, Burma, between 4,000 and 7,000 feet (Kurz).

Structure of the Wood.—Pinkish-white, hard, even-grained; warps and has an unpleasant scent.

WOOD.

1974

1975 *C. sanguinea*, Linn; *Fl. Br. Ind.*, II, 744.

THE DOGWOOD, DOGBERRY, or HOUNDS' TREE, a name given in consequence of a decoction of the bark having been formerly used for washing mangy dogs; sometimes also called the CORNEL TREE

References—Brandis, *For Fl.*, 253; Gamble, *Man Timb.*, 212; O'Shaughnessy, *Beng Dispens.*, 375; O'Shaughnessy, *Beng Pharm.*, 39; Cooke, *Oils and Oilseeds*, 38; Smith, *Dic.*, 156.

Habitat—A shrub or small tree found in Europe, Siberia, and in Kashmir, in the last-mentioned country at 7,000 feet in altitude. The writer found the plant also growing near a village in Chumba State, but it may there have been only cultivated. The young shoots are red in spring, and the leaves turn of that colour in autumn, hence the specific name.

OIL.

1976

black
amps
rmas
seful
ould

WOOD.

1977

the
&c

Coromandel or Calamander-Wood, see *Diospyros guzista* and *D. hirsuta*

Coroxylon Griffithii, a misprint which appears in Balfour's *Cyclopaedia* and in the writings of other authors. See Caroxylon and also Haloxylon.

Corrosive sublimate, see Mercury.

1978

Corundum.

EMERY STONE, Eng; L'EMERI, Fr.; SCHMERGEL, Germ.; SMERIGLIO, Ital

Vern.—Aurund, HIND; Samada, GUJ

... alumina, with
... very freely
... but the
... a few and

C. 1978

Corundum or Emery Stone.

CORYDALIS
Govaniana.

far between The finest quality of Corundum is perhaps that obtained

Punyghee in the Bellary district, North Arcot district, Kistna and Godavari, and Hyderabad territory, and on into the Central Provinces

1979

Coimbatore, p 23) Emery is said to be largely exported to Bombay (Madras Manual of Administration, II, 38, Settlement Report of Upper Godavery Dist., 42, Balfour, Cyclopædia of India, 816)

CORYDALIS, Linn, Gen Pl, I, 55

[Ill, t 16, f 2, FUMARIACEÆ

Corydalis Govaniana, Wall, Fl Br Ind, Vol I, 124; Royle,

1980

Vern.—*Bhutkis, bhutken*, HIND & BENG; *Bhutakesi*, SANS (Dutt, Mat Med Hind)

Some doubt seems to prevail as to the source of the *budhes* of the drug shops Stewart says that in the Ravi basin that name is given to the root of a *Ptychotis*

References—Stewart, Pl Pl, 10, 109 Pharm Ind, 23, O'Shaughnessy, Beng Dispens, 185, U C Dutt, Mat Med Hind, 294

Habitat.—A small herbaceous plant, found in the North-West Himá-

MEDICINE.

Root.

1981

Corydalis.

1982

in solution to dogs without inconvenience"

"The *Corydalis tuberosa* and *fabacea* in Europe have a bitter acid root, usually sold as *ARISTOLOCHIA* root, and used chiefly as an external

C. 1982

CORYLUS
Avellana.

The European Hazel.

1983

application to indolent tumors. The small quantity in our possession alone prevented the *Corydalis* and its salts from being extensively tried in the treatment of ague. The chemical properties of the salts are closely analogous to those of morphia and anarcotine, an interesting fact, as it strengthens the resemblance already detected by botanists between the PAPAVERACEÆ and FUMARÆ." It might be added also that the relation of these orders to the RANUNCULACEÆ, through *Coptis* and to BERBERIDEÆ through the berberry or *rassout* extract, is similarly borne out by their chemical and medicinal properties. (See the next species and compare with the remarks under *Coptis Teeta*, C. No 1789, and *Berberis Lycium*, B. No. 460; also *Picrorhiza Kurooa*).

The Turkey-corn or Turkey-pea (*Corydalis formosa*) contains in its roots, according to Mr. W. T. Werzell, the alkaloid *corydaline*, formic acid, bitter extractive, an acrid resin with volatile oil, a tasteless resin, and

MEDICINE.
1984

the alkaloid (*Corydaline*) found in the European species—*Corydalis tuberosus*.

The roots of all these plants are supposed to be tonic, diuretic, and alterative, and are prescribed in syphilitic, scrofulous, and cutaneous affections, in the dose of from 10 to 30 grains. The drug is also often used in the form of a decoction or tincture.

Corydalis ramosa, Wall, Fl Br Ind, I, 125.

Dr Aitchison, in his *Flora of the Kuram Valley* (Linnæan Soc Jour, XIV, page 145), says that in Kuram this common Himalayan scrambling annual is employed medicinally by the natives in the treatment of eye diseases, like all other plants with yellow sap. It is there called *mimran*. It would be interesting to know if this plant is used medicinally in other parts of the Himalaya, but these properties are not attributed to it in Kulu, where the plant is abundant. (See remarks under the preceding species and compare with the account of *Coptis Teeta* C. No 1789).

CORYLUS, Tourn, Gen Pl, III., 406.

1985

Corylus Avellana, Linn, CUPULIFERÆ.

THE EUROPEAN HAZEL.

VERN.—*Findak*, Hindak, Hind, PERS, *Chal'gost*, PERS

References.—*Brandis For Fl*, 494, *Gamble, Man Timb* 320; *O'Shaughnessy, Beng Dispens.*, 609, *U. S Dispens*, 15th Ed., 571; *Esler Powell, Pb Pr*, 2d, 385

MEDICINE
Nuts
1986
FOOD
Nuts.
1987

C. 1987

... in the Caucasus
on the
volcanic
or semi-

s, &c. It is found
and sold in the
Upper and Central

The Fan Palm of South India.

CORYPHA
umbraculifera.

Corylus Colurna, Linn

1988

Syn — C LACERA, Wall

Habitat — A moderate sized tree of the North-West Himālaya, be-
appear in March and April, and
bear every third year, and yield
(Atkinson).

OIL.
1989

mention is however, made
although the plant is suffic-
much so as to bestrew the ground for miles with the nuts

Medicine.—The nuts are not uncommon in drug-sellers' shops, being
considered tonic

MEDICINE.
Nuts
1990
FOOD
Nuts
1991

Food.—The nuts are smaller than the European variety, but are
the various
anistan and
cognised by
It is thus
rom a culti-
vated superior stock of C Colurna. As seen in the forests in the Simla
district, the actual nuts are small and rarely mature their kernels, but
they are encased in a large coarse outer coat and form large succulent
heads

Structure of the Wood — Pinkish white, moderately hard. It is only
used locally, but it is well grained and does not warp, and deserves to be
better known, especially as many specimens shew a fine shining grain
resembling Bird's eye Maple

WOOD
1992

C. ferox, Wall; Gamble, Man Timb, 390

Vern — Curri, NEPAL, Langura, BHUTIA

Habitat — A small tree of Nepal and Sikkim, 8,000 to 10,000 feet

Food — The fruit is covered with a prickly cup, the kernel is edible

Structure of the Wood — Pinkish white, moderately hard, even-
grained

FOOD
Nuts.
1993
WOOD
1994

CORYPHA, Linn, Gen. Pl., III, 922

Corypha umbraculifera, Linn; PALME

THE TALIPOT PALM OF CEYLON AND THE FAN PALM OF SOUTH
INDIA

Vern — T
panai,
Bajar

1995

29 Voigt, Hort Sub
Fl Burm., II, 524,
Bomb Fl., Supp. 63,
74, t 8, Sw Walter

C. 1995

CORYPHA
umbraculifera.

The Fan Palm of South India.

Elliot, Flora Andhrica, 169, Madras, Man Admin, 37, Moodeen Sheriff, Supp Pharm Ind, 116, Drury, U Pl, 159, Royle, Fib Pl, 98; Kew Offl Guide to the Mus of Ec Bot, 71; Kew Offl Guide to Bot Gardens and Arboretum, 33

Habitat.—A large tree of Ceylon and the Malabar Coast, cultivated in Bengal and Burma. But Roxburgh says it is "a native of Bengal,

misleading.

Fibre—The leaves are made into fans, mats, and umbrellas, and are

FIBRE
Leaves.
1996

Fibre-bundle.
1997

Paper (olas).
1998

employed, the leaves are taken whilst tender, and after separating the central ribs, they are cut into strips and boiled in spring-water. They are dried first in the shade and afterwards in the sun, then made into rolls and kept in store, or sent to the market for sale. Before they are fit for writing on they are subjected to a second process. A smooth plank of areca palm is tied horizontally between two trees each *ola* is then damped, drawn backwards and becomes perfectly moisture dries up, it is necessary to renew it till the sheet is dry etc. The writer

Braids.
1999
Hats.
2000
FOOD
Sago.
2001

construction of straw or Leghorn hats.

Food—A kind of sago is yielded by the pith. Little information of a definite kind can be discovered as to the extent in which this starch is used in India as an article of food, nor as to the methods adopted in its

C. 2001

Sago Palm, the *Coscini***COSCINIUM
fenestratum**

preparation Knox says of Ceylon that the people "beat it in mortars to flour, and bake cakes of it, which taste much like white bread, it serves them instead of corn before the harvest is ripe"

Structure of the Wood—Soft with a hard rind composed of black vascular bundles. The vascular bundles in the centre of the stem are soft. Roxburgh remarks "I do not find that the wood is put to any useful purpose"

The tree often grows to a great size before flowering, one whose measurements were given in the *Indian Agriculturist* for November 1873 as flowering at Peradeniya, Ceylon, measured height of stem 84 feet, of flower panicle 21 feet, total 105 feet, girth at 3 feet from the ground round the persistent bases of the leaves 13 feet 9 inches, at 21 feet from the ground 8 feet 3 inches. age about 40 years. The leaves are very large, often 10 to 16 feet in diameter.

Domestic and Economic Uses—In addition to what has been said of

WOOD
2002DOMESTIC
Beads
2003

Europe they are now largely employed in the manufacture of buttons. The trade in these nuts is chiefly carried on by Arabs.

Corypha Tallera, Roxb, Cor Pl, t 255

A closely allied species to the preceding, which bears most of the vernacular names given above and is put to the same industrial purposes, is a native of the north eastern coast of Madras especially in Coromandel. A third species may here be mentioned by name *C. elata, Roxb, Fl Ind 298*, a stately palm and native of Bengal, where it is known as *bajur*, but Roxburgh views *C. umbraculifera* as the intermediate form between *Tallera* and *elata*, so that even if future botanists continue to view all three as distinct species, for industrial purposes, they may be regarded as but forms of one plant. It would, indeed be impossible to separate under these plants the various properties assigned to them.

Ornaments
2004
Buttons
2005
2006**COSCINIUM, Colebr, Gen Pl I 35****Coscini** *fenestratum, Colebrooke, Fl Br Ind, Vol I, 99,*

[MENISPERMACEÆ

2007

VERN—*Wateh* (Malay)

..

Habitat—An extensive climber, met with in the forests of the Western Peninsula, and distributed to Ceylon and the Straits.

The Costus	COSTUS speciosus
<p><i>Silem</i>) "This has been in use for some years in the hospital and found to be a fairly useful medicine in certain cases of dyspepsia. I think it a fairly good substitute for calumba. It has been used in the form of powder and infusion. Preparations, &c.—The same as calumba" (<i>Apothecary J. G. Ashworth, In Medical charge, Kumbakonam</i>)</p> <p>Trade—The root is sold in Madras at $\text{Rs } 1\frac{1}{2}$ per maund, and retailed at 2 annas a pound. There are no foreign exports of the root from India but it may be had in every large bazar throughout the country, so that there must be a considerable local demand.</p> <p>Cosmetic Bark, see <i>Mutraya exotica</i>, <i>Linn.</i></p> <p>COSTUS, <i>Linn.</i>; <i>Gen Pl</i>, III, 646</p> <p>Costus arabicus, see <i>Saussurea Lappa</i> and <i>hypoleuca</i>, <i>COMPOSITÆ</i></p> <p>C. speciosus, <i>Sm</i>, <i>Wight, Ic</i>, 2014, <i>SCITAMINÆ</i></p>	<p>TRADE 2012</p>
<p>" " " " " " " " " " " "</p>	<p>2013</p>

Tsana speciosa, *Gmelin*, IV, and the *Herba spiralis hirsuta* of

Hort Sub Cal,

a
T
h

easily enough be exported from Bengal were some effort made to bring this root before the perfumers of Europe. There is a strong probability,

doubt however, that the latter and not the former is the drug sold in Indian bazars, but it is curious how the mistake of confusing two so widely distinct plants could ever have occurred. It has been deemed

COSTUS
speciosus

The Costus

2015

MEDICINE
Tubers.
2016

desirable to leave the available information in its present form, since it is by no means established that *Costus speciosus* is not used as a substitute for *Saussurea*

§ "Piesse's remarks must apply to *Aplotaxis* (= *Saussurea*), not to roots are quite insi-

as *Saussurea* there seems no doubt but that a certain amount of the tubers of *Costus speciosus* are regularly used by the natives of India both as food and medicine. The late Dr U. C. Dutt wrote on the margin of a copy of the of India, showing *Costus speciosus* at the Calcut (where a brief ing *Costus* and *Saussurea* is given), — this root is said to be bitter, astringent, and

pain in the marrow the p writer and was *Costus*, not *Saussurea* alluded to, Dr Dymock says — root is described as depurative

The *kust* *saurea*, but e been confused (in the literature of the subject, although they bear no resemblance to each other) perhaps for the past 200 years, but at the same time there is a certain amount of *Costus speciosus* root deliberately used, and not from any idea of adulteration with the supposed *Costus* of the ancients. Sir Walter Elliot gives several Sanskrit synonyms for *Costus speciosus*. He may have been mistaken as to these synonyms but he clearly recognised what the *Costus speciosus* of botanists meant, as he describes the plant. He refers to *Roxburgh's Flora Indica* Vol. I, p. 50 and to the Coromandel plants, page 126 and states that while Roxburgh in these works gives *Bomma kachchika* as the Telugu for *Zingiber roseum*, "in Vizagapatam, it (that name) is invariably given to *Costus speciosus*, which abounds in the forests of that province. The Sanskrit synonyms *Pushkara malaka* in Wilson's Sanskrit Dictionary, p. 545, and *Kasmira* (Wilson p. 219) and Brown's Telugu Dictionary, p. 224 are both applied to *Costus*." He further gives *Kasmiramu* as another Sanskrit synonym.

the root of *Costus speciosus* is the root of a plant near water and is (sic) used in massalas inodorous, and tasteless. Here there seems no reason to doubt we have an illusion to *Costus* and not to *Saussurea*.

Food.—The tuber is cooked in syrup and made into preserve in some parts of India, the natives consider it whole me. This information regarding India was first published by Roxburgh but Alsted drew attention to the fact that in *Brown's Hortus Jamaica*, Vol. II, p. 231,

FOOD
Tubers
2017
Sweetmeats
2018

C. 2018

Cotula or Babuna Alpine Stocks

COTULA
anthemoides

the root stock is said to be used as a substitute for ginger Dr Dymock
commenting on this statement remarks The rhizome resembles the

subject to favour it The Revd A Carpuce says it is used
by the Santals

COTONEASTER, *Medik Gen Pl I 627*

[ROSACEÆ

Cotoneaster acuminata, *Linll Fl Br Ind Vol II 385,*

2019

Vern — *Ru rauns r us ruish* HIND

References — *Brand's For Fl 209 Gamble Mal Tinb 171*

Habitat — A deciduous shrub of the Himalaya from the Beas to Sikkim and occurring between 4500 and 13000 feet

Structure of the Wood — Hard like that of *C bacillaris* used for walking sticks

WOOD
2020

C bacillaris, *Wall Fl Br Ind, Vol II 384*

2021

Vern — *Ri ru l n l nu lehan khars luni rau reish reus rish stelu*

R

Habitat — A small deciduous tree of the Salt Range above 1500 feet of the North West Himalaya from the Indus to the Sarda between 5000 and 10000 feet and of Sikkim and Bhutan

Structure of the Wood — Wood smooth very hard close and used for making walking sticks

WOOD
2022

usually made of this wood and the sticks are used to support it to the plants from many points along the Himalaya This is the *Cotoneaster obtusa* alluded to in the Settlement Report of the Simla district in which it is said the hill tribes use the sticks as goads (*chalta*). The larger pieces are made into jampan poles axe handles &c Baden Powell suggests that it is suitable for turning

C microphylla, *Wall Fl Br Ind II 385*

2023

Cotton and Cotton Manufactures see the article *Gossypium* in Vol III

Fruit
2024COTULA, *Lin Gen Pl II 48*

Cotula anthemoides, *Lin Fl Br Ind III 316* COMPOSITE

2025

Vern — *Babana* PB HIND

C 2025

CRAMBE
cordifolia

The Cow Tree

MEDICINE.
Fowers.
2025

Hab. tat.—A small herbaceous plant found in the Gangetic plain, from Rajmahal and Sikkim westwards, to the Panjab.

Medicine.—It furnishes part of the official *babina*, which is heated with oil and applied externally in rheumatism, &c. Compare with *Anthems nobilis*, *Linn.*, A. 1185.

§ "The infusion is used as an eye wash, in most diseases of the eye (*Surgeon-Major C W Culthrop, M.D., Morar*).

Country Borage, see *Coleus aromaticus*, *Ben'l.*; LABIATÆ.

Cotyledon laciniata, *Roxb.*, see *Kalanchoe laciniata*, *DC.*

COUSINIA, *Cass.*, *Gen. Pl.*, II., 467.

2027

Cousinia minuta, *Bour.*; *Fl. Br. Ind.*, 359; COMPOSITÆ.

Syn.—*C. CALCOTRAPHIFORMIS*, *Jaub & Spach.*; *C. AYLENSIS*, *Bunge*.
Vern.—*Lakheri*, *po.*; *kand eri*, or *kand ari*, *Ps.*

Reference—*Stewart*, *Pb Pl.*, 125.

Hab. tat.—A small rigid herb, found in a wild state in some parts of the Western Panjab plains, and distributed to Afghanistan, Baluchistan, and Persia.

FOOD
2023

Food.—The young plant is used as a vegetable in the Salt range (*Stewart*).

Covellia glomerata, see *Ficus glomerata*, *Roxb.*; URTICACEÆ.

Cow itch or Cowhage, see *Macuna pruriens*, *DC.*; LEGUMINOSÆ.

Cownie, Kawrie or Cowdie Pine, commercial name for *Dammara australis*, see under *Dammara*, *Hopea*, and also *Canarium*, C. 273.

Cownie or Cowry, see *Shells*, also *Beads*, B 380.

2029

Cow Tree.—Many plants, with milky sap, receive the name of Cow Tree. Perhaps the only peculiarity that more especially justifies that name is when the sap contains very little Caoutchouc and is wholesome. The Cow Tree of most writers is *Brosimum Galactodendron*, to which Humboldt was the first to draw special attention. It is a member of the Bread-fruit family (*Artocarpæ*). Several fruitless efforts have been made to introduce this plant into India, see the *Indian Forester*, IX., 517.

Crab's Eye, see *Mela Azedarach*; also *Abrus precatorius*, A. 73.

Crab Tree, see *Pyrus Malus*, *Linn.*; ROSACEÆ.

Crabs, see *Crustacea*.

CRAMBE, *Linn.*, *Gen. Pl.*, I., 93

2030

Crambe cordifolia, *Steud.*, *Fl. Br. Ind.*, I., 165; CARYOPHYLLACEÆ.

Habitat.—A tall herbaceous annual, with leaves nearly a foot in diameter. Frequent in the North-West Himalaya, Quetta, Western Tibet, &c., at 10,000 to 14,000 feet.

FOOD
2031

Food.—The young leaves are, in the Salt range, eaten as a pot herb (*Stewart*), and in Baluchistan the root is eaten (*Stewart*).

C. 2031

Hawthorn The Bel Fruit of some Writers

CRATÆVA
religiosa.

CRATÆGUS, Linn, Gen Pl, I, 626

Cratægus Clarkei, Hook f, Fl Br Ind, II 384, ROSACEÆ

2032

A species of hawthorn met with in Kashmir, which may be viewed as intermediate in type between the two following species

2033

C. crenulata, Roxb, Fl Br Ind, Vol II, 384

THE HIMALAYAN WHITE THORN

Syn — C PYRACANTHA, Persoon; MESPILUS CRENULATA, Don

Vern — *Gingaru gianru*, HIND, *Gengaru*, PBReferences — *Roxb, Fl Ind, Ed C B C, 406* *Voigt Hort Sub Cal*
198 Brandis, For Fl, 208 *Gamble Man Trib 170* *Dalziel & Gibb,*
Bomb Fl Supp 132 *Baden Powell, Pb Pr, 576* *Drury, U Pl*
208 *Catfour, Cyclop, 836* *Treasury of Bot, 344*

Habitat — A large spinescent shrub of the Himalaya, from the Sutlej to Bhutan, found at altitudes from 5 000 to 8,000 feet, but in Kumáon at 2 500 feet

Structure of the Wood — White, hard, very close and even grained, used as axe handles, staves, &c

WOOD

2034

C. Oxyacantha, Linn, Fl Br Ind, II, 383

2035

THE HAWTHORN

Vern — *Ring, ringo ramnia pingyat, or pinyat, phindak, patákhān*
ban sanlyi sursinlyi or singly PB HIMALAYAS, *Ghwansa, or ghwardsa,*
TRANS-INDUS *Durana* AFGH

Habitat — A small tree (20-30 feet) met with in the North West Himalayas from Quetta to the Rávi basin Cultivated eastwards near villages, and in Afghanistan is a favourite tree planted near tombs

FOOD
Flowers

2036

Fruit

2037

WOOD

2038

poses as the preceding

CRATÆVA, Linn, Gen Pl, I, 110

Cratæva religiosa, Forst, Fl Br Ind Vol I, 172, CAPPARIDÆÆ

2039

Syn — CAPPARIS TRIFOLIATA, Roxb, C ROXBURGHII, Ham, C NUR
VALA, HamVern — *Barna barun b lāsi, bila biliana* HIND, *Barun, tikto-shak*
BENG, Tailadu, bunboronda, MECHI, Purdong, LECHA, Barna,
darnahi, PB, RAJ, Bela, bel, C P, Vāyavarna bhatavarnā hāda
varna, kumla varuna karvan, BOMB, Kumla karwan, MAR, Māra
lingam, varulinga, harcala, TAM, Nirvāla vitusi KAN, MAL,
Ustia usiki, usiki mānu ulimidi urimidi urumitti, tella ulimidi tella
ula, TEL, Nirujani COORG, Āadet katat BURM, Varuna asma
vighna SANS Roxburgh says that it is the *Tikta-shaka* of Sanskrit
writersHistory — L
Ægle Marmel
Cratæva Marm
the same vern
ances and in

HISTORY.

2040

C. 2040

CRATÆVA
religiosa.

Cratæva or Bel.

HISTORY

Ind.), under *Cratæva religiosa*, gives the following vernacular names as

that the medicinal leaves sold at the present day are those of *Ægle* rot of *Cratæva*.

A brief review of the confusion which exists, in the literature of India, between *Cratæva* and *Ægle*, will be given.

scribes *Ægle* Marmelos, quotes the same botanical description, the same name, and the same fruit.

Ainslie's meaning when he says "The species in question I have never seen," nor can we presume that he was labouring under the idea that *Cratæva* Marmelos was a different plant from *Ægle* Marmelos, seeing that in his two articles upon the medicinal product discussed he quotes the

name for the plant. It is worth noting that the use of the word "Bengal" practically implies that the Madras supply was imported from that province. Roxburgh wrote his *Flora Indica* about the same time as Ainslie produced his *Matéria Indica*, and the latter author frequently admits that he had seen the MS. of Roxburgh's work. In the *Flora Indica* it is stated of *Ægle* Marmelos that it is a native of the mountains of Coromandel, "and is also found sparingly, in the low lands." Is it thus possible that, before the *Bel* fruit was cultivated to the extent it now is, *Cratæva* took its place (at least as a medicinal yielding tree) and was displaced from popular favour, the *Bel*, as we now know it, receiving many of the older names? If so, the botanical name *religiosa* may rest on a stronger basis than the mere fact that the tree is grown near temples and tombs. Lisboa says: "So far as my enquiries go, it is not mentioned in Hindu religious books, nor used in their worship." But does it not rest on the evidence in classical literature of descriptions that refer unmistakably to *Ægle* Marmelos, or simply on the words *Bilva*, *Bel*, &c.

The Bel Fruit and *Cratæva***CRATÆVA religiosa.**

Botanical evidence would point to *Ægle* being almost insular in its character, and it may be doubted if it is even grown to any extent in the present day beyond the limits of peninsular India, it does not succeed, for example, in Northern Panjáb. But *Cratæva* is more continental in its distribution, and is therefore more likely to have been known to the ancients.

The writer's object, however, in suggesting a doubt regarding the bel fruit will be gained if greater attention is paid to the two most useful plants—*Ægle Marmelos* and *Cratæva religiosa*.

References—*Roxb., Fl. Ind. Ed. C.B.C., 426* Brandis *For Fl., 16* *Kurs., For Fl. Burm., I., 66* *Gamble, Man. Timb., 15*, *Dals & Gibs, Bomb. Fl. 8*, *Stewart, Pb. F. Elliot, Flora Andhrica* *Mat Ind., II., 86, 19* *deen Sheriff, Suppl. F. 115 323* *Dymock, Ma. Drugs 13, Pl. and Dr.*

HISTORY

tree near temples and tombs

Varieties—The *Flora of British India* refers the forms of *Cratæva* to two varieties, which seem in a measure to correspond with the species of that genus alluded to by authors on Economic Botany.

Var 1st, Nurvala *Leaflets ovate-lanceolate, taper-pointed berry ovoid-oblong*—This appears to be the *C. Nurvala* of Hamilton and the *Nurvala* of Rheede. Dalzell and Gibson say this form is the true "*Varvanna*" and is met with in the Caranjah Hill, Warree country. Wight and Arnott (in their *Prod. Floræ Penins. Ind. Or.*) speak of it as "frequent in rich moist soil on the banks of ditches and rivers on the Malabar coast, also in Mysore, where it grows to the height of 15 or 20 feet." They also state that it is the *C. Tapia, Burm. (in part)*, and also the *C. inermis, Linn. (in part)*.

VARIETIES.

Var 1st,
Nurvala
2041

With the exception of the middle paragraph (p. 111) in the *Marmelo* (p. 459) the *Ægle* in Sanskrit that writes of *Ægle* the most pellucid glands in the tissue would be proof positive of the leaf not being *Cratæva*. Ainslie further states however, of his plant that "the root, as

ub-aromatic and bitterish taste, quality." He further observes of Rheede, and the *lunu-arua* *atalogue of Ceylon Plants*, affirms the next variety. This is therefore, the only serious mistake made by Ainslie in his attempt to distinguish the two forms of *Cratæva*.

Var 2nd, Roxburghii *Leaves small ovate-lanceolate abruptly acuminate, berry globose*—This is *C. Roxburghii, Br.* and the *C. odora religiosa*, and *unilocularis* of Hamilton, and the *Capparis trilocularis* of

Var. 2nd
Roxburghii
2042

CRATÆVA
religiosa

Forms of Cratæva

VARIETIES

Roxburgh, Dalzell and Gibson say it is common on the banks of the Nerbudda, Roxburgh, that it is

Varana, Sans. He further calls it the "Smooth Tapia or Garlic Pear," the latter name, as he explains,

tea-spoonful twice or thrice daily." Sir Walter Elliot alludes to this form in his *Flora Andhrica* (pp 180, 185, 187), and gives it the Telugu names of *ulimidi*, *usiki manu*, *tella-ulimidi*.

It may be worth pointing out that it is the leaves of variety Nurvala that he does not tell us whether or not the natives of India were in his

another Jamaica species, *C. gynandra*, he says "that the root blisters like cantharides."

These facts are of the greatest importance, in the confirmation which they afford to the opinions, expressed on a further page, by Dr. Moodeen Sheriff, as to the rubefacient properties of the leaves. It would be instructive to learn whether these properties were common to both forms of *C. religiosa*, or only possessed by the form which bears Dr. Roxburgh's name. There is also another point of some importance. Ainslie in his article on "*Cratæva Marmelos*" (*Mat Ind*, I, 86), which is clearly an account of *Ægle Marmelos*, and again, in the 2nd paragraph of his article on "*Cratæva religiosa*," refers to a resin found within the fruits, which he regards as of great value "in clearing foul ulcers." It is also used, he informs us, "in the arts as a cement." This resin and cement is well

Cratæva becomes possible

Gum and Dye — "Aitchison states that at Jhelum the fruit is mixed with mortar to form a strong cement, and the rind as a mordant in dyeing" (*Stewart*)

Medicine — From what has been said it may be inferred that some doubt still exists as to whether the medicinal products of *Cratæva* can be spoken of as afforded by the one species or two species. The writer must

* A name which does not appear now to be in use in Hindustan, although mentioned by the older writers.

The Nurvala

CRATÆVA
religiosa.MEDICINE.
Bark
2048
Leaves,
2049

common complaint of a somewhat obscure nature. The leaf-juice is given in rheumatism in the Concan in doses of $\frac{1}{2}$ to 3 tolas, mixed with cocoanut juice and *shu*. In caries of the bones of the nose the leaf is

Juice,
2050

lingai, lam.," that "the leaves, bark, and roots are used medicinally."

will
irras.
not

light-
boards, combs, and in turnery. In Trichinopoly it is also used "for making planks and as firewood"

FOOD.
Fruit.
2052
WOOD.
2053

C. 2053

CRINUM.

Famine Food

CRATOXYLON, *Blume, Gen Pl, I, 166*

[HYPERICINÆE

- 2054 *Cratoxylon formosum*, *Benth et Hook, Fl Br Ind, I, 258*,
A large tree, met with in the Andaman Islands, yields a useful timber,
but the tree is rare (*Kurz, For. Fl. Burm, I, 84*).

- 2055 *C. nerifolium*, *Kurz, Fl Br Ind, I, 257*

Vern — *Baidya* BURM

- WOOD
2056 Habitat — A moderate-sized tree, found in Chittagong and Burma
Structure of the Wood — Dark grey, hard, close-grained. According
to *Kurz*, it is used for building purposes, for ploughs, handles of chisels,
hammers, and other implements

CRESSA, *Linn, Gen Pl., II, 881*

- 2057 *Cressa cretica*, *Linn, Fl Br Ind, IV., 225*; CONVOLVULACEÆ

Vern — *Gul*, *Sind*, *Khardi*, *BOMB*, *Chavel*, *NASIK (BOMB)*, *Uppu*
sanna TEL (Sir Walter Elliot remarks regarding the above Telugu
name that "the plant is so called from frequenting salt lands near the sea,
where it has much the look of young *Chenna* or *Cicer*")

References — *Roth Fl Ind, Ed. C B C 265, Dals and Gids, Bomb*
Fl 162, Voigt, Hort Sub Cal, 363, Grah, Cat Bomb Pl 133, Dym-
mock, Mat Med W Ind, 2nd Ed, 569, Walter Elliot, Flora And
hrica 186, Bomb Gaz (Cutch), V, 27, Stocks Account of Sind;
Aitchison, Cat Pb and Sind Pl p 98, Sakharani Arjun, Bombay
Digs, 93

Habitat — A small erect shrub, common throughout the warmer parts of
India especially near the coast from Multan, Baluchistan, and Sind, through
Gujarat southwards to the Coromandel coast, and distributed to Ceylon
Appearing in the fields after the rains

- FOOD
Seeds
2058 Food — *Stocks* mentions that in Sind the seeds of this plant are
mixed with wheaten flour
Dymock mentions that in
aten during the famine of

MEDICINE

- 2059 Medicine — *Dr Sakharani Arjun* says "It is used as a tonic and
is believed to possess expectorant properties" *Dr Dymock* remarks "
"It is found in Greece, and is supposed by some to have been one of the
two kinds of *ἀνθλῆς* described by *Dioscorides*"

- 2060 CRINUM, *Linn, Gen Pl, III, 726*

A genus so named from the Greek *κρίνον*, a lily (*Theophrastus*) It con-
tains about sixty species mostly natives of the tropical regions in the old and new

Toxicarium.

CRINUM
asiaticum.

Crinum amœnum, Roxb, *Fl Ind*, Ed. C.B.C., 283; *Herbert*, 255;

2061

Vern.—Gocinda, SYLHET

References.—*Drury*, *Fl. Ind*, III., 454; *Voigt*, *Hort. Cal*, 590

Habitat.—A native of Nepal, Sylhet, and Burma, flowering nearly all the year, but mainly in the hot and rainy seasons; the flowers are large and white.

C. asiaticum, Linn.; var. *toxicarium*, *Herbert*, *Bot. Mag.*, 1073.

2062

Syn.—*C.* ...

Roxb;

Bot. M

V. p

Vern.—Chundar, kanwal, pindar, kanmu, HIND, Nagdown (according to Dymock), BOMB, Nagdamani, Guz, Nagadivana, MAR, Nagin-ka patta, Duk, Bara kanur, nagdown, boda kanod (gac-honar-patta, according to Bidie, Dymock, &c), BENG, Vishi-mungil, TAM,

Koyangi,

Ian sy lan,

Vishaman

References.—Roxb, *Fl Ind*, Ed. C.B.C., 283, *Voigt*, *Hort. Sub. Cal*,

Drury, III., 454, *Drury*, I., 11, *Drury*, III., 19, *Drury*, Cal. Ind

Habitat.—A fairly abundant cultivated plant, its erect stems with their crown of large graceful leaves forming almost a characteristic feature of

erect stem in distinguishing it from *C. defixum*, and he expresses the opinion that it may be a native of Ceylon. Speaking of that region Thwaites remarks that "it is very abundant on the sea-coast of the island," and "frequently planted as a fence for native gardens near the sea."

Although thus not established in the synonymy of the Economic Botany give the *aticum*. This idea has been probable future investigation may relegate to *C. defixum*, *C. amœnum*, or *C. pratease* much of what is here given under the popular name *C. asiaticum*.

Medicine.—Ainslie wrote in 1826: "The succulent bitterish leaves of this plant, which are about 2 inches broad and 3 feet long, the natives bruise and mix with a little castor-oil, so forming an application which they think useful for repelling whittows, and other inflammations that come at the end of the toes and fingers, the juice of the leaves is employed

MEDICINE
Leaves
2063
Juice
2064

C. 2064

CRINUM
pratense

Toxicarium—a useful Emetic.

MEDICINE
Root
2005

for the ear-ache in Upper India. In Java, by Horsfield's account, this plant is reckoned one of the most satisfactory emetics the inhabitants have "It is the root (? bulb) chewed that is the emetic, provided a little of the juice is swallowed" Sir William O'Shaughnessy, who wrote some 20 years later, says: "into a paste, a emetic after a

Extract
2006

phoretic, we have never known it to occasion any untoward symptoms. The dried sliced roots are also an efficient emetic, but require to be given in double the dose of the recent article" The extract, whether watery or alcoholic, is very uncertain in its action. In the form of a syrup it may probably be found to retain the native principles of the recent plant. The tincture of the fresh plant does not succeed, doubtless in consequence of the large quantity of spirit counteracting the emetic effect by its stimulating energy.

These two passages express all that has since appeared, as for example, in the *Pharmacopœia of India*, Drury, Murray, K. L. De, and indeed most subsequent writers, repeat in other sentences the same facts. Dr. Dymock adds "I have not met with any account of this drug in native works on

Bulb
2007

... relating the juice being used for ear-ache, Dr. Dymock gives as a footnote "A well known popular use of the plant, the leaves are slightly roasted, and the juice is then expressed and a few drops poured into the ear"

The bulb of the so-called *Crinum asiaticum* is made official in the

... (Dr. H. W. Hill, Manbhoom)

2008

Crinum defixum, Ker (and of Gawl), *Herbert*, p. 255; *Bot. Mag.*, [2208.

Syn—*C. ASIATICUM*, Roxb (non Linn), *Fl Ind*, Ed C B C, 283; *C. ROXBURGHII*, Dals, *Fl Bomb*, 275, BELUTTA POLA TALLY, *Rhede*, XI, t. 38, *RADIX TOXICARIA SECUNDA*, Rumph, VI, 156

Vern.—*Suk darshan*, BENG, Nagdown, BOMB; *Kesar chettu*, TEL; *Hintolabo*, SING (according to Ainslie)

References—*Dals & Gibs*, *Bomb Fl*, 275, *Lisboa*, *U Pl Bomb*, 204

Habitat—A native of the Concan, of Coromandel, and of many parts of Bengal, as, for example, the Sunderbunds. Flowers large, sessile, white, fragrant during night, flowering time, the close of the rainy season. Dalzell and Gibson say it is common on the banks of

MEDICINE
2009
2070

C. pratense, *Herbert*, *Amaryll.*, 256.

Syn—*C. LONGIFOLIUM*, Roxb, *Fl Ind*, Ed C B C, 283; *C. LAURIFOLIUM*, *Herbert & Roxb*; *C. ELEGANS*, *VENUSTUM*, and *CANALIFOLIUM*, *Carey*

Vern.—*Pa tang*, BURM

References—*Voigt*, *Hort Sub Cal*, 590, *Bot Mag*, t. 2392 and 2122

C. 2070

The Common Crocodile

CROCODILUS
palustris.

Habitat.—A native of the interior of Bengal, Sylhet, Pegu, &c, flowering time the rainy season. Flowers large, white, fragrant. A variable plant, some of the names given above belonging to what may prove recognisable forms.

2071

Crinum, sp. (found in Chutia Nagpur)

2072

Mr. C. B. Clarke writes of this plant that he is unable to name it and presumes it may be an undescribed species. In that case it should bear the discoverer's name—the Rev A. Campbell. Mr. Clarke also informs the writer that he has collected another species in the tanks of Chutia Nagpur which flowers in November; he views this as distinct from the common Sunderband species, which flowers in May.

Vern.—*Sikyom baha*, SANTAL.

Habitat.—High and dry situations in Chutia Nagpur, flowering during the hot season before the leaves appear. In some respects, this resembles *C. latifolium* as described in Roxburgh's *Flora Indica*.

2073

Medicine.—Mr. A. Campbell says "The bulb is sometimes as large as a good-sized turnip, and of the same shape. A decoction prepared from it is given internally and pounded and made into a paste, it is also applied externally by the Santals in dropsy. It is used for the diarrhoea of cattle."

MEDICINE
Bulb
2074

C. zeylanicum, Linn., Wright, Ic. 2019-20

2075

Syn.—*C. ORNATUM*, Herbert. *C. ZEYLANICUM*, Roxb. *C. LATIFOLIUM*, Roxb. *C. MOLUCCANUM*, Roxb. *C. HERBERTIANUM*, Herb., p. 253, also Wall., *Pl. As. Rar.*, 2 p. 145.

Vern.—*Sukh-darsan*, BENG. *Gadambikanda*, BOMA. *Goda manil*, SING.

Re.—

Habitat.—A very variable plant, some of the above synonyms corresponding to well marked varieties, which in a work on economic products,

conference

Medicine.—Dymock remarks of this species "The bulb is extremely acrid and is used for blistering cattle, a slice being bound upon the skin. When roasted it is used as a rubefacient in rheumatism."

MEDICINE.
Bulb
2076

CROCODILE (CROCODILUS, Cuv.)**Crocodylus palustris, Less.**

2077

THE COMMON CROCODILE, often vulgarly called in India, the Alligator—an American Reptile.

C. 2077

CROCUS sativus.

The Crocodile ; Saffron.

There are apparently two other species besides the above met with in India, viz, *C. porosus*, *Schneid*, and *C. trigonops*, *Gray* The long snouted Gaval lives on fish and turtles, and frequents the rivers of India along with the Crocodile

Vern.—*Magr, kumhr*, HIND, *Sisan*, SIND

Habitat.—Found throughout India and Ceylon, affecting rivers, lakes, marshes, and even the sea coast It may be recognised by its shorter and broader snout than that of the Gaval, and by the first and the fourth tooth of the lower jaw fitting into the upper

Although held sacred in many parts of India (and sometimes even

great size, being from 15 to 30 feet in length, and although it is reported to eat the dead bodies thrown into the rivers, it lives mostly on live animals, taking human beings when pressed for other food

Economic Products —OIL, SKIN, MUSK, and FLSSII.

Crocodile Flesh —It many parts, Crocodile flesh is said to be eaten or

Africa appear to regularly extract Forbes Watson, in his *Industrial* sample of this substance procured

from Travancore

Crocodile Oil.—The oil of the Indian Crocodile contains a larger quantity of solidifiable fat than either neat's-foot or any fish-oil It is prepared by the Sanif tribe, in the Panjáb, who eat crocodile flesh, and is also said to be procurable in abundance at Agra (*Spons' Encyclop* 5136)

CROCUS, Linn, Gen Pl, III, 693

This is the *κροκος* of *Dioscorides* It is not alluded to by the earl er Sanskrit writers, but Arab an authors speak of it as cultivated in the tenth century at Darband and Ispahan and Chinese writers state that it was introduced into their country by the Muhammadans in the Yuen dynasty (A D 1280)

Crocus sativus, Linn, Royle, Ill Him Bot, t 90, IRIDEE.

SAFFRON

Vern —*Yafran*, BENG ; *Kesar, safran*, HIND, *Safran, kessar, kecara*, BOMB *Kecara*, MAR, *Keshar*, GUZ, *Kuikuma, kasmirayanma* (*Ainslie*), *kumkuma* (*Dutt*) *saurab* (*Dymock*), SANS ; *Zaafaran*, ARAB, PERS, *Kungumapu*, TAM, *kunkum apare*, TEL, *Thanwai* (Mr Oliver, Forest Officer in Burma, informs the writer that this is the

Re,

Sheriff, Supp Pharm Ind, 118, U C Dutt *Ind A* *Pharmacog*, 663
11, 274
Murray, 11
14; *Baker*
118, 118, 68,
118, 118

monds, Trop Agri, 379

Saffron, Indian Crops	CROPS.
Habitat.—The European supply of this plant comes from France.	SAFFRON.
	DYE. 2084
highly thought of as a remedy for catarrhal affections of children, and is used in certain Indian dishes as a colouring agent. Mullahs (priests) make a kind of ink with this substance with which they write charms (<i>Dr Emerson</i>). In over doses it is generally reported to act as a narcotic poison. Ainslie gives perhaps the most complete account of the native uses of this drug, and of the opinions which prevailed among	MEDICINE 2085 2086
torius)	
Chemistry —§ "The colour of saffron is due to the presence of a glucoside polychrom, which is decomposed by acids, with the formation of a new colouring principle <i>Crocin</i> " (<i>Prof Warden, Calcutta</i>). For full particulars as to the chemistry of this drug see the <i>Pharmacographia</i> , p 666.	CHEMISTRY 2087
Trade in Saffron —The imports of foreign saffron were in 1882-83, 226 cwt valued at R4,25,224, and in 1886-87, 268 cwt valued at R5,50,383. Of the Indian imports the bulk comes from France	TRADE 2088
CROPS.	
An important feature of Indian Agriculture is the fact that, through the presence of extensive montane tracts, India possesses considerable areas that are under temperate influences, as well as vast expanses that are purely tropical. Between these two conditions almost every possible gradation exists in which the tendency to extreme humidity or extreme aridity modifies the general character. From this point of view alone	2089
sometimes three harvests a year. This is modified in certain provinces through the rains not occurring at the same period. Thus, in Bengal, Bombay, the greater part of the Central Provinces, and in Berar, the rains	

CROPS

Indian Crops.

occur in June, July, August, and September, being preceded by the hot

seen that to study the crops of India, the closest attention must be paid to this shifting of

In the regions

marked crops

The temperate mountains within these regions have accounts of the
des),
son
hus
art-
own
of
be
hief
ical

place in this work

2090 1st, CEREALS — This includes Wheat, Rice, Oats, Barley, Indian-corn, Millets (various kinds), and Coix (Job's tears) (Conf. with Cereals)

2091 2nd, PULSES — Such as Gram, Peas, Beans, Lentils, &c (Conf. with Pulses.)

2092 3rd, GRASSES — Such as Rice, &c (Conf. with Gramineae)
or eaten boiled
he GRAMINEAE
ains)

2093 4th, SPICES AND CONDIMENTS — Turmeric, Ginger, Cumin, Coriander, Caraway, Pepper, Betel-leaf, Capsicum, Cardamum, &c, &c (Conf. with Spices)

2094 5th, STARCHES AND SUGAR — Sugar-cane, Arrow-root, Sago, &c (Conf. with Starches)

2095 6th, GARDEN PRODUCTS AND VEGETABLES — Potatoes, Yams, Colocasia, Cabbage, Gourds, Melons, Cucumbers, &c, &c (Conf. with Vegetables)

The above might be grouped as edible products, but there are other crops some of them of even great importance, such as—

2096 7th, FIBRES — Cotton, Silk, Jute, Sunn-hemp, and many others, the fibre from Hibiscus cannabifolius being, after sunn-hemp, the next most important of fibre crops (Conf. with Fibres)

2097 8th, DYES — Indigo, Safflower, Al (Morinda tinctoria), Madder, &c (Conf. with Dyes and Tans)

2098 9th, NARCOTICS — Opium, Ganja, Tobacco, Tea, and Coffee. (Conf. with the separate accounts of each of these products and with the article Narcotics)

Crops; Sunn-Hemp.

CROTALARIA
juncea.

10th. OIL-SEEDS—Ground-nut, Rape, Mustard, Cotton-seed, Linseed, Opium-seed, Castor-oil, Gingelly or Sesame oil, &c (Conf. with Oils)

2099

These are the principal crops of India, but the agriculturists have

CROTALARIA, *Gen. Pl.*, I, 479.

2100

A genus of plants closely allied to the Broom, the generic name being derived from the Greek *κροτάλον* (a castanet), in allusion to the rattling noise made by the loose seeds within the inflated pods. This same idea, according to Sir Walter Elliot, is implied by the Sanskrit name *Ghantar atamu*.

Crotalaria Burhia, *Hamilt.*, *Fl. Br. Ind.*, II, 66, LEGUMINOSÆ

2101

Vern.—*Sis, sissai, mesut, pola, khippi, buta, kkep, khip, khif bhata, bti, lathia, kharan kauridia, PB, Ghagari, MAR, Ghugharo, Güz, Druuna, SIND*
References.—*Pl. Br. Ind.*, I, 479, II, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935, 936, 937, 938, 939, 940, 941, 942, 943, 944, 945, 946, 947, 948, 949, 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, 984, 985, 986, 987, 988, 989, 990, 991, 992, 993, 994, 995, 996, 997, 998, 999, 1000

Rajputana Gaz., 301

Habitat.—A low under-shrub, abundant in the sandy plains of Sind Panjab, Rajputana, and Cambay, ascending to 4000 feet in altitude.

Fibre.—Is said by Mr. Baden Powell to yield a good fibre for cordage, used, to some extent, in the Panjab in place of the *Sunn-hemp* (*C. juncea*) of other provinces

FIBRE.
2102

MEDICINE.

Branches.

2103

FODDER.

2104

2105

Medicine.—The branches and leaves are used as a cooling medicine

Fodder.—The Rajputana Gazetteer states that the plant is much valued as a fodder.

C. juncea, *Linn.*, *Pl. Br. Ind.*, II, 79

SUNN OR SUNN HEMP OR INDIAN HEMP, FALSE HEMP, BROWN HEMP, BOMBAY OR SALSETTE HEMP, WLCKOO NAR (OR TRAVANCORE FLAX), JUBBULPUR HEMP, &c, &c

Syn.—*C. tenuifolia*, *Roxb.*

Vern.—*Sun, sanai, sani* (or *sun, shon*), *HIND, BENG, Ausá, suna, Assam, San, Al, - - -*

ing to Sir Walter Elliot), *SANS*

According to some writers the name *Ambadi* or *ambari* is, in Western India, given to this plant, but it seems probable that that name should be restricted to *Hibiscus cannabinus*. Indeed, it has been found difficult to arrive at any definite idea regarding the present area under *sunn-hemp* cultivated on from the fact that the above *Hibiscus* appears to be confused with it. In Bengal, and indeed in some parts of the N-W Provinces,

CROTALARIA juncea.	History of Sunn-Hemp
FIBRE.	<p>nus are separately reported It would thus appear that the term "Bombay hemp" is often, though incorrectly, given to the Ambadi fibre, <i>Hibiscus cannabinus</i> It is thus unfortunate that, in modern commerce, the term "hemp" should ever have come to be applied to any but the true hemp plant, as, by this usage, widely dissimilar products have been almost hopelessly confused The <i>sunn</i> is a bush closely allied to the English broom or the Indian <i>dal</i>, while the <i>ambari</i> is a <i>Hibiscus</i> or cotton-looking plant with sharply-cut leaves not unlike those of the hemp plant,—hence the specific name <i>cannabinus</i> The true hemp has its nearest affinity, of fibre yielding plants, in the common nettle The hemp fibres thus afforded by these three plants have little or nothing in common</p> <p>References —<i>Roxb, Fl Ind, Ed C B C, 545, Voigt, Hort Sub Col,</i></p> <p>Habitat —The <i>Flora of British India</i> gives the habitat of this plant as "Plains from the Himalaya to Ceylon, but often planted for its fibre" The writer is not aware of <i>Crotalaria juncea</i> having been recorded as found in a wild state anywhere in India although it may sometimes exist as an escape from cultivation Kurz says of <i>C juncea</i> in Burma "like wild along the banks of the larger rivers, especially the Irrawaddy," and Griffiths that <i>C. juncea</i> is met with in Afghanistan Roxburgh describes a form (by modern botanists reduced to the present plant, viz, <i>C. tenuifolia</i>) which he states is a native of Coromandel Many writers however, familiar with the living plants, still affirm that <i>C. juncea</i> and <i>C. tenuifolia</i> are distinct They seem at least to be cultivated recognisable states which, owing to the reputed superiority of the fibre of <i>C tenuifolia</i>, it might, for historic point reported the as next found be cultivated</p>

History of Sunn-Hemp

CROTALARIA
juncea

to this day, although as yet it has not been reported as found anywhere between these remotely distant regions. At the same time *C. juncea* is he Panjab and Sind which yields a fibre a superfluous.

SUNN (or SAN) HEMP FIBRE.

Under the heading *Cannabis sativa* the suggestion has been offered that the Greek and the Latin *cannabis* may have been derived from the

FIBRE.

2106

to be the same as the references to hemp in Sanskrit literature is that Even ication, r than of flax

of hemp, such names as *shesh* implying an intoxicating power—a property of the hempen fibres possessed alone by *Cannabis sativa*. The *sana* fibres of the *Sann* are a fact that closely agree with

of hemp, such names as *shesh* implying an intoxicating power—a property of the hempen fibres possessed alone by *Cannabis sativa*. The *sana* fibres of the *Sann* are a fact that closely agree with

CROTALARIA
juncea.

History of Sunn-Hemp

FIBRE.

the name for gr
for the *ksliaur*
it was made
made, the *patt*
probability the sunn hemp made garment Later writers speak of *sana*

The hill tribes of the North-West Himālaya weave a proportion of their clothing of hemp, but although the plant springs up wild all over the plains

Sacred
Threads
2108

sana has been carried, at the present day, to the extent of violating even this injunction. Lisboa (*Bombay Useful Plants*, p 290) states "It is a native of India, and is cultivated in the hills and mountains, where it is used for making threads."

it appears
to justify
can it be
even said to be a native of Persia, though it may possibly be of China, as it is of Russia, Siberia, and Kirghiz. On the other hand, *Crotalaria juncea*, while met with to-day almost exclusively under cultivation, would appear to be a native of India, and possibly also of Central Asia, many other species of the same genus being indigenous to those countries. This fact added to the conclusion

is and
whole
general
as the
There
itively
mited
o not
but
prefer to cultivate *sunn* hemp (*Crotalaria juncea*) or *san-pai* (*Hibiscus cannabinus*) for the cordage and sacking required for agricultural purposes. There is still a further consideration, and one of some importance, viz., that on the plains of India the hemp plant does not produce fibre of any value. Unless, therefore, we are to presume that it has degenerated, or that the climatic conditions of India have altered, the ancient people of the plains were not likely to have obtained their *sana* fibre from *Cannabis sativa*.

We may conclude this brief historic review of the hemp plants by giving the opinions that prevail regarding the origin of our word "hemp."

Cultivation of Sunn-Hemp

CROTALARIA
juncea

FIBRE

Royle in his *Fibrous Plants of India* traces hemp from *sana*. Speaking of *sunn*-hemp, he says "Its name, Shanapam or Janapa on the Madras side, is not very unlike Canapa, Hampa, Hennip, and Hanf. From these we derive our own name 'Hemp'." In Mysore it is known as *sanabu* and may be found in the Veda in all parts of India and in the Greek and Latin, and *kannab* in Arabic.

CULTIVATION.

CULTIVA-
TION
2109

Sunn is grown by itself or at times is cultivated in strips or around the margins of fields. It is never cultivated as a mixed crop. Throughout India as a whole it is a *kharif* crop,—that is to say, it is sown about the commencement of the rains and cut at the end of September or beginning of October. It is thus off the ground to allow of being followed by a *rahi* crop in the same year. But in some parts of India there are two crops of *sunn* hemp. Thus in the Thana District of Bombay it is sown in November after the rice harvest, and the stalks are pulled up by the root in March. "It is also sown as a rainy season crop in sandy soils" (*Gas*, *XIII*, *I*, 290). This system has prevailed in

Or Hove, writing in
"reached to the height
d that it was sown
een gathered in"

In Kolaba it is
he stalks are up-
t and harvested in

December by being cut when the plants are full grown. In Poona it is sown in July and ripens in October. In the Central Provinces and the North-West Provinces it is a *kharif* crop, being sown with the advent of the rains, but in Bengal it is sown a little earlier, namely,—from the 15th April to 15th June, in Madras the sowings take place even still earlier. In the experiments performed at the Saidapet farm Madras, *sunn* was sown on the 2nd of February. In the *Ain-i Akbari* the plant is described

mean period of sowing is about the beginning of the rains (or in June),
it and occupies the soil for
in view of the possibility of
throughout the whole year
fect this varying period of
fibre produced. Indeed, it
other crops sown at two or
more seasons each year) there may be different cultivated forms of the
plant produced as the result of ancient cultivation. We are ignorant of
this subject, and it seems desirable that a thorough investigation should
be made. Although, as stated, everything points to *sunn* hemp being a

CROTALARIA
juncea.

History of Sunn-Hemp

FIBRE.

the name for gr
for the *kshaur*
it was made
made, the *patt*

lution in popular opinion took place until (as in the present day) *sana* and

seeing that, as far as
sypium (cotton) is tri
(*Book II*, 44) we have
of the Brahmin must l
strings, that of a Cshatriya of *sana* thread only, and that of a Vaisya of
woollen thread" It is believed that the substitution of cotton for the
sana has been carried, at the present day, to the extent of violating even
this injunction Lisboa (*Bombay Useful Plants*, p 290) states "It

to be a wild state over the greater part of India there is little to justify
China, as
ia juncea,
appear to
be a native of India, and possibly also of Central Asia, many other

that on the plains of India the hemp plant does not produce more of any
value Unless therefore, we are to presume that it has degenerated, or
that the climatic conditions of India have altered, the ancient people of
the plains were not likely to have obtained their *sana* fibre from *Cannabis*
sativa

We may conclude this brief historic review of the hemp plants by
giving the opinions that prevail regarding the origin of our word "hemp."

Cultivation of Sunn-Hemp

CROTALARIA
juncea

Royle in his *Fibrous Plants of India* traces hemp from *sana*. Speaking of *sunn*-hemp he says: "Its name, Shanapam or Janapa on the Madras side, is not very unlike Canapa, Hampa, Hennip, and Hanf. From these we derive our own name 'Hemp'." In Mysore it is known as *sanabu* and in Ceylon as *hanz*. On the other hand, the root of the word *an* or *ang*

FIBRE.

Greek and Latin, and *kannab* in Arabic.

CULTIVATION.

CULTIVA-
TION.
2109

Sunn is grown by itself or at times is cultivated in strips or around the margins of fields. It is never cultivated as a mixed crop. Throughout India as a whole it is a *kharif* crop,—that is to say, it is sown about the commencement of the rains and cut at the end of September or beginning of October. It is thus off the ground to allow of being followed by a *rabi* crop in the same year. But in some parts of India there are two crops of *sunn* hemp. Thus in the Thana District of Bombay it is sown in November after the rice harvest, and the stalks are pulled up by the root in March. "It is also sown as a rainy season crop in sandy soils" (*Gaz.*, VIII, I, 290). This system has prevailed in Thana and Surat. In the latter place, the native name for *De. Ho.* is *1787*, says:—

stalks are uprooted in March. In Koinapur it is sown in August and harvested in December by being cut when the plants are full grown. In Poona it is sown in July and ripens in October. In the Central Provinces and the North-West Provinces it is a *kharif* crop, being sown with the advent of the rains, but in Bengal it is sown a little earlier, namely,—from the

mean period of sowing is about the beginning of the rains (or in June), *sunn* hemp may be sown in almost any month and occupies the soil for 4½ to 5 months. This is an important feature in view of the possibility of securing a continuous supply of fresh fibre throughout the whole year. It remains to be ascertained, however, what effect this varying period of cultivation has on the quality and quantity of fibre produced. Indeed, it is probable that (as is the case with rice and other crops sown at two or more seasons each year) there may be different cultivated forms of the plant produced as the result of ancient cultivation. We are ignorant of this subject, and it seems desirable that a thorough investigation should be made. Although, as stated, everything points to *sunn* hemp being a

CROTALARIA

juncea

Cultivation of Sunn Hemp

CULTIVATION
OF
FIBRESoll
2110

native of India, it may be doubted if the plant has ever been found in a truly wild state. And the existence of distinct cultivated forms might not only help to confirm the opinions given of an ancient cultivation, but might also establish the superiority of certain crops over others for textile purposes. To what extent the form *C. tenuifolia* is cultivated is not known still less do we know how far it affords the superior *sunn* hemp referred to by writers on this subject.

Nature of the Soil recommended for Sunn hemp—It requires a light but not necessarily rich soil, and it cannot be grown on clay. It is therefore sown on the high sandy lands less suited for the more important crops. This is the opinion which prevails in Bengal, but Messrs Duthie and Fuller, writing of the North-West Provinces, say "Authorities differ as to whether a rich soil is necessarily required, and y in the soil is necessary to et it cannot be contested that st any other crop. One pos- ury that plants of this order"

(the pea family) "can assimilate nitrogen direct from the atmosphere, and are hence less dependent on the soil for nourishment, and another explanation may be deduced from the fact that its roots penetrate deeper than those of most other crops, and can hence draw supplies from a larger body of soil." At the same time the practical experiments performed at the Saidapet farm, Madras, tend to prove that the plant would not produce so much fibre on rich as on poor soil. Speaking of these experiments Mr Benson says "The seed germinated well, and the plants grew with great luxuriance, but when they had reached the time for cutting was a unfav

performed, the seed being sown on "a light and very sandy loam recently levelled." The land was manured with "12 loads or about 4 tons per acre" of horse manure and the results were most favourable. In the *Mysore Gazetteer* it is stated that the best soil for *sanabu* is the red or black used for *ragi* cultivation. Wisset remarks that clay soils are injurious, but that on a rich soil the fibre is c dry high situations. On the ot the cultivation in the Northern C is sown towards the end of the ra strong clayey soil suits it best.

Rotation
2111

Effects of Sunn Cultivation and the Rotation of Crops Pursued—It is all but universally believed by the Indian cultivators that *sunn*, like *grim* (see Cicer, C No 1067), improves the soil. In the *Bombay Gazetteer* (Kolhapur District, p 172) it is stated "As it is supposed to refresh the exhausted soil, it is considered a good *bevad* or preparatory crop, and is grown as such every second or third year in some of the fields required for sugar-cane, tobacco, and other rich crops. Sometimes it is sown as a second crop and ploughed in when young as a green manure." From Poona it is reported that the leaves are considered "excellent manure." In gardens and occasionally in dry crop lands it is grown solely for manure, the plants being ploughed into the soil when ready to flower." The Director of Agriculture in Bengal states "It is considered by the people of the Lower Provinces to be a renovating crop, and is sometimes used as a green manure to enrich poor paddy land and land that has been infested with weeds." He adds "It comes after one of the pulses or mustard, and is followed by a pulse, sometimes by *shara* onions. When *sunn* is grown on good soil, it is sometimes followed by potatoes. It is not necessary to prepare the land well for *sunn*. Three or four

C 2111

Cultivation of Sunn-Hemp.

CROTALARIA
juncea.

ploughings are sufficient" "Sometimes also paddy and *sunn* seeds are sown together in the same field. When the plants have properly grown, the field is lightly ploughed and the ladder (a kind of harrow) is passed over it. The paddy plants mostly recover themselves, but the tender and juicy *sunn* is buried underground and dies. A few *sunn* plants remain on the surface and are used for food."

CULTIVATION
OF
FIBRE.

Messrs Duthie and Fuller say of the North-West Provinces "Ploughing in a green crop of hemp is known to add considerably to the fertility of the surface soil by increasing its stock of nitrogen, and it is extraordinary that this is not a general practice with native cultivators" In Bombay *tag* (*sunn*) is not considered a good green manure for wheat.

Tillage, Sowing, and Harvesting—As indicated above, the opinion prevails all over India that high cultivation is not necessary for sunn-hemp. Of Kolaba (*Bomb Gas*, XI, 97) it is said "The soil is roughly ploughed twice and the seed sown broadcast" In Bengal "the seed broadcast it is necessary become bushy and coarse"

TREATMENT.
2112
Bombay.
Bengal.

N-W Provinces

Madras.

o plots and watered twice
ombatore, by Nicholson,
allowed no manure, and

the seed is sown broadcast on the ground, without any previous cultivation, at the season when the rains become what the natives call male,—that is to say, when they become heavy. After being sown the field

Mysore.

SEED per
acre.
2113

eighty to a hundred pounds weight to the acre were used in h e t m a t e

CROTALARIA
juncea.

Production and Cost of Sunn-Hemp.

CULTIVATION
OF
FIBRE.
Left standing
for a month
Steeped at
once.

these are supposed to injure the colour of the fibre it allowed to rot in the water of the tank. With regard to *sunn* hemp, the general rule may be almost safely laid down that in moist regions, like Bengal, rapid submer- sion is preferred, and in dry regions, like Madras, stacking the crop is preferred. Roxburgh, from actual experiments arrived at the opinion

hand, states that the strongest opinions have been expressed in favour of first drying the plants before retting, the probability being, as indicated above, that both theories are correct, but applicable to different climatic

Fibre not
removed
from bark till
required.

PRODUCE
2115

THE PRODUCE PER ACRE.—Is so variously stated that it is feared little reliance can be put on the figures. Wisset says that it varies from 3 cwt.

640 lbs per
acre.

In the Kolhapur District the average acre outturn of experiments made at the flower, cut level with the ground, on the 4th December 1887, yielded in the roots on the same day 325lb, on the 11th 487 5lb; and on the 24th 5lb. The average given by Wisset is thus most likely to be a high one and the Kolhapur returns incorrect. Duthie and Fuller say of the North-West Provinces. "The average outturn is about 8 maunds (or 640lb) of clean fibre to an acre, worth about Rs20."

COST
2116

Duthie includes in the cost as "The net re-

that in 1877 its price was as high as 6 seers (12lb) per rupee, whilst a few years ago it was only 1 Re 1 maund.

Area under Sunn-Hemp

CROTALARIA
juncea.

acre The produce was sold by the cultivators to the Telinga Chitties or manufacturers by the thousand handfuls of the dried stems, tall plants fetched two rupees per thousand handfuls, and short plants a rupee and a half. But another crop, he says, was sown in January. This crop had to be watered and more labour spent upon it, but the produce was more valuable. An acre, he says, required $4\frac{1}{2}$ bushels of seed, and its produce was sold for about £1 2s 10½d.

AREA UNDER SUNN-HEMP.—As may be inferred from what has been stated regarding the ambiguity in the Indian literature of this subject, it is next to impossible to discover the extent of sunn-hemp cultivation. Messrs. Duthie and Fuller, from special returns furnished for their *Field and Garden Crops*, state that in the North-West Provinces there are about 40,000 acres under the crop. But in the *Land Administration Report for 1885-86* (page 163 A) it is stated that there were 198,728 acres under "Sana: or Til (sic)." But it is further remarked that the total area under "fibres other than cotton and jute" was in that year only 123,403 acres. This last return would include hemp (proper) *sana:* and *Hibiscus cannabinus*. The Settlement Reports of Oudh show about 800 acres under *sana:*. In *Spon's Encyclopædia* it is stated that there are 50,000 acres in the Panjâb. It is not known from what source that statement was derived, but it seems highly true.

than in the North
about 40,000 acre
hemp plant, how
as sunn hemp, it
acres of brown hemp (*Crotalaria juncea*) grown in Bombay. Full particulars regarding Madras cannot be obtained, but of the districts for which returns are available there were last year 775 acres under "sunn" and 83 acres under "Bombay hemp." What this Bombay hemp may be cannot be learned, but in most works on the subject Bombay hemp is a synonym for sunn-hemp. In 1834-85 there were 380 acres of "Bombay hemp."

duced, but it is not known to what extent the plant is cultivated. In the Central Provinces there were 24,800 acres under "False or San hemp" and in Mysore under "hem." The former explains the name *Crotalaria juncea*. In Burma and Assam there are about 500 acres, in each province, of land entered as under "fibres other than cotton and jute." No returns are available for Bengal, but from personal observation the writer would be disposed to think there must be as much in the Lower, as in the North-West Provinces.

It will thus be seen that the actual area under sunn-hemp cannot be absolutely determined, since the fibre is not included among the agricultural products regarding which regular annual statistics are furnished. But it seems probable that there are at least 150,000 acres annually under the crop in India as a whole.

SEPARATION OF THE FIBRE

The question as to whether the plant should or should not be dried before being placed in the retting tanks having been discussed above, there remains to be given here a brief account of the various modes of retting or of peeling the fibre and of cleaning and boiling it after it has been separated from the stems. In some localities the stems are recom-

SEPARATION
OF
FIBRE.AREA.
2117
N. W. P.
40 000 acres.Panjab
50 000 acres.Bombay
28 614 acres.Madras,
Travancore.

C. P.

Burma
500 acres.
Bengal,India
150 000 acres.
SEPARATION
2118

CROTALARIA
juncea

Methods of separating sunn-Hemp Fibre

SEPARATION
OF
FIBRE.

mended to be buried in the mud at the margin of the tanks, in others, to be submerged in the water by being weighted. In others stagnant water is condemned as destroying the colour and lustre of the fibre, running streams being urged as preferable (*Gibson's account of the Bombay fibre*). But practical and comparative experiments not having been performed in the other provinces similar to those made at the beginning of the present century by Roxburgh, in Bengal a definite opinion for or against the different methods pursued cannot be offered. After removal from the ground, the stems are tied in bundles (20 to 100 in each), but the leaves are generally stripped off and left on the field. When the stems are left until quite dry, the leaves either fall off naturally or are removed by the stems being beaten. It is a common practice to place the bundles of stems erect in 2 or 3 inches of water for 24 hours, so as to give the thicker and lower ends a longer submersion. But the length of time required for retting depends largely on the temperature

Leaves
strippedLength of
submersionStems placed
erect in
water then
horizontal.

laid down lengthways in the water and are kept submerged by being weighted with earth. The time required for retting varies from three days

mentation, while it whitens the fibre, injures its strength. Roxburgh,

"All that seems necessary is to caution the plant, which they are apt to do the bark from the stalks easier, but clear water, well exposed to the sun's

beams, seem best suited for steeping in, because heat hastens maceration, consequently preserves the strength of the fibres, while the clean water

Deep water
Running
water

Damp Mud

the margins of tanks, referred to by some, is even more objectionable, as it seems impossible to adopt this mode of retting without serious loss to the colour of the fibre.

Cleaning of
Retted Fibre

Having discovered that the cultivator, of the stems in his gives way and the

being partially washed, to dry for some hours. This practice, while it is

fibre has been separated and approximately cleaned. In Salsette Island and other parts of Bombay, little or no retting is

Cleaning Sunn-Hemp Fibre.

CROTALARIA
juncea.

employed "The plant while moist is peeled by the hand, and immediately dried in the open air or under cover, according to the state of the weather. By peeling, the fibres are better kept in their natural state of arrangement, and give support and strength to each other, whereas, by the process of the Bengalese, they get so materially entangled that a great loss is always sustained. If they are restored to their natural situation by the heckle, there is a loss of nearly one half of the original quantity, which renders the heckled *sunn* of Bengal of a high price" (Royle). The writer cannot discover any recent description of this Bombay process of separating the fibre without retting, but, as Roxburgh stated, the superior quality of Bombay over Bengal *sunn* hemp seems likely to be due to the fact that the fibre has not been subjected to strong fermentation.

Washing the fibre is very tedious, and a man rarely works for more than three hours at a time but is relieved by turns, he will clean 15 seers a day, which represents the fibre obtained from 5 or 6 maunds of stems. Of Khandesh it is said a man earns Rs for cleaning 40lb of fibre.

Reference has incidentally been made to the period when the crop should be cut, and before proceeding to discuss the further treatment of the fibre it may be as well to add here that the period of cutting will depend on the state of the weather. A softer and more pliant fibre is obtained just as the flowers appear. A few plants are always left by the cultivators to mature seed for the next year's crop, and from the stems of these they extract a strong, though coarse, fibre. On the other hand, it seems to be the habit of some cultivators (the Wunjaras of Bombay) to allow the whole crop to ripen its seeds, this coarse fibre being all they desire, together with the seeds, which are valued as a food for buffaloes. Old stems are also used for making ropes.

required for textile purposes, eg., ropes and twine—it is while hanging over the water that it receives all the treatment grows as "breaking" cleaning is never used fibre that the Native generally separates from the stem.

quotes a report of a sample of *sunn* hemp experimented with at Hull of which it was stated that "by using more care in the steeping and exposure, it will be fully equal to the Baltic." Such opinions are current in the reports of this fibre which appeared while the error existed of supposing it to be Indian grown hemp or *Cannabis sativa*. It is impossible to avoid the impression that *sunn* hemp fell into disfavour when this error was exploded. An expert in 1842, for example, says "Your hemp is very clean—a material point—but it wants more beating and dressing, and I think the natives have not proper implements to do it with. You cannot improve in your mode of packing, it is decidedly superior to the Baltic. I do not despair of seeing the produce of the Baltic supplanted by that of India, as that defect appears to me solely to arise in the management of it—it stands too long before it is pulled or cut, or is too much steeped or exposed, to get the fibre to separate from the stalk." Unfortunately the advances of scientific exploration told all such writers that the defects they complained of were due to the fact that Bombay hemp was not hemp at all, and instead of the fibre supplanting

SEPARATION
OF
FIBRE.
Not Retted.Wages for
cleaning.Period of
cutting.

Soft fibre.

Strong Coarse
fibreSeed used as
buffaloe foodFURTHER
CLEANING.
2119Breaking.
ScutchingSaid to be
nearly as
good as Baltic
Hemp.

CROTALARIA
juncea.

Properties of Sunn-Hemp

PROPERTY OF
FIBRE.
2120

the Baltic hemp it is to-day in the same position commercially as it was a hundred years ago. While not hemp, it is a hemp substitute that deserves a better position than it has as yet obtained.

PROPERTY AND STRENGTH OF SUNN HEMP

£35 a ton

of the fibre by growing and manufacturing it carefully, and Royle mentions a sample of heckled fibre sent to London by the Company that

EARLY
RECORDS.
2121First
Exported.

1791-92. Although numerous favourable reports appeared shortly after this date, the whole interest in the fibre gradually died out, and the

No	Names of the Plants.	Average weight each line broke with when dry	Average weight each line broke with when wet	Average weight gained by wet- ting the lines
4	Sunn (<i>Crotalaria juncea</i>) cut before the plants were in blossom and steeped immediately	112	158	46
5	The same as No 4, but dried, or rather kept some time before they were steeped	60	78	30
6	Sunn cut when in full blossom, and steeped immediately	130	185	42
7	No 6 kept drying for some time	100	166	66
8	Sunn, winter crop cut when the seeds were ripe and steeped immediately	150	201	35
9	The same as No 8, but dried	110	163	43
10	Sunn, winter crop cut when the seeds were ripe, and steeped immediately	160	209	31

Properties of Sunn-Hemp

CROTALARIA
juncea

No	Names of the Plants	PROPERTY OF THE FIBRE.		
		Average weight each line broke with when dry.	Average weight each line broke with when wet	Average weight gained by wetting the lines
1	W. ... Co's	158	190	20
2	"	248	343	38
29	Böhmia nivea) alloté,	240	278	16

SHORTS.

but the new trade is from Bombay, not Bengal

Roxburgh tried the properties of sunn hemp in another way in order

Roxburgh's

NAMES OF THE PLANTS	AVERAGE WEIGHT AT WHICH EACH SORT OF LINE BROKE					
	When fresh			After 120 days maceration		
	White	Tanned	Tarred	White	Tanned	Tarred
English hemp, a piece of new tiller rope	105			Rotten, as was also the English log line		
Hemp from the Company's farm near Calcutta	74	139	45	All rotten		
Sunn hemp of the Bengalese	68	69	60	rotten	51	65
Jute (<i>Bungha pát</i>)	68	69	61	40	42	60

CROTOLARIA
juncea.

Properties of Sunn-Hemp

PROPERTY
OF THE
FIBRE.Deterioration
with age.Removal of
Export Duty

According to these experiments *sunn* hemp stood the action of the maceration better than did either of the samples of true hemp. It has further been shown that a cord 8 inches in size of best Petersburg hemp broke with 14 tons, 8 cwt, 1 qr., while a similar rope of *sunn* only gave way with 15 tons, 7 cwt, 1 qr. Dr. Wight found that a rope of coir of a certain thickness broke with a weight of 224lb, of cotton with 346lb, of American alooe with 362lb, of *sunn* hemp with 407lb, of *Calatropis gigantea* with 552lb, and one of Ambári (*Hibiscus cannabicus*) with 290lb. Royle has shown the slight deterioration which *sunn* hemp undergoes in the following statement: "A rope made in 1803 broke with a weight of 6 tons 2 cwt 3 qrs. whereas a rope made in 1804 broke with a weight of 7 tons 2 cwt 3 qrs."

cent century the bulk of the exports of raw hemp (? *sunn* hemp) went from Bombay and not from Bengal, in spite of the efforts made a few years before that date to create a Bengal trade. This would seem to point to a superiority possessed by the Bombay as compared with the Bengal *sunn* hemp. It seems probable that had this fact been realised by the East India Company, their efforts to establish an Indian hemp industry would have been more successful than was the case with their attempts in Bengal.

In a Report on the Indian Fibres by Cross, Bevan, King, and Watt, recently published by E and F. Spon, the following passage occurs: "It is impossible to urge too strongly the claims of this much-neglected fibre—a fibre which seems to have suffered severely through the immense

that so little of the better qualities of *sunn*-hemp were procurable. Mr. Goffyer and several other Brokers and Merchants stated that their only

Future Pros-
pects.

Chemical Properties of Sunn

CROTALARIA
juncea

actual experiment not to be the case, then there must be something in the climate or soil of Madras and of Bombay more favourable to *sunn* hemp than exists in Bengal

FIBRE

CHEMICAL AND MICROSCOPIC PECULIARITIES OF SUNN

2123

soda, it loses 8.3 per cent, and after an hour only 11.7 per cent. Among Indian fibres it occupies the third or fourth place in point of amount of cellulose. According to this classification Girardinia or Nilgiri nettle heads the list with 89.6 per cent, then Marsdenia with 88.3 and after that *Crotalaria juncea* and *Sida rhombifolia* equal, each with 80.0 per cent of cellulose. "The percentage yield of cellulose of the raw fibre is the most important criterion of its composition and value." It may be worth

Percentage
of cellulose

Europe, there still remains the practical fact that, under the crude methods adopted in India, they are valued as strong and durable fibres. It will be received with no small surprise by many that so humble a position should be assigned to the famed Poya fibre of Assam, and thus in concluding these remarks a possible explanation may be sought in the mode of hydrolysis (or washing and bleaching) employed. The Poya was found to lose 62.7 per cent by being boiled in caustic soda the residue being the cellulose upon which the low opinion of its properties is based. May it not

retains all its properties and under nitration attains a great weight (150.5) being in this respect third in the list of the Indian fibres experimented with by Messrs Cross and Bevan. A writer in *Spon's Encyclopædia* says of *sunn* hemp: "Samples of the fibre, exposed for two hours to steam

CROTOLARIA
juncea.

Trade in Sunn-Hemp

CHEMISTRY
of the
FIBRE

at 2 atmospheres, boiled in water for 3 hours, and again steamed for
inst fix, 3'50; Manila
(without the aid of an
in point of durability
under moisture and under caustic alkali (processes of washing and bleach-

MICROSCOPIC
FORM,
2124Re-examina-
tion desira-
ble,

mean, 0'0015 in. These measurements are in round numbers double

by the process of drying before retting.

TRADE.
2125

TRADE IN SUNN HEMP.

Little or nothing can be learned of a definite nature regarding the
extent of the trade in this fibre. It is grown in every province, and
nearly universally used by the people of India; but, as already stated,
definite information in the use of the
another, and true
same reason we are

Exports
2126

* Presumably sunn hemp or sunn hemp along with a certain amount of the fibre of
Hibiscus cannabinus—*anpal* or *ambadi*.

Imports and Uses of Sunn-Hemp.

CROTOLARIA
juncea

FIBRE.
TRADE IN.

factured Hempen Goods other than cordage. This continued to expand until, in 1870-71, when it was valued at Rs. 64,433, of which Bengal had assigned to it Rs. 53,330. The bulk of these exports went to the Straits Settlements, Ceylon, and Mauritius. From 1871-72, this trade began, however, to steadily decline, and in 1874-75 was valued at

Hempen
Goods.
2127

Ropes and
Cordage
2128

the bulk of the raw fibre so reported may be the Manila hemp used up in the Indian rope factories, and of the hempen goods used in the fabrics of true hemp. 7641 cwt. of hemp

Imports.
2129

USES OF
2131

Canvas.
2132

CROTOLARIA
juncea.

Trade in Sunn-Hemp

CHEMISTRY
of the
FIBRE

at 2 atmospheres, boiled in water for 3 hours, and again steamed for
 1 hour, and then dried at 100° C. for 24 hours. Manilla

MICROSCOPIC
FORM
2124

Mr. King, who worked out the microscope form of the fibre bundles con-

Re-examina-
tion desira-
ble.

ied
 lian
 He con-
 ell marked
 nm, ends
 in *Spon*
 e with the
 plant may
 tent He
 in, min,
 0 001 in,
 rs double
 ility of the
 aving samples of
 pared It would
 - - - - - made

maturity of seed, b
 by the process of drying before retting.

TRADE IN SUNN HEMP

TRADE.
2125

Little or nothing can be learned of a definite nature regarding the
 extent of the trade in this fibre. It is grown in every province, and
 nearly universally used by the people of India, but, as already stated,
 the confusion which exists
 ase, Hibiscus in
 ed to) For this
 foreign trade in
 ports to foreign
 whole of

Exports
2126

* Presumably sunn hemp or sunn hemp along with a certain amount of the blue
 Hibiscus cannabinus—sarpal or ambadi

Imports and Uses of Sunn-Hemp.

CROTOLARIA
junceaFIBRE,
TRADE IN.Hempen
Goods.
2127

Straits Settlements, Ceylon, and Mauritius. From 1871-72, this trade began, however, to steadily decline, and in 1874-75 was valued at

probable that this native industry may have been ruined by the remark-

Ropes and
Cordage
2128Imports
2129

in the Indian rope factories
fabrics of true hemp
7641 cwt of hemp

hemp from all Indian ports to other Indian ports, and these are returned as valued at Rs 24,303, the trade having steadily increased since 1882-83, when it was valued at Rs 6,087.

USES TO WHICH SUNN HEMP IS PUT.—The chief purpose for which this fibre is utilised at the present day is the manufacture of a coarse cloth (*tatputi*) or *charkas* used chiefly for sacking. A large amount of the fibre

USES OF
2131Canvas,
2132

CROTALARIA
juncea

Uses of Sunn-Hemp.

FIBRE
Paper
2I33

paper is regularly made of this material, and large quantities are annually used up by the Indian paper-mills. The paper made by the natives of Bombay is principally of *sunn*. In S. regarding *sunn* paper " "

paper, weighing 39 grs, made from "raw" fibre, was 64lb, as compared with Bank of England note pulp, 47lb. One batch was reported to make a nice, clean, smooth paper, of good colour, but not taking ink well."

For European purposes the fibre may be used as a substitute for hemp or for flax. Speaking of the special form of the fibre produced in Travancore, Dr Royle says "The appearance of this fibre is totally different from any other which comes from India, as it is in the state as if prepared for spinning into thread, and must have been combed or heckled. The fibres are brownish in colour, about 3 to 4 feet in length, clean and shining, not so fine as flax, but still resembling some of the coarser kinds. A very competent judge informed the author that it might be sold for the purposes of flax, or as a kind of flax, and was worth £35 a ton, so some specimens sent to Dundee were valued at the same sum, and it was said could be used for the same purposes as flax, though rather too dry." So, again, "This hemp, when prepared with the patent liquid, became soft, white, and so fine when heckled as to bear the closest comparison with flax at £80 per ton. It is better than any Russian flax for fine spinning. Bombay hemp, rough and dark, and valued at £20 per ton. This article, being similarly prepared, was considered equal in value with the Madras hemp."

Sunn stalks (after removal of the fibre) are used chiefly as fire-wood. But of the Kolaba district, Bombay, it is stated "Hemp torches" stalks with hes round, into about

STALKS
2I36

Torches

2I37

Matches

2I38

C. P. Fibre,

2I39

Bengal

2I40

Bombay

2I41

Madras

2I42

N. W. P.

2I43

Immature

Process

2I44

Deferred

Process

2I45

as well to e obtained tter being the fibre most probably of the form known as *Crotalaria tenuifolia*) as superior to the ordinary *sunn* hemp. We possess so little definite knowledge regarding the cultivated forms of the *sunn* plant that it can only be

mination Were such specimens to be accompanied with samples of the

s of ret- s where ble to courage the natives to adopt the process of preparation of the fibre which was

s of *Crotalaria* ighly probable

Sunn-Hemp-yielding Plants

CROTALARIA
retusa

Food and Fodder—It has already been incidentally remarked that in some parts of India the seeds of this plant are collected and given to cattle. Roxburgh says "This plant—and it is the only one—is also cultivated by the natives of some parts of the Northern Circars to feed their milch-cows with during the dry season. I have found that it is

FODDER

Seeds
2146

MEDICINE
Seeds
2147

Crotalaria laburnifolia, Linn, Fl. Br. Ind., II, 84

2148

A shrubby plant met with in the Western Peninsula, particularly in the South Concan. Properties similar to those of the next species. It is known in Hindustani as *muna*, the *pedda-galligista* of Telegu. Sir Walter Elliot gives it the further Telegu name of *Chiri giligichcha*, and the plant is often seen in gardens on account of its flowering throughout the year.

C. Leschenaultii, DC, Fl. Br. Ind., II, 76

2149

ns of the
ant used
zell and

Satara
Paper
2150

C. medicaginea, Lank, Fl. Br. Ind., II, 81

2151

Vern—*Gulabi*, Pb

A diffuse perennial abundant in the tropical regions of India from Kashmir to Burma, ascending to 6000 feet in altitude.

Medicine—This plant is officinal in the Panjáb being sold in the bazárs under the name of *gulabi* (*Biden Powell, Pb. Pr., 343*).

MEDICINE
2152

C. prostrata, Roxb, Fl. Br. Ind., II, 67

2153

A slender creeping weed, common on the drier plains of India ascending to 6,000 feet.

This is known to be used by them it is used known in Bengal as Roxburgh says this

2154

C. retusa, Linn, Fl. Br. Ind., II, 75

2155

form of sunn hemp
may be passed of

FIBRE
2156

instructive to possess definite information as to the comparative value and property of this fibre with the true sunn-hemp. In Bengal it is

C. 2156

Sunn Hemp yielding Plants

Crotalaria sericea, Retz , Fl Br Ind , II, 75.

although its note is sometimes prepared Kurz alludes to it (p. 111).
Bengal and Roxburgh
but Dr Uday Chandra
the Bengali name of

C. striata, DC., *Fl Br Ind*, II, 84

A low growing shrub, with robust, sulcate, thinly silky branches and large yellow flowers striped with red. Fairly abundant throughout the warmer parts of India.

The Rev. A. Campbell states that this is cultivated by the Santals in Chutia Nagpur on account mainly of its fibre. The plant is known to them as *Son jhunka* and to the Hindustani speaking people of that region as *Son, San*. He adds that the root or a small portion of the stem is tied to the wrists and neck of a person suffering from dropsy. Roxburgh remarks this is known to the Telegu speaking people of Madras as *Mung*.

C. tenuifolia, Roxb., *Fl Ind*, *Fd C B C*, 546

This has been reduced by most botanists to a synonym for *C. juncea*, Linn., which see

C tetragona, *Roxb*, *Fl. Br. Ind*, II, 78

A stiff very handsome shrub, often 68 feet in height, met with on the lower Himalaya (up to 3 500 feet in altitude) from Kumāon to Assam and Pegu. Kurz alludes to this plant and gives it the Burmese name of *Chu Yain*. The shrub flowers in October and November. Mr Gamble in his *List of the Trees and Shrubs &c of the Darjeeling District*, says it is known by the Paharia names of *Kengni*, *kotulkasub* and to the Lepchas as *Suhutung rung*.

C. verrucosa, Linn, *Fl Br Ind II*, 77, Wight, *loc. cit* 200

Vern —
killup
Ainsl
spec e
according to Trimen

according to Frimen
 11 feet A 1 lb 1 lb 1 lb 1 lb plant 2 2 feet in height
 , , , , ,
 , , , , ,

Medicine — Ainslie says "I have given this a place here, on the
 but not growing
 externally).

C 2164

The Croton

CROTON
Joufra.

CROTON, Linn, Gen Pl, III, 293

2165

The generic name *Króton* (a tick) was given by Linnæus to this assemblage of plants in allusion to the shape of the seed. The chief medicinal

Croton argyratus, Bl, Fl Br Ind, V., 383, EUPHORBIACEÆ

2166

Syn — C BICOLOR, Roxb

Vern — *Chonoo*, BURM, *Talib da*, AND

References — Roxb, Fl Ind, Ed C B C, 687 Gamble, Man Timb, 359 Kurz, For Fl Burm, II, 372

Habitat — A moderate sized or small evergreen tree of Martaban, Tenasserim, and the Andaman Islands

Structure of the Wood — Hard, yellow, close and even-grained, seasons well. It is worthy of notice and weighs 46 to 48 lb per cubic foot

TIMBER.

2167

C. aromaticus, Linn, Fl Br Ind, V, 388

2168

Syn — C LACCIFERUS, Linn, ALEURITES LACCIFERA, Willd

Vern — *Welkeppitiya*, SING, *Vid pund*, TAM (names used in Ceylon for C. aromaticus, the form C. laccifera being *Keppitiya* in SING)

References — Beddome, Forester's Man, 204, Wight, Ic, t 19 15, Lisbon, U Pl Bomb, 121, Trimen, Cat Ceylon Pl 81; Gamble, Man Timb, 358, O Shangknessy, Beng Disp, 553

Habitat — An aromatic shrub or small tree, met with in the Dekhan from the Concan southward

Medicine — Said to be used medicinally. Thwaites remarks that the lac obtained from C. lacciferus "is employed by the Singalese for medicinal purposes."

MEDICINE

2169

Lac

2170

2171

C. caudatus, Geisel, Fl Br Ind, V, 388

Syn — C DRUPACEUS, Roxb

Vern — *Nan bhanur* BENG, *Takehabrik*, LEPCHA, *Husta URIYA*

References — Roxb, Fl Ind, Ed C B C 688 Voigt, Hort Sub Cal, 156, Kurz, For Fl Burm II, 375 Gamble, Man Timb 358-359 and XXII

Habitat. — A large straggling, more or less scandent shrub of Bengal Assam, Burma, and South India, found chiefly on the banks of streams. Roxburgh states that it is a native in the country about Dacca, and flowers in March, the seeds ripening in September

Medicine — Mr Home Conservator of Forests, writes, the leaves are applied as a poultice to sprains

Structure of the Wood — White or yellowish-white, hard, close grained. Home says it is used for fuel

MEDICINE.

Leaves

2172

TIMBER.

2173

2174

C. Eluteria, Bennett, affords Cascarilla Bark, — an imported drug

C. Joufra, Roxb, Fl Br Ind, V, 387

2175

Vern — According to Roxburgh *Joufra* is in Sylhet the name of this small tree or shrub

C. 2175

The Purging Croton

CROTON
Tiglum.

the properties having been but recently understood. There is no good Hindi nor a Bengali name for the plant. It is not referred to by U O Dutt nor by Ainslie, and while Roxburgh describes it he makes no mention of its medicinal products. On the other hand, there is nothing to justify a

by European writers

Structure of the Wood.—Whitish to yellow, close-grained, moderately hard and heavy, liable to crack in seasoning.

Domestic Uses.—The plant is frequently employed for fences

TIMBER.

2186

DOMESTIC.

2187

2188

Croton polyandrus, Roxb, see under *Baliospermum montanum*, Muell, Vol I, B 28

Hooker, in the *Flora of British India*, V, 461, reduces this to *B axillare*, Blume. Consult also O'Shaughnessy's *Bengal Dispens*, 555. U C Dutt's *Mat Med of the Hindus*, 229, and Dymock's *Materia Medica, West Ind*, 2nd Ed, 688, the last work has appeared since the issue of the 1st volume of this publication

C. reticulatus, Heyne, *Fl Br Ind*, V, 386

2189

Syn — *C. HYPOLEUCUS*, Dals, *C. ZEYLANICUS*, Muell -Arg

Vern — *Pándhari* or *pandharisala*, MAR

References — Dymock, *Mat Med West Ind*, 2nd Ed 684, S Arjun, *Bomb Drugs*, 122 Thwaites, *En Ceyl Pl*, 276, Dals and Gibs, *Bomb Fl*, 231, Lisboa, *U Pl Bomb*, 121

Habitat — A shrub with slender branches, met with in the Dekhan Peninsula from the Koncan southwards, distributed to Ceylon

Medicine — Sakharam Arjun says the bark is "used as a bitter and stomachic"

MEDICINE

Bark

2190

2191

C. sebiferum, Linn, and *Sapium sebiferum*, Roxb, are synonyms for *Stillingia sebifera*, the Chinese Tallow Tree. This is now cultivated to some extent in India, and, according to Roxburgh, is known in Bengal as *Momchina*

C. Tiglum, Linn, *Fl Br Ind*, V, 393.

2192

THE PURGING CROTON

Syn — *C. PAVANA* (or *PARANA*) Hamilton

Vern — *Jayapala kanakaphala* (in Ainslie dunt, bija) SANS, *Jaypal*

References. — Roxb, *Fl Ind*, Ed C B C 688 Voigt, *Hort Sub Cal*,

The Purging Croton.	CROTON Tiglium.
the properties having been but recently understood There is no good	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	
.	

Croton polyandrus, Roxb, see under *Baliospermum montanum*, Muell, Vol I, B 28

Hooker, in the *Flora of British India*, V, 461, reduces this to B

C. reticulatus, Heyne, *Fl Br Ind*, V, 386

Syn — *C. HYPOLEUCUS*, Dals, *C ZEYLANICUS*, Muell - Arg

Vern — *Pandhari* or *pandharisala* MAR

References — *Dymock Mat Med West Ind*, 2nd Ed, 684, S Arjun, *Bomb Drugs*, 122 *Thwaites, En Ceyl Pl*, 276, Dals and Gibs, *Bomb Fl*, 231, *Lasboa*, U Pl *Bomb*, 121

branches, met with in the Dekhan Pen distributed to Ceylon

ys the bark is "used as a bitter and

stomachic

C. sebiferum, Linn, and *Sapium sebiferum*, Roxb, are synonyms for *Stillingia sebifera*, the Chinese Tallow Tree This is now cultivated to some extent in India, and, according to Roxburgh, is known in Bengal as *Momchisa*

C. Tiglium, Linn, *Fl Br Ind*, V, 393

THE PURGING CROTON

Syn — *C. PAVANA* (or *PARANA*), Hamilton

Vern — *Jayap la kanakaphala* (in Ainslie *duntis biya*) SANS, *Jaypál*

by Ainslie

datun ARAB, *Bed anjire khatai*, *habbe khatai*, PERS (according to Moodeen Sheriff)

Referred to in the *Fl Br Ind* CROTON for *Lat Hort Sub Cal*

CROTON
Tiglium.

The Purging Croton

and Arboretum, 67, Simmonds, Trop Agri, 424

Wah tat. A small tree (up to 30 feet high) with the odor of lemon

OIL
Nuts,
2193

Bombay
2194
Cochin
2195
Chinese,
2196
European
Expressed
2197

2198

MEDICINE
Seeds,
2199
oil
2200

2201

nuts are exported chiefly from Bombay and Cochin (often being also Chinese re exports), and the oil is expressed in England. Dr Dymock informs the writer that the oil is expressed at the Government Medical Store Depôt at Bombay. It costs about 12 annas a lb, whereas in 1825, the same oil was sold for about 10 shillings an ounce in England. The plant used to be grown for the purpose of its seeds at Hewra, but the supply is now imported from China *via* Singapore. The nuts sell for Rs1 per maund of 41lb.

It is necessary to be cautious in handling the nuts or the oil, owing to their blistering the skin. The oil is frequently used for colds in the chest as an external application, causing a severe blister. It is much resorted to as a domestic cure but is not recommended by the profession.

§ "The drastic principle of the oil has not yet been isolated, it appears to exist not only in the seeds but also in the leaves and wood" (*Professor Warden, Calcutta*).

Medicine.—The SEEDS are used as a powerful drastic purgative, and the OIL is regarded as a valuable medicine. In overdoses they act as an acro narcotic poison. When externally applied the oil is a stimulant rubefacient and counter-irritant. Croton oil is said to possess powerful hydragogue cathartic properties. It is also useful in dropsy, obstinate

fever, constipation, intestinal worms, enlargements of the viscera, ascites, anasarca, &c."

C. 2201

The Purging Croton

CROTON
Tigilium.MEDICINE.
Grana Tiglia.
2202

opinions of a few Indian medical officers who re-made known the properties of this drug at about the beginning of the present century or the close of the last. Practically all subsequent writers have but slightly altered the

2203

biting the oil at first in larger doses than one or two minims, to adults,

the oil highly useful as an emmenagogue

"Rumphius informs us that the root of the plant is supposed, by the inhabitants of Amboyna, to be a useful drastic purgative, in cases of dropsy, given rasped in doses of a few grains, or as much as can be held between the thumb and finger" "Rheede, who speaks of the plant under the name *caddé avanacu*, says, that the leaves rubbed and soaked in water also are purgative, and when dried and powdered are a good external

Root
2204Leaves
2205

2207

their uses as a drastic purgative the seeds are applied in the form of liniment to the penis in cases of impotence and have a high reputation in this disease amongst the natives" (Lal Mahomed, 1st Class, Hosp'l. Asstt.,

C. 2207

CROTON
Tiglium.

The Purging Croton

Habitat—A small tree (15 to 20 feet high) met with under cultivation throughout the greater part of India, probably indigenous or only naturalised in Eastern Bengal and Assam and southward to Malacca, Burma, and Ceylon.

OIL
Nuts.
2193

Bombay
2194
Cochin
2195
Chinese.
2196
European
Expressed
2197

Nuts are exported chiefly from Bombay and Cochin (often both as Chinese re-exports), and the oil is expressed in England. Dr Dymock informs the writer that the oil is expressed at the Government Medical Store Depot at Bombay. It costs about 12 annas a lb, whereas in 1825, the same oil was sold for about 10 shillings an ounce in England. The nuts are sold in the market of its seeds at Hewra, but in Singapore. The nuts sell

2198

the nuts or the oil, owing to its being frequently used for colds in the chest, it produces a severe blister. It is much resorted to as a domestic cure but is not recommended by the profession. § "The drastic principle of the oil has not yet been isolated, it appears to exist not only in the seeds but also in the leaves and wood" (*Professor Warden, Calcutta*).

MEDICINE
Seeds.
2199
Oil
2200

Medicine.—The SEEDS are used as a powerful drastic purgative, and the OIL is regarded as a valuable medicine. In overdoses they act as an acro narcotic poison. When externally applied the oil is a stimulant rubefacient and counter-irritant. Croton oil is said to possess powerful hydragogue cathartic properties. It is also useful in dropsy, obstinate constipation, and apoplexy. When the nuts are boiled in water (as at the present day) to extract the oil, the nuts boiled (as at the present day) to extract the oil, and according to many authorities, the seed leaves (or seed leaves) are also used.

2201

The Purging Croton

CROTON
Tiglium

MEDICINE
Crana Tigha.
2202

tion, they have been long banished from modern practice For the same

2203

biting the oil at first in larger doses than one or two minims to adults,

the oil highly useful as an emmenagogue

"Rumphius informs us that the root of the plant is supposed, by the inhabitants of Amboyna to be a useful drastic purgative, in cases of dropsy, given rasped in doses of a few grains or as much as can be held who speaks of the plant under s rubbed and soaked in water powdered are a good external

Root
2204

Leaves
2205

tion and drops
tion, amountir
bor is in Benga
Shib Chunder

CROZOPHORA
plicata.

The Indian Turnsole.

MEDICINE.
2208

Mam Dispensary, Hoshangabad, Central Provinces) "The seeds, half roasted over a lamp or candle flame, and the smoke inhaled through the nostrils, relieves a fit of asthma" (*C. I. E., Madras*) "I used oil or olive oil to . . . (Dojil Chunder Shro rubefacient" (*D. Pic.*) is frequently applied (Surgeon-Major Robb, Civil Surgeon, Ahmedabad)

2209 *Croton tinctorium*, *Turnsol*, see *Crozophora* (*Chrozophora*) *tinctoria*, A. Juss.

Crown Bark, see *Cinchona Condaminea*, *Huml.*, RUBIACEÆ C. 1129.

2210

CROZOPHORA, A. Juss.; *Gen Pl*, III., 305

error in the spelling of the name
then arranging the material for
effect of placing it in the wrong
from γρωζω the word should of

course be *Chrozophora* as corrected by Necker.

2211

Crozophora (*Chrozophora*) *plicata*, A. Juss.; *Fl. Br. Ind.*, V., 409, EUPHORBIAEÆ.

Syn.—C. ROTTLERI, A. Juss.; C. Plicatus, Vahl; C. ROTTLERI, Geisel, C. TINCTORIUS, Wall, *Burm*, C. Plicatum, Willd. (*in Roxb., Fl. Ind.*)

Vern.—Shadrei, saba'i sonballi, HIND, SIND and Okharada, GUZ., Khadi-
okra, BENG., Pango nari, SANTALI, Surjavarzi, SANS., Pat kanda,
nilkharis, nil-ak rai, PB.; Neal boti, TANN., Gurugu chettu, lingua
miryam, TEL.

References.—Roxb., *Fl. Ind. Et. C. B. C.*, 657, *Thwailes, En. Ceylon*
Fl., 443, *Dalz. & C.*, *Fl. Ind.*, *Fl. Andrica* 66,
Chutia Nagpur, 18
Fl. Ind., 2nd ed.

Si . . . is the Indian Turnsol—
Wilson, Brown, Pid-
dington, and others have imagined the plant to be the sun flower, and
still further to increase the confusion, they have turned the old Greek name
Chrozophora tinctoria, L. (ἡλιοτροπιον μικρον) into the modern
Heliotrope, and explained the various Indian names of *Croz plicata* by
Heliotropium (*Tiardinum*), *Indicum*, *Lin II*, *Lee King*, p. 281 "This
mistake has been repeated by O'Shaughnessy, who says that *Chro-*
zophora tinctorium, the Turnsol (*Turnsole*) is the ἡλιοτροπιον
μικρον of Dioscorides."

2212

Habitat.—There are two well marked forms of this plant—(1) a small
procumbent annual, found in sandy damp situations, such as on the
banks of . . .
rial bus
the Flor
the warri
Burma,
dit on et
phora ti

C. 2212

The Turnsole

CROZOPHORA
tinctoria.

Madras, and Burma, and is of no interest from an economic point of view, since the properties described below are alone applicable to the erect plant, and to *Chrozophora tinctoria*. The confusion alluded to by Sir Walter Elliot may be accounted for by the fact that the crumpled leaves of the procumbent plant are remarkably Borag. naceous in their

on to
dye
—“It
sules,
becomes blue after exposure to the open air, they, no doubt, contain colouring matter, which might be turned to good account in the arts”
O Shaughnessy, who wrote 20 years later still, says—“The summits of the plant and the fruit are a far the proper portion of the dye named”

on to
dye
—“It
sules,
becomes blue after exposure to the open air, they, no doubt, contain colouring matter, which might be turned to good account in the arts”
O Shaughnessy, who wrote 20 years later still, says—“The summits of the plant and the fruit are a far the proper portion of the dye named”

species

Fibre.—The Santals prepare a strong and useful rope fibre from the bark, but it is difficult to separate (*Campbell*)

Medicine.—The ASHES of the root are given to children in coughs. The LEAVES are considered depurative, and are officinal under the name *milkhanthi*. The SEEDS are used as a purgative. The Revd A Campbell states that the Santals mix the root with that of *Carissa Carandas* for blistering purposes. “This is a plant which Dr F Hamilton (MSS) had brought to him in Behar, as one of those which was supposed to have virtues in leprosy affections, the dry plant is made into decoction, to which is added a little mustard” (*Ainslie*)

Timber.—The stems of both this and the next species are regularly collected as fuel. Dr Stewart says of *C. tinctoria* “It is cut and carried into the city of Lahore to be used as fuel in ovens” This fact may be

to are both perennate annuals” The and to be most prominent with in rice

fields of Bengal, as distinct from the bushy perennial found in Chutia Nagpur and Upper India

Crozophora tinctoria, *A. Juss*, *Fl. Br. Ind.*, V, 408

TURNSOLE Eng

Vern.—*Shalevi*, *sonballi subali* HIND & SIND *Toppal buti*, *nilan kukranda*, *Pa*, *Kap-o-chist* in the Harirod Valley, Afghanistan (*Aitchison*)

Habitat C. — — — — —

FIBRE
2214
MEDICINE.
Ashes
2215
Leaves,
2216
Seeds,
2217
Root
2218
Dry Plant,
2219
TIMBER
Fuel,
2220

2221

CROZOPHORA tinctoria.		The Turnsole.
DYE Blue 2222	less woolly leaves than either <i>C. plicata</i> (procumbent form) or <i>C. tinctoria</i> , but is covered with a granular mealy substance Dye—Although it seems probable that most Indian authors who allude to having observed the fruits of <i>Chrozophora</i> yielding a purplish dye, speak of the erect perennial form of <i>C. plicata</i> , still <i>C. tinctoria</i> doubtless affords the same dye in this country as it is cultivated for in India of the dye principle be of some practical utility, industry in this dye-stuff, opean uses and methods of preparation The researches of Dr Joly (<i>Ann de Chim et de Phys</i> , VI, 111) have shown that the dye principle occurs in all parts of the plant and not in of the	
Yellow. 2223	cles. blue to from 50° to 60°, that liquid assumes a rather deep violet blue colouration, and deposits, on being evaporated, a beautiful azure-blue resinous	
Green 2224 Litmus on Rags 2225	without the aid of mordants, a violet-red upon wool, silk, and cotton tissues, and that this colour may be rendered fast by steaming and the simultaneous : however, turn the colour more blue” 383). “This dye is called Turnes ie plants—little herbs they yield about omes purple by It is chiefly exported to Honanu, and is prepared for exportation by soaking coarse linen rags or sacking with it, the rags being previously washed clean After soaking they are allowed to dry, and are exposed to the influence of ammonia by being suspended over heaps of stable manure They are then packed in sacks and are ready for shipping to Holland” (<i>Treasury of Botany</i>) “The red colour of the outer crusts of some kinds of Dutch cheese is due to the presence of some lactic and butyric acids in that substance No good substitute for this ‘litmus on rags’ for the last named purpose has as yet ever been found A sum of £10,000 is annu- tants of Grand Gillar- would like to be any ing after having been en used to rub cheese t the old rags take up dily than new ones”	
Powder. 2226		
Sacking Impregnated 2227	(Crookes)	
TRADE. 2228	reason to suspect that a very extensive trade might be done in it the plant is wild everywhere on the waste lands of India, luxuriating on both dry sandy tracts and river margins, it might be grown at a small cost anywhere, and the subject thus seems well worthy of attention, as there are many purposes to which it might be put in India The writer	

The Crustacea—Prawns, Shrimps, Crayfish, &c.

CRYPTERONIA
pubescens.

can discover no evidence of its ever having been utilised by the natives of India but it is a remarkable coincidence that in Bengal, at least, it bears a name (*okra*) now given to several introduced American plants

Dr
Bixa
t be
the
ogue

TURNSOLE-
DYE.

2229

grown as hedges around their mango trees, thus affording a possible extra revenue, while serving a purpose for which they are eminently suited, since no herbivorous animal has as yet been observed to browse either on *Jatropha glandulifera* or *Chrozophora tinctoria*.

CRUSTACEA.

2231

FOOD
Crabs
2232
Prawn
2233
Lobsters.
2234
Cray fish.
2235
Shrimps
2236

2237

MEDICINE.
2238

animal food "

CRYPTERONIA, Bl., *Gen Pl*, I, 782

2239

[*Man Timb* 199, LYTHRACEÆ
Crypteronia pubescens, Blume, *Fl Br Ind*, II, 574, Gamble,

2240

Vern — *Anando*, BURM

Hab tat — *A tree 20 feet high*

TIMBER.
2241

for fuel.

C. 2241

CRYPTOMERIA
japonica.

The Cryptomeria

CRYPTOCARYA, *R. Br ; Gen Pl, III, 150*

Several species afford valuable timber.

- 2242 *Cryptocarya amygdalina*, *Nees ; Fl Br Ind, V, 118 ; LAURINEÆ*
 Vern — *Patmaro*, NEPAL ; *Kaledso*, LEPCHA
 Habitat — A tree with spreading branches, found from Nepal eastwards to the Khasia hills and south to the Andaman Islands
 Structure of the Wood.—Strong and useful
- TIMBER.
 2243
 2244 *C. ferrea*, *Bl. ; Fl Br Ind, V, 119*
 [*Lisboa*, *U Pl. Bomb, 115.*
- 2245 *C. Wightiana*, *Thwaites, Fl. Br Ind, V, 120 ; Wight, Ic, t 1829.*
 Vern — *Golu mora*, SING
 Habitat — A tall tree, frequent in the Dekhan peninsula from Kanara southwards to Ceylon
 Structure of the Wood.—Strong and durable, useful for building purposes
- TIMBER.
 2246

CRYPTOLEPIS, *R Br ; Gen Pl, II, 740*

- 2247 *Cryptolepis Buchanani*, *R & S, Fl Br. Ind, IV, 5, Wight,*
 [*Ic, t 494 ; ASCLEPIADEÆ*
uruga pila-tige, a tree it is called Ma sti-like
India from Kashmir ascending the Himala
Vizianagram make
the Santals rest on the "Doctrine of Signatures."
- References — *Roxb, t*
Dils & Gibbs 148
II, 109, Elliot Fl
Chutia Nagpur, 49 ;
II, 113
Fl, 330 ;
Burm,
Pl,
Bomb
- FIBRE
 2248
 MEDICINE.
 2249
- 2250

CRYPTOMERIA, *Don ; Gen Pl, III, 428.*

- 2251 *Cryptomeria japonica*, *Don, CONIFERÆ*
 Habitat — A handsome tree, native of China and Japan, but largely cultivated throughout the districts of Darjeeling, Simla, and occasionally in other hill stations
- C. 2251

Caoutchouc-producing trees.

CRYPTOSTEGIA
grandiflora.

Structure of the Wood—White, soft, with a brown, often almost black, heart-wood; very uniform, with narrow bands of darker and firmer tissue at the edge of each annual ring.

TIMBER.
2252

CRYPTOSTEGIA, R. Br.; Gen. Pl., II, 742

[ASCLEPIADACEÆ

Cryptostegia grandiflora, R. Br.; Fl. Br. Ind., Vol. IV., 6,

2253

Vern.—*Vilarjuti vakundi*, MAR (according to Dr. Sakham Arjun in a letter to the author), *Palay*, MAL (according to Sir George Birdwood).

Habitat—An extensive climber, cultivated in various parts of India, supposed to be a native of Africa or Madagascar.

Caoutchouc.—Dalzell and Gibson (*Bomb. Fl. Sp.*, 55) say "the whole h is like India-rubber, ng made to extend the lombay (*See Agri-Hort. Hyderabad, Sind, 1882*, A sample of the Sind s grown in the Botanic

CAOUTCH-
OUC
2254

much blackened by oxidation; a very small portion only had retained the light colour of Ceara rubber. The whole had become agglomerated by the adhesiveness of the little separate masses of which the sample was composed

"The sample was carefully torn to pieces and examined, a separate examination being made of the lighter and darker portions. The only difference found is in the much larger quantity of moisture met with in the lighter portions

"It might have been possible to have given some assurance on this point if the time was stated how long this sample had been collected. In its present condition it is hardly equal to Ceara rubber from Brazil, although

con-
stantly

the lighter portions lost 15.0 per cent, the darker portions lost only 2.9 per cent. The amount of ash obtained from the lighter portions was before washing 4.3 per cent, after washing 2.7 per cent. The darker portions yielded before washing 4.2 per cent, after

2255

vulcanized, as compared with the darker portions, but in this respect no difference could be perceived"

The Conservator of Forests, Northern Circle, Bombay Presidency, wrote on the 16th January 1885, that *Cryptostegia grandiflora* "is cultivated in gardens in nearly every station in India, and can be easily propagated. The cost of collecting the sap would be so great that a plantation is not

2256

The Cucumis or Melon.

Crystal Rock, see **Carnelian**, C. 616.

CTENOLEPIS, *Hook. f. : Gen. Pl., I, 832.*

2257

Ctenolepis Garcini, Naud., *Fl. Br. Ind*, II, 630; CUCURBITACEÆ.

Vern.—*Gud: murals*, TEL.

References—*Roxb*, *Fl Ind*, Ed C B C, 703; *Dalz & Gibs*, *Bomb Fl*, 99; *Atkinson*, *Econ Prod*, V, p 12

Habitat—An annual climber, met with in Bundelkhand and the Dekhan. Grows on rubbish heaps and hedgerows.

MEDICINE.

2258

Medicine.—Atkinson says the fruit, seeds, and roots are used in medicine.

Cubeba officinalis, *Miq.*, see *Piper Cubeba*, *Linn.*; **PIPERACEÆ**

Cubebs, see **Piper**.

2250

CUCUMIS, *Linn* , *Gen Pl* , I, 826.

A man of almost no knowledge of the subject was asked to give an account of the

HISTORY

2260

History—Much confusion still exists regarding the Indian so-called wild and cultivated species and varieties. Roxburgh was the first author who systematically examined and described the Indian forms. In his *Flora Indica* he gives the distinctive characters of what he regards as nine species, two of which, by all subsequent botanists, have been removed to other genera, and the remaining seven reduced to three species. De Candolle, however (*Orig. Cult. Pl.*, p. 259), seems to be of opinion that they represent but two species—*C. Melo*, Linn (embracing all the wild and cultivated Indian, African, and American forms of the Melon) and *C. sativus*, Linn (the Cucumber). Referring to Roxburgh's nine species, Ainslie says they are all natives of India "except the Melon,"

seems probable that *molam* or *mulam-pandu* is but a modern corruption from the English word *melon*. There are, however, many ancient and

(curbita and
lassification
ross fertilize

C. 2260

The Sweet Melon.

CUCUMIS
Melo.

with the production of fertile seeds, the plants so experimented with may be viewed as varieties or even only cultivated races derived from a com-

HISTORY.

2261

fertile individuals, as we see, for example, in the human species, they must

monly stated that a fertile mule exists between the two species of Camel—*Camelus dromedarius* and *C. bactrianus*—but the progeny is more unmanageable than the mule itself, and is accordingly very little bred (see article on Camel, C 203). But Naudin's physiological classification

2262

India.

[*Mono Phanerog*, III, 462; *CUCURBITACEÆ*.Cucumis Melo, Linn, *Fl Br Ind*, II, 620, *Cogniaux*, in *DC.*,

2263

The SWEET MELON (Stewart and also Baden Powell call this the Musk Melon, but by giving it at the same time the name *Kharbua* they remove the suspicion of *Cucurbita moschata*. The information furnished by these authors under "*C. Melo*, *L*—musk melon" has accordingly been compiled under this species)

Vern — *Kharbua* or *kharbua* *khurbuj* or *kharbua*, HIND, *Kharmuj*, BENG, *Tarbuja*, SANTAL, *Dungra*, C P, *Khurbua*, KANGRA (in Seidl Rept, 25), *Aharabua*, *khurbuj*, *chibuda*, BOMB, *Chibunda*, MAR,

seems probable that in Bombay *Tarbuja* and *kharbua* are applied to distinct forms of the melon

References.—Roxb, *Fl Ind*, *Fd C B C*, 701, *Voigt*, *Hort Sub Cat*, 58, *Thunberg*, *En Ceylon Pl*, 127, *Dale & Gibb*, *Bomb Fl*, 103, *Supp*, 36, *Stewart*, *Pb Pl*, 96; *Aitchison*, *Cat Pb and Sind Pl*, 63; *DC*, *Orig Cult Pl*, 258, *Naudin*, *Ann des Scien Natur*, 4th Series,

CUCUMIS
Melo.

The Sweet Melon.

Trop Agri, 423.

Habitat.—Extensively cultivated on account of its fruit in the sandy basins of rivers. Said to be a native of North-West India, Baluchistan, and west tropical Africa (*DC.*). Ainslie wrote in 1826 that *C. Melo* "has been said to be a native of Calmuc Tartary, an opinion adopted by Willdenow; in India it is cultivated by seed brought from Persia (see *Tavernier*).

is called *ki*

Hindustan

molam pu

which pre

methods of cultivation see under a further paragraph. A good plate of this plant occurs in Duthie and F

OIL
2264

Oil.—The flattened and ellip
fact, the seeds of most of the mem
and gourd family, contain oil, bu

any considerable extent are those of the Sweet-melon (*Cucumis Melo*) and the Water-melon (*Citrullus vulgaris*). From West Africa large quantities of melon seeds are exported to France. China also does a considerable trade in them, but in India the fruit is chiefly eaten as such, and not allowed to ripen its seeds, and accordingly the supply of melon oil is not extensive.

MEDICINE
Seeds,
2265

Medicine.—The seeds are used as a cooling medicine. They are

Mixed seeds.
2266Pulp.
2267

diuretic, very beneficial in chronic, and also in acute eczema.

FOOD
2268

forcing beds. This is the practice in growing melons in the alluvial rivers such as the Ganges and Jumna, which consist wholly of white sand. Where the river deposit is of richer quality and contains a mixture of organic matter, a much less amount of manure is required, and it is

C. 2268

The Sweet Melon.

CUCUMIS
Melo.

FOOD

reported that occasionally manure is altogether dispensed with. The melon beds commence fruiting in April and continue yielding until they are

Madras all stated to be rather watery, but Moorcroft declares the people fatten on them 'as horses are said to do in Bokhara'. Vigne states that the melons of Tibet here he saw grow to 10,000 feet are

2269

it), several varieties of melon are extensively grown, and Davies' Trade Report states that 300 mule-loads are annually imported thence via Dacca. The melons which, It has enerate melon such the

In Manipur the melon is cultivated by the Nagas and is of a spherical form with ten segments. The pulp of the fruit is usually sweetish and pleasant, and is eaten by Europeans as well as natives.

Cultivation.—Firminger refers to two good forms of melons, one of which—the Afghan—has been alluded to above. He says "the kind which ranks as finest of all, called the *surdah*, is a native of Cabul, and has not that I am aware, been cultivated with success in any part of India." "The seeds of this kind are at once to be distinguished from those of any other, being fully four times larger." "The next kind, second perhaps only to the *surdah*, and superior to any other with which I am acquainted, is, I believe, also from Cabul. Like the *surdah*, too, it is of the green flesh sort. It is of a large oval form, with very smooth, pale green exterior, traced here and there with a delicate network. This succeeded most satisfactorily at Ferozepore, and was the one which I cultivated exclusively. The seeds of this also may be known by the largeness of their size." Quoting from the Agri-Horticultural Society's Journal Mr Firminger gives an account of a melon sent from Buxar by a Mr W H Bartlett, who writes "with culture in a manured soil, the smaller of these melons may be grown in a sub-tropical climate."

CULTIVA-
TION
2270

2271

CUCUMIS Melo.	Indian Forms of the Melon.
CULTIVA- TION.	<p>system by which the Cabul melon might be grown It was, however, troublesome and expensive though attended with success. The chief features of this system were the selection of an open situation even by the sun, the soil being a light sandy loam, the seed sown to the depth of 1 inch; a great point being the steeping of the seeds in warm water for 24 hours; afterwards retaining them in wet ashes or a wet cloth until they sprout; as soon as sprouted to be sown about a foot apart and an inch and half deep; lastly, to be deluged with water every day from sowing until the</p> <p>beetle. The usual preventatives adopted by the native gardeners is to dust the plants with wood ashes. This must, however, be highly injurious, and since in most cases with age the plants cease to be attacked by the beetle a better course is to cover the seedling plants with a muslin frame. The following two forms are the cucumber-like plants which, by modern European botanists, are treated as melons, and are not even allowed the position of varieties from the type.</p>
2272	(i) Cucumis Melo, Linn : var. Momordica. This form does not appear to be referred to in the Flora of British India. It is a very common plant in the gardens of the natives, and is cultivated by them for its fruit, which is a large, elongated, somewhat flattened, and is covered with a scabrous or bumpy surface. It is frequently mottled. As Roxburgh says, the plant is more like the cucumber than the melon, except that it is less scabrous and larger. Atkinson remarks: "C. Momordica, Roxb, the phunt
2274	ously when ripe; it is then from a foot to 2 feet long and from 3 to 6 inches in diameter, and weighs 4 to 8 lb. The seeds are smaller than C. 2274

Indian Forms of the Melon.	CUCUMIS Melo.
<p>those of the common melon. A good drawing is given of the plant by Duthie and Fuller in <i>Field and Garden Crops</i></p> <p>Vern.—<i>Phut</i> or <i>phunt</i> (ripe), <i>kachra</i> (when unripe), <i>tuti</i>, HIND ; <i>Phuti</i>, BENG ; <i>Kakari-kai</i>, TAM ; <i>Pedda kai</i>, <i>pedda-dosrai</i>, TEL ; Dr U. C. Dutt says this is the <i>Ertaru</i> of Sanskrit writers Kurz in his Report on Pegu gives <i>Tha khwahumzay</i> as the Burmese</p> <p>Habitat.—Cultivated here and there throughout India : Roxburgh remarks that in the Carnatic it is a cold season crop. According to</p>	
<p>Medicine.—The seeds are used as a cooling medicine.</p> <p>Food.—Roxburgh writes.—“The fruit is much eaten both by Natives</p> <p>very wholesome.</p>	<p>OIL 2275 MEDICINE. 2276 FOOD. 2277</p>
<p>2) <i>Cucumis Melo</i>, Linn. ; var. <i>utilissima</i>.</p> <p>Syn.—<i>C. UTILISSIMUS</i>, Roxb.</p> <p>Vern.—<i>Kakri</i>, <i>kakni</i>, HIND ; <i>Kakkur</i>, or <i>kankur</i> (<i>Kakri</i>, according to</p> <p><i>Takkua</i>, BURM.</p> <p>Re</p>	<p>2278</p>
<p>Description.—The various writers who have described the Indian melons, cucumbers, &c., give somewhat conflicting accounts of this fruit.</p>	<p>DESCRIPTION 2279</p>
<p>some varieties of cucumber. white, usually changing to a</p>	<p>Seeds. 2280 Fruits. 2281</p> <p>C. 2281</p>

CUCUMIS
sativus

The Melon; The Cucumber

and Fuller) In the Gazetteer of the Khandesh District, Bombay, it is stated that the fruit of this plant is one of the most valuable of the district.

CULTIVA-
TION
2282

It is cultivated throughout the district, and is one of the most valuable of the district. It is cultivated throughout the district, and is one of the most valuable of the district. It is cultivated throughout the district, and is one of the most valuable of the district.

OIL.
2283
MEDICINE
2284

It is cultivated throughout the district, and is one of the most valuable of the district. It is cultivated throughout the district, and is one of the most valuable of the district. It is cultivated throughout the district, and is one of the most valuable of the district.

FOOD.
2285

It is cultivated throughout the district, and is one of the most valuable of the district. It is cultivated throughout the district, and is one of the most valuable of the district. It is cultivated throughout the district, and is one of the most valuable of the district.

2286

It is cultivated throughout the district, and is one of the most valuable of the district. It is cultivated throughout the district, and is one of the most valuable of the district. It is cultivated throughout the district, and is one of the most valuable of the district.

2287

Cucumis sativus, Linn.; Fl Br Ind., II, 620.

THE CUCUMBER

The larger form of this fruit, but for the appendant structures on the young stem, often closely resembles *C. Melo*, var. *Momordica*, and also var.

The Cucumber.

CUCUMIS
sativus.

utilissima, more nearly in fact than they approach the melon. Hence a certain confusion in the vernacular names

Syn — C HARDWICKII, *Royle, III, 147*

Vern — *Khira* HIND, *Kaknai*, ORISSA, *Sasa*, *khira*, BENG *Khira*, *Ahiyar*, PB, *Aokri* SIMLA, *Kakri*, *kankri* BOMB, *Kakdi*, MAR, *Aakari* GUZ, *Muhetehti*, TAM, *Dosa kina* TEL, *Sante kayi*, KAN, *Trapusha* (according to Dutt), *Sukasa* (according to Piddington),

Dispens, 32, *S Arjun*, *Bomb Drugs* 58 *Hunter*, *Orissa*, II, 188, *Firminger Man Gard Ind*, 126 *Baden Powell*, *Pb Pr*, 347 *Duthie & Fuller Field and Garden Crops*, 53, *Li boia*, *U Pl Bomb*, 159, *Bird wood*, *Bomb Pr* 283 *Plates* 51 52

2283

antiquity of the species in Europe. There is even an Esthonian name, *Uggurits ukkurits, urits*. It does not seem to be Finnish, but to belong to the same Aryan root as *aggouria*. If the cucumber came into Europe before the Aryans, there would perhaps be some name peculiar to the Basque language, or seeds would have been found in the lake dwellings of Switzerland and Savoy, but this is not the case. The peoples in the neighbourhood of the Caucasus have names quite different to the Greeks; in Tartar *kisar*, in Kalmuck *chaja*, in Armenian *karan*. The name *chisar* exists also in Arabic for a variety of the cucumber. This is, therefore,

THOSE OF CUCUMIS UTILISSIMA

"In sunstroke pieces of cucumber are put on the bed so that the patient may breathe the moistened air in order to neutralize the heat of the sun." (A Surgeon)

UCUMIS
sativus.

The Cucumber.

FOOD,

is much less eaten than one of a dark green, and the other of a creamy-white colour. The rainy season varieties are the most common, and are universally eaten by natives of all classes as well as by Europeans. The other varieties are much less common, and are not so much eaten. The small

and

The rainy season varieties are the most common, and are universally eaten by natives of all classes as well as by Europeans. The other varieties are much less common, and are not so much eaten. The small

2292

and if so the further suggestion might be offered that it may after all prove but a peculiar form of *Cucumis sativus*. Most if not all the forms of

might be tried in addition to the preparation of carefully dried specimens both of the natural and hybridised plants.

CULTIVA-
TION
2293

Cultivation—These plants are alluded to by many writers, but it is scarcely necessary to repeat all their statements. The following abstract from the *Indian Forester* (written by Mr. Gollan, Superintendent, Botanic Gardens, Saharanpur) gives some particulars regarding the cultivation of hot season cucumbers or gherkins:—

"This is a variety of the common cucumber, with small egg-shaped

should be sown along both sides of the drill, and if the soil be dry, water should be given immediately after sowing. After germination, water every ten days, but like the *kakri* this vegetable should not be watered too often." (*Vol. IX, 162*)

Regarding the rainy season forms Mr. Gollan (*Ind. For.*, IX, 201) says they have much larger fruits and are more like the English cucumber; there are two forms,—“when in a young state the colour of one is a dark green, and of the other creamy-white; when full grown, both are about a foot long, and the colour changes to a rusty brown. These two, although not equal to the commonest varieties met with in England, are not to be despised. They thrive with little care and are always sure of yielding a crop.”

Firminger, in his article on Cucumber, deals fully with the two forms of the rainy season plant, but was apparently ignorant of the hot season one or did not view it as a cucumber. Speaking of the rainy season forms, he observes of the bitter sort that it “is of smaller growth and of a creamy-white colour when young, turning to a rusty colour at the ends as it ripens. This answers nearly to the description of the one called the ‘White Turkey.’ It is the better of the two for stewing, cooked in which

C. 2294

The Cucumber.

CUCUMIS
trigonus.CULTIVA-
TION.

way it affords a very delicious dish during the rains, when so few other

grown in October it may be made to yield. This is a point of some interest, since, if derived from the Indian wild stock, cultivation in Europe has completely changed the character of the plant. A writer in the Agr. Horticultural Society's Journal (IV, 21) says, however, that in importing seed of cucumbers, only those grown in the open air should be got, frame cucumbers are useless for India. He recommends that they

2295

DOMESTIC.
2296

on *Shraavan shudh 5th* (*Nagpanchmi* day). It is likewise employed in the worship of many other gods" (*Lisboa, U. Pl Bomb*, 285).

C Hardwickii, Royle, has been alluded to as most probably only the wild state of the cucumber. At the same time it bears separate vernacular names and is collected and sold for so very different purposes that it deserves an independent notice. It is known as the *air-dlu* in Kumdon

2297

Cucumis trigonus, Roxb., Fl Br. Ind., II, 619

2298

Syn — C PSEUDO COLOCYNTHIS, Royle, C TURBINATUS, Roxb., C MADER ASPATANUS Roxb., C MELO, Linn, var AGRESTIS, Naud., C PUBESCENS Wall., C ERIOCARPUS, Boiss., BRYONIA CALLOSA, Herb. Rottler

These are the synonyms of *air-dlu* in the *Flora of D. & L. I. 3* L. 111

This may be indicated thus —

C. Melo, Linn

2299

Var α agrestis, Naud.; SYN C MELO, var PUBESCENS Kunt (Trans

101 000

Var β culta, Kunt, SYN C DLDAM, Linn. C FLEXIOSUS Linn.; W. & A. Prod., 342, C AROMATICUS, Royle, Ill. Hb. Bot., pl. 2, p. 220

C 2299

Wild Forms of Cucumis.	CUCUMIS trigonus.
<p>those of <i>C. trigonus</i> or <i>C. turbinatus</i>, and in fact are almost reniform and often small is fruit other 'anjáb, fruits ood by the natives and much esteemed, yet they never take the trouble to cultivate the plant" Atkinson states of the North-West Provinces, that "<i>C. pubescens</i>,</p>	2308
<p>he says it is much less bitter than <i>var. pseudo colocynthis</i>, "and is commonly used as a vegetable after having been soaked in salt and water, the seeds of these cucumbers (sic) are considered cooling and are applied to Herpes, after they have been beaten into a paste with the juice of the <i>Daria</i> (<i>Cynodon Dactylon</i>)"</p>	MEDICINE, 2309
<p>4 <i>Cucumis pseudo colocynthis</i>, Royle. Syn — <i>C. PUBESCENS</i>, Willd ; <i>C. ERIOCARPUS</i>, Boiss ; <i>C. CUCUTRISATUS</i>, Stocks Vern — <i>Indrayan</i> (= <i>colocynth</i>), <i>bislámbe</i> in Northern India (O'Shaugh- nessy) ; <i>Kant Boun</i> ; <i>C. Hardwickii</i> (= "<i>var. pubescens</i>") ces be</p>	2310
<p>properties of this plant and of <i>C. Hardwickii</i> "the po these</p>	2311
<p>properties of this plant and of <i>C. Hardwickii</i> "the po these</p>	MEDICINE, 2312
<p>properties of this plant and of <i>C. Hardwickii</i> "the po these</p>	2313
C. 2313	

The Squash Gourd

CUCURBITA
maxima.

Botanic Diagnosis—Leaves, 5 palmate, lobes rounded, sinus narrow; petiole, nearly as long as the blade, not prickly; fruiting peduncle, round smooth, corolla lobes, curved outwards, calyx segments, lanceolate-linear.

2317

Habitat.—C of the globe.

2318

as the musk-m

find either C

the other hand a writer in the *Indian Forester* (IX., 202), and apparently, Mr. Gollan of Saharanpur, says—"Kudu (pumpkin) *Cucurbita maxima*" is

2319

chata), and Voigt, who wrote after Roxburgh, describes only *C. maxima*, to which he reduces Roxburgh's *C. Melopepo*. Stewart gives an account of all three plants collectively under *C. maxima*.

Oil.—The seed yields an oil.

OIL.
2320
MEDICINE.
2321

Medicine.—The seeds are used medicinally; the oil as a nervine tonic. The pulp of the fruit is often used as a poultice

§ "Also called in Panjáb *Ghia kaddu*. The fruit cut into small circular chips is a good application to relieve the burning of hands and feet in fevers" (*Asst. Surgeon Bhagwan Dass (and), Surgeon, Rawal Pindi, Panjáb*). "The pulp is used as a poultice to boils and carbuncles"

Early appearance of *C. maxima*—L

Food.—This plant produces the largest known cucurbitaceous fruit, in some cases weighing as much as 240 lb, and measuring nearly 8 feet in circumference. The fruit is wholesome, and when young is used as a vegetable. It is sweetish and yellow. When mature it will keep for many months if hung up in an airy place. It is largely used by natives of all classes in curry. "When very young and tender it may be employed as a pleasant vegetable for the European table, by being boiled, press-

FOOD.
2322

C. 2322

CUCURBITA
moschata.

The Musk Melon.

ed down to extract the water, and served warm, with butter, salt, and pepper" (*Mr L. Liotard*).

Mr Gollan says of "*kudu* (pumpkin) *Cucurbita maxima*" that there are several varieties of this plant common in the gardens as a rainy season vegetable. The commonest one is a large globular gourd and of a brown colour. The young fruit resembles the vegetable marrow in flavour but the full grown fruit is also very good. The seeds should be sown from April to June. The plant requires very rich soil and the general treatment is the same as that for *Lagenaria vulgaris* (the *Al kudu*).

2323

Firminger remarks of the "Red Gourd" or *sufuri-kumra*, also *Lal-*

carrots are, it can hardly be distinguished from them either in appearance or flavour. An annual seed sown in the rains; vegetable in use during the cold season, not often cultivated in gardens. It may be suspected that Firminger alludes in the above to *C. moschata* (forma

2324

mon Gourd
dian writers
vulgaris the
"Cucurbita
of the same
kind, while

2325

Cucurbita moschata, *Duchesne, Fl. Br Ind, II, 622.*

THE MUSK MELON, *Eng*, POTIRON, *Fr.*

Syn — *C MELOPEPO, Roxb*

Vern — *Sitaphal, saphari kumhra, kumra, kaddu, mitha kaddu, N-W P.; Kali-dudhi, BOMB*

This is said to be the *Abobrade Guinea* of the Portuguese in India.

2326

Botanic Diagnosis — Leaves as in the preceding but very often marbled with whitish blotches. Petiole hairy but not prickly. Fruiting peduncles angular and furrowed, calyx segments of the female flower large foliaceous.

There are two primary forms — one with the fruit smooth but mottled brown and yellow (*C. moschata* proper), and the other with the fruit torulose or fluted, with 15 to 30 ridges (*C. Melopepo, Roxb*).

Habitat — Very extensively cu

2327

The long account given by Firminger (*Van Gar for India, 128*) under the heading "*C. Melopepo, squash*" has reference to imported seed of *Squash, Gourd* or *C. moschata*.

North-West
not live in the
(in *Field and*
of *Cucurbita*
cultivation,

The Pumpkin or Vegetable Marrow.

CUCURBITA
Pepo.

season, &c. They state that only the *Cucurbita* there figured appears to occur in the North-West Provinces. Their plates seem to represent the form Roxburgh called *C. Melopepo* and not his *C. moschata* proper, if the idea be correct that the fluted fruit is *C. Melopepo*.

OIL.
2328
FOOD.
2329

C. Melopepo which would have answered to Mr. Powell's description of *tindá*.

2330

Cucurbita Pepo, DC.; *Fl. Br. Ind.*, II., 622.

2331

THE PUMPKIN, VEGETABLE MARROW.

Syn.—*C. Pepo*, Roxb

Roxburgh included this plant (the pumpkin) as well as *Benincasa cerifera*, Savi (the white melon) under one species. Atkinson, Drury,
P. M. Mendenhall, &c.

mouth, and the anthers are more or less united. The fruits of *Benin-*

does

Botanic Diagnosis.—Leaves 5-palmate, sinus, broad and segment pointed, petiole as long as the blade, the hairs of the lower surface

OIL.
2332
MEDICINE.
2333

nal applications for burns.

2 T

C. 2333

CUMINUM
Cuminum.

The Pumpkin or Vegetable Marrow.

MEDICINE.

Special Opinions —§ "The seeds are anthelmintic and used in cases for round worms though uncertain in action" (*Civil Surgeon F. H. Thornton, B A, M B, Monghyr*) "Grubler has isolated from pumpkin seeds a crystallisable variety of albumen Hemp and castor oil seeds also contain a similar crystallisable substance" (*Dr. F. H. Thornton, Calcutta*)

FOOD
2334Rolled
2335Twigs
2336Sherbet.
2337DOMESTIC
2338

seen the pumpkin (*C. Pepo*) in Assam, although the two fruits named are common in Assam, Cachar, and Manipur. The system of boiling in *klar* water is, however, very interesting to whichever fruit it applies and so also is the fact that the young twigs are eaten as a pot-herb. Under the names "*C. Pepo*, *DC*, pumpkin or white Gourd—*kumhra*, *kumara*, *kadimih peth* (in places), *kondur* the *lauka* and *kaddu safet* of Bijnor," Mr Baden Powell, and after him Mr Atkinson record an interesting fact which most probably should be given under *Benincasa cerifera*. "A sherbet is made by filling the hollow centre with sugar and exposing it to the sun until it becomes acid."

Domestic and Sacred Uses —"The seeds are recommended for the use of this plant considering it and *Narad* priest of the gods tell of this cucurbitaceous plant (*vide page 310 of Vaidya in Sanskrit from P. 1. 1. Puran*) Its fruit is also cut with some ceremony, called *kohala muhurt*, a day or two before a marriage" (*Lisboa, U Pl Bomb, 285*)

CUMINUM, Linn, Gen Pl, I, 926

2339 Cuminum Cuminum, Linn, Fl Br Ind, II, 718; UNDELLIFERE.

CUMIN, Eng, the *Κυμινον* of DIOSCORIDES, CUMINUM of HORACE and PERSIUS

Vern —*Zird*, HIND; *Jiraka*, *jiraka* or *ajdi* (Ainslie) "*Jiraka*, *jirana*" (Elliot), SANS; *Jira*, BENG; *Jira*, *jira utmi*, GR; *Jire*, *jire* MAR; *Kamin*, ARAB; *Zira*, PERS; *Zero* SIND; *Shiragam*, TAN; *Jiraka*, "*jirakarra*" (Elliot), TEL; *Jiringe*, *jirage*, KAN; *Jirakim*, MAL; *Duru*, *sudu duru*, SING; *Ziya*, BURM

A considerable amount of confusion exists in the vernacular names for this plant, *Zira* or *Jira* being also applied to *Carum Carui* (See C 681).

The Black Cumin of the Bible—*Nigella* of Hippocrates and Dioscorides, and the *Gill* of Pliny is *Nigella sativa*

Habitat —More or less cultivated in most provinces of India except perhaps Bengal and Assam. There seems no doubt the plant is not a native of India. Roxburgh is silent on this point, but Ainslie, who wrote

The Cumin

CUMINUM
Cuminum.

about the same period says of the Calcutta Botanic Gardens (which were then under Dr Roxburgh) that "the plant, however, is growing in the Bo

2340

and *and*, the quantity seems enormous. The same authority also gives 25
maunds as exported by that route. Atkinson makes no mention of
his Cata-
in Fe-
North-
the plant

References—*Roxb, Fl Ind, Ed C B C, 271, Vahl, Hort Sub Cal,*

Oil—A medicinal oil is prepared from the seeds (=fruits)

Medicine.—As a medicine Cumin seeds are considered aromatic, car-
minative, and stimulant. They are also stomachic and astringent, and
useful in dyspepsia and diarrhoea. The *Pharmacopæia of India* says

OIL.
2341
MEDICINE
2342

CUMINUM
Cuminum.

The Cumin

MEDICINE

or Persian, Nabti or Nabthean, Kirmāni or black Cumin, which they say is the Bāsilikon of the Greeks and Shīnū of Syrian. They consider it to have the same properties as the cumin (Dymock). Dutt says that "made of cumin seeds with the" to be applied externally for

...ive and stomachic, half drachm doses, in combination, never alone (*Assistant Surgeon Nehal Sing, Saharunpore*) "Seeds mixed with lime juice are used in bilious nausea in pregnant females" (*Surgeon-Major J. J. L. Ratton, M.D., M.C., Salem*) "Sinfud sirā is taken internally shortly after child-birth to increase the secretion of milk" (*Civil Surgeon R. Gray, Lalore*). "A quantity of the seeds lightly smeared with ghee put into a pipe and smoked relieves hiccup" (*Surgeon-Major D. R. Thomson, M.D., C.I.E., Madras*) "A reputed galactagogue" "Practitioner" (Nov. 1881, Vol. XXVII, p. 385, and p. 161 (quoting *Lancet*, 1872) however denies this action" (G. B.)

CHEMISTRY.

2343

Chemistry.—The chemistry of cumin has been dealt with fully by Flückiger and Hanbury (*Pharmacos*, 332), and their account reproduced in Dymock's *Materia Medica* (2nd Ed., 369). It is not necessary therefore to repeat the information there given, since either of the works referred to is likely to be in the hands of the student of Indian Materia Medica. Professor Warden has, however, contributed the following brief note for the present publication:—

"The fruit contains an essential oil, which is a mixture of Cymol and Cumimol, and other hydrocarbons. Cymol is also a product of the dry distillation of coal tar."

FOOD.

2344

the natives

TRADE

2345

Trade.—Cumin (or Cummin) would appear to have been known to the ancients, at least there are names for it in most of the classical languages. During the middle ages it was one of the most favoured of spices. In one instance it is recorded that during 716 A.D. an annual provision was made for the army of most of Europe. In the 14th century it was in frequent use, and in 1453 it was one of the commodities which the English had to pay for. The weighing and oversight

Foreign

Trade.

2346

At the present day the European demand has greatly declined, the export of Cumin from Bombay 20,040 cwt. from Calcutta in the year 1870-71. The quotations, since only about one-fourth of those amounts left India, the remainder represented the coasting traffic, and hence a further error, since some of the coasting imports into each of the ports named would have

2347

The Weeping Cypress.

CUPRESSUS
funnebris

reappeared again in the foreign exports therefrom. Thus of the exports from Calcutta 14,037½ cwt went to other Indian ports, nearly 2,000 cwt going to Bombay, an amount which must have greatly influenced the Bombay exports of the year. These remarks have been considered necessary owing to its being customary to find India assigned a far larger share in the world's trade in Cumin than is justified by the official returns. An analysis of the figures for the year 1875-76, compared with those for 1886-87, will remove this misconception. Last year the total exports were—Indian grown Cumin 9,051 cwt + foreign imports re-exported 1,260 cwt, or a total of 10,311 cwt. This amount was valued at Rs 41,486. In 1875-76 the total exports were 8,120 cwt, valued at Rs 4,919. The foreign trade in Cumin has thus slightly improved, but it falls far short of what most readers would infer from the amounts quoted above as exported from two of the Indian ports. Of the foreign imports, India received in 1875-76 only 538 cwt, and last year 2,020 cwt, so that deducting the re-exports, 760 cwt was thus added to the amount locally produced in 1886-87. But of the foreign imports 1,994 cwt came from Persia and the remainder from Turkey in Asia.

TRADE.
Foreign
Trade.

2348

to Sind
to Bomba
cwt C
and Ea

cwt, and the United Kingdom only 95 cwt.

The Indian internal trade in Cumin must be at least four times as extensive as the foreign, but the ramifications of road, rail, river, and coast-

Internal
Trade.
2349

Indian market

Dr Dymock says of the Bombay traffic in Cumin that it "comes from Jubbulpore, Guzerat, Rutlam, and Muscat. Value, Rutlam, Rs to Rs per Surat maund of 37½ lb, Muscat Rs to Rs½, Guzerat, Rs to Rs½, Jubbulpore, Rs to Rs."

2350

Domestic and other Uses—By the ancients smoking Cumin seeds was considered to produce pallor of the countenance.

DOMESTIC.
2351

Cuprea Bark, the bark of *Ramya purdicana* or *R. pedunculata*, see Cinchona, C 1152

CUPRESSUS, Linn, Gen Pl, III, 427

[Timb, 410, CONIFER

Cupressus funnebris, Endl, Brandis, For Fl, 534, Gamble, Man.

2352

THE WEEPING CYPRESS

Vern.—Chandang, Ichenden, BHUTIA

Habitat—A handsome tree with pendulous branches, and a fibrous brown bark, often planted in Nepal, Sikkim, and Bhutan, near temples and monasteries, and in China (*Gamble*)

C. 2352

CUPRESSUS
torulosa.

The Cypress

2353

Cupressus glauca, LamHabitat—Very generally cultivated in Western India above the Ghats (*Dils & Gibs, Bomb Fl Supp, 83*)

2354

C. sempervirens, Linn

THE CYPRESS

Vern—Sara, saras, N-W INDIA, Firath, SIND, Sarābake, MAR.

Reference—*Fl. Ind. Brit. Mus. Nat. Hist.*

Habitat—A large tree growing on the outer ranges of North-West

Himalaya, from Chamba to Nepal, scattered or in numerous isolated localities of greater or less extent, chiefly on limestone, between 5500 and 9000 feet. Common on the north of the Shikhar, Simla, and at Nainital.

Medicine.—Wood and fruit are regarded as astringent and antihelmintic.

Stem—The wood is very hard.

Very hard.

It is used for trunks and boxes, the contents of which are proof against most insects (*Brandis*).

2358

C. torulosa, Don

HIMALAYAN CYPRESS

Vern—Dev-dhar, RAVI, Deodar, KULU, BHAJI, Gulla, gulrai, kailain, SIMLA, Leauri, JAUNSAIR; Kaisalla, sarai, KUMAON, Sarai, sarahyu, TIDET.

References.—*Vogel, Hort. Sub. Cal., 558; Brandis, For. Fl., 533; Fl. Ind. Brit. Mus. Nat. Hist.*

Habitat—A large tree growing on the outer ranges of North-West Himalaya, from Chamba to Nepal, scattered or in numerous isolated localities of greater or less extent, chiefly on limestone, between 5500 and 9000 feet. Common on the north of the Shikhar, Simla, and at Nainital.

RESIN

2359

TIMBER

2360

sacred ark. It is often burnt as incense in temples. The *Indian Forester* (Vol. 1, 63) gives the following analysis of the ash:—

Soluble potassium and sodium compounds	0.004
Phosphates of iron, calcium, &c	0.039
Calcium carbonate	0.041
Magnesium carbonate	0.008
Silica with sand and other impurities	0.004

TOTAL . . . 0.099

C. 2360

Copper.	CUPRUM.
CUPRUM or COPPER.	2361
<p>Cuprum; <i>Man. Geol Ind., III., 239, IV., 4</i> COPPER, MINERAL DE CUIVRE, <i>Fr.</i>; KUPFFERERZ, KUPFER BLENDE, <i>Germ.</i>, MINERALE DI RAME, <i>Ital.</i> Ve—</p>	
Re	
<p>Consult also the numerous publications referred to by Ball (<i>Man Geo Ind., III., 611</i>) DISTRIBUTION OF COPPER ORES IN INDIA—The following brief note has been furnished for the present publication by H. B. Medlicott, Esq.,</p>	DISTRIBUTION. 2362
<p>mining has been practised on a large scale, but is now almost extinct. In Afghanistan, copper ores have been mined to a considerable extent at various places. In the Kumaon and Garhwal districts of the North-West Provinces, copper deposits occur which have been several times unsuc- cessfully worked by the natives. In the Darjiling hills, copper occurs in small quantities, and has been worked in the Karnai and Nellore districts of the Madras Presi- dency."</p>	2363
<p>For detailed information regarding the Indian mines and sources of copper ore the reader is referred to Ball's account in the <i>Manual of</i></p>	
C. 2363	

CUPRUM	Copper.
DISTRIBUTION	
	<p>and also in several of the groups of transition rocks, as, for example, in the Cuddapah, Bijawar, and Arvali groups. In extra peninsular India they are found for the most part in highly metamorphosed rocks, the precise age relations of which to those of the peninsula are not in all cases clearly made out as yet.</p> <p>"The ore of most common occurrence is the copper or pyrites but towards the outcrops it is commonly altered into carbonates or oxides. The associated minerals are in general identical with those which are found under similar circumstances all the world over. Recent analyses by Mr. Mallet have tended to clear up much of the uncertainty which</p> <p>tions, the copper ores of India are not generally in the form of true lodes but are either sparsely disseminated or in the form of small, irregular, massive bunches and nests in cracks and fissures traversed by veins. In not a few cases it is best described as a copper-bearing matrix in which the ore is disseminated. In not a few cases it is best described as a copper-bearing matrix in which the ore is disseminated.</p> <p>flows through tertiary metal, reaching up to the surface, and were common in Ladakh, see Geological Museum (weighing about 21 oz) cut from a lump of some</p>
FOREIGN TRADE 2365	<p>old copper, unwrought and wrought copper, amounted to 615,039 cwt, valued at Rs. 1,99,40,085. For the past 20 or 30 years the imports of copper have steadily increased with the increased agricultural prosperity of the people, but within that period they have borne a marked relation to the fluctuations of agriculture. In the year 1885-86 the imports amounted to 1,00,000 cwt and in 1892-93 to 1,00,000 cwt.</p>

Copper Sulphate.

CUPRI
Sulphas.FOREIGN
TRADE

of this year it had further fallen to
ne much lower, falling below £45
than it has ever been, being more
price ever known, and authorities
state 30 per cent below what the trade had previously considered a
safe and moderate price. This decline is due to a greatly increased
production in the United States, and it would seem to those who are in
a position to estimate the conditions of future production there and else-
where in Calcutta,
and it has
50 per cent
three-fourths
every year
is trade is,

2366

however, more apparent than real, as a large proportion of it is due to the
fact that it comes direct to India instead of to England. This direct ship-
ment is of great value, as it means that the commercial relations of India
with Australia are becoming more intimate.

Cupri Sulphas.

2367

COPPER SULPHATE OF BLUE STONE

Vern — *Nīla thutha*, *nīla tuta*, *nīla tuti*, *a*, HIND, *Mōr tuttā* or *mōdr-*

References — *Pharm Ind*, 378; *Moodeen Sheriff's Supp to Pharm Ind*,
123, *U C Dutt, Mat Med Hind*, 66, *Waring's Basar Med*, 46

MEDICINE,
Salts
2368

vapradāsa says it contains some copper and therefore possesses some of
the properties of that metal. It is described in this work as astringent,

2369

2370

porating to crystallization

2371

According to European Medical practice pure sulphate of copper is
tonic, astringent, emetic; in large doses an irritant poison. Locally ap-
plied in substance to a denuded or granulating surface, mildly caustic,
styptic, and in solution stimulant. The article so used is imported from
Europe. It is largely used in chronic dysentery, diarrhoea, epilepsy,
chorea, and hysteria. Locally, it is applied in solution in gonorrhoea,
leucorrhoea, purulent ophthalmia, weak ulcers, superficial hæmorrhage,

2372

C. 2372

CURCULIGO
Orchioides.

Copper Sulphate

MEDICINE.

and, in substance, to crancrum oris, aphthous ulcerations, exuberant granulations, and granular conjunctivitis (*Pharm. Ind.*) Waring recommends an emetic of 5 grains of sulphate of copper in tepid water for Opium, Datura, Nux Vomica, Cocculus Indicus, Bish (Aconite), Arsenic, or other poisoning cases. If it does not operate in half an hour it may be repeated.

Spe C
is a re
nally "
calcine
expecte

Robb, Ahmedabad) "Sulphate of copper is used internally as astringent in chronic dysentery and diarrhoea in dose of $\frac{1}{4}$ to $\frac{1}{2}$ of grain, also applied externally" (*Asstt Surgeon Nehal Sing, Saharunpore*) "Copper coins, on which there is a deposit of verdigris, are kept for an hour or two in a mixture of (ripe) tamarind and water, and then rubbed on parts of body attacked by urticaria" (*Honorary Surgeon P Kinsley, Chica-cole, Ganyam, Madras Presidency*) "Useful as an emetic in cases of poisoning" (*Civil Surgeon J H. Thornton, B A, M B, Monghyr*) "Copper foil (Shabari, Swahili, E. Africa) cut into small pieces about an inch or more square, which are spread over the chest before and behind is the native (African) treatment of cough and all general chest troubles. Two dozen of these thin copper plates were counted in a case that came up for other treatment; their application is on the principle of a series of small blisters or counter-irritants" (*Zanzibar*).—*Surgeon-Major John Robb, M D, Surat, Bombay Presidency.*

Plates.
2373

Leaf
2374

COPPER LEAF—A thin copper foil is sold in the Muscat bazar as an external application to unhealthy ulcers. It is applied like thin Gutta-percha tissue over the surface of the ulcer and secured for days by means of a bandage.

CURCULIGO, *Gartn*, *Gen Pl.*, III, 717

[p 124, AMARYLLIDÆ.

2375

Curculigo orchioides, *Gartn*, *Baker, Linn Soc Jour.*, XVII,

Most authors refer the native medicinal tuber known in the Panjāb as *siyah musli* to this plant, but Stewart says it is obtained from *Anilema tuberosa*, *Ham*, and Dymock describes it under *Hypoxis orchioides*, *Willd*, giving *Curculigo orchioides* as a synonym. In Bengal the tuber is generally known as *Tal lura*.

Syn.—CURCULIGO MALABARICA *Wight, Ic*, t 2043, HYPOXIS ORCHIOIDES *Kunt*, in *Ann Mus Lug Bat*, IV, 177, ORCHIS AMBOINICA *MAJOR*

BOMB.
(varaki,
"TAM;
iti gauda,

i, I, 242
rm Ind,
fat Med
m Ceylon
Rhetde,

Siyah Musli

CURCULIGO
orchoides.

Habitat—A small herbaceous plant with a rosette of radial leaves and tuberous root, native of the greater part of the hotter regions of India and Ceylon. Roxburgh says that in cultivation it flowers all the year round.

Medicine—In most Hindu and Muhammadan works on Materia

MEDICINE
Black root

white *Asparagus adscendens*. According to some writers the young roots of *Bombax malabaricum* constitute one of the white *muslis*, and by others the black and white forms are obtained from one and the same plant during different stages of its growth. Dr. Moodeen Sheriff remarks that in South India a false *sufed musli* is sold which is obtained from *Asparagus sarmentosus* (A 1577). On the other hand Dr. U. O. Dutt says "The roots of *Bombax malabaricum* and of *Asparagus racemosus* are sometimes sold by the native druggists of Calcutta under the name of *sufed musli*. These are clearly not the same as the *musli*." The tuberous roots of

C. ensuifolia as the *kali musli*. He further states that much of the latter root sold in the Bombay Presidency is *Ancilema scapiflorum*, Wight (Conf. A 1122). Dr. Dutt says of *C. orchoides*—The tuberous roots of

2379

and sometimes given with milk and sugar, in doses of two drachms in the twenty-four hours in cases of gonorrhoea.

says "A
gonorrhoea
disiac, &

Medical

Rutiam
maund of 3 1/2 lb

TRADE
2380

Wild Arrowroot.

CURCUMA
angustifolia.

found wild in the North-West Himalaya. The flowers are large and yellow, longer than the bracts, they expand in the morning and wither in the evening of the same day.

Cultivation of East Indian Arrowroot.—Perhaps the most complete accounts of the cultivation of this plant are those which will be found in the Reports of the Saidapet Experimental Farm, Madras. The following

CULTIVA-
TION
2386

Madras Root-
stocks.
2387

the above yield would represent an outturn of 493lb of flour per acre. In another case in the College Experimental Garden, a plot, measuring 1,160 square yards, planted with this crop yielded 1,798lb, or at the rate of 7,500lb per acre. The culture of this crop is very simple: it is only necessary to plant the sets in properly prepared soil, and to water them occasionally during the dry season. The removal of the crop is tedious unless the tubers can be ploughed out, as potatoes are done in England, which is seldom possible owing to the dryness of the soil, so that the tubers have to be dug up. The preparation of the flour is also very simple and easy. The TUBERS have only to be reduced to pulp on a grater, after being well washed to remove soil and dirt, and then the pulp is mixed thoroughly with water so as to separate the starch completely from the fibrous matters. The whole is afterwards strained through cloth, through which the STARCH and water passes, and the fibre left behind. After this the STARCH has only to be thoroughly washed by decantation with clean water, and dried in the sun. It is then rolled on a table to break it up thoroughly into fine flour and is ready for sale. The flour can be produced at a very low price; it could be sold profitably at 4 annas per pound. And thus 400 rupees per acre could be realized. This is a remarkable return and should also be published for the information of the public."

Starch,
2388

Profits
Rs. 400
an acre.

"The following extract from a letter from the Collector of South Kanara, dated 10th March 1882, No. 517, will be found interesting: "With reference to paragraph 48 of your report on the Saidapet Farm, recorded with the Board's Proceedings dated 10th December 1881, No. 3182, I

South Kanara.
2389

plant in this district (with its annual rainfall of about 130 inches between June and November) would be thankfully received. The plant, I believe,

a angusti-
nation and

CHEMISTRY,
2390
Inferior to
Maranta.
2391

yielded by sample marked '1st sort' is of a superior description and nearly as good as that of the Maranta. This sample is susceptible of further

C. 2391

Wild Arrowroot

improvement it contained a number of extraneous matters, black particles, and the process of The three sam presence of slight ion of the starch the Farm sample

Solar heat to be avoided

Use of Caustic Soda

immediate conversion of caustic soda water for steeping found useful in Thorough washers soda."

The arrowroot is said to be largely manufactured at Cochin Travancore, and Hanari Royle says that "a very excellent kind called *tickar* is also made at Patna and Baglupore from the tubers of *Batatus* (*Ipomoea*) *edulis*."

Medicine—The arrowroot is used medicinally in some parts of the country.

Food—A good quality of arrowroot is prepared from the tubers especially in Travancore, where the plant grows in abundance. Roxburgh observes that a sort of starch or arrowrootlike secula is prepared which is sold in the markets of Benares, and is eaten by the natives. The flour, when boiled in milk forms an excellent diet for patients or children. It is largely used for cakes, puddings &c., though it is often complained of as producing constipation. The granules much resemble the ones of arrowroot, but are smaller and always stratified.

men in Bombay use it to thicken milk which is often watered'. The edible properties of the tubers of this plant are alluded to in most of the

Thicken milk
2308

PREPARA-
TION OF
ARROWROOT
Travancore
2300

prepared The process adopted in the Upper Godavari District (No. 505) is thus referred to "*Tankir* or *Tikhar* is a description of 'tro' "

abbed
e ther
azars

Wild Turmeric

CURCUMA
aromatica.

for export" (For further particulars see the paragraph on Cultivation)

PREPARA-
TION OF

2403

Malabar.
2404

be trusted as referring to this or to the true arrowroot See *Maranta arundinacea*

Dymock remarks of Turmeric (*Curcuma longa*) that the starch "of the young tubers at the end of the radicles, which are nearly colourless, forms one of the Last Indian arrowroots. It is to be observed that the tubers that yield only starch when young will yield turmeric when old, the colouring matter and aromatic principles are deposited in the cells at a later period of growth."

Turmeric.
2405
Starch.
2406

Curcuma aromatica, Salisb., Roxb., Fl Ind., Ed C B C., 8.

WILD TURMERIC, YELLOW ZEDOARY, COCHIN TURMERIC

Syn — *CURCUMA ZEDOARIA*, Roxb.

Vern — *Jangli haldi*, *ban haldi*, *ban haridra* (*jedwar*?). HIND *Ban*
halud, BENG *Kasturi man*
kattu manna
Roxburgh

SING, *Aiydsanoin*, BURM

References — *Voigt*, *Hort S*
Ainslie, *Mat Ind* 1, 49
125, *U C Dutt*, *Mat Me*
Ind 769, *Year Book P*
regarding *Pharm Ind*,
clap 859

Habitat — Roxburgh says of his *Curcuma Zedoaria* "This beautiful species is a native, not only of Bengal (and common in gardens about Calcutta), but is also a native of China, and various other parts of Asia and the Asiatic islands. Flowering time, the hot season, the leaves appear about the same period or rather after, for it is not uncommon to find the beautiful, large, rosy, tufted spikes rising from the naked earth before a single leaf is to be seen." "The plant when in flower is highly ornamental, few surpassing it in beauty, at the same time it possesses a considerable degree of delicate aromatic fragrance."

The flowering spikes are quite distinct from the leaf bearing stems,

Bengal
2407

Malabar.
2408

Concan.
2409

observe that the leaves when young have a central purple stain which

C. 2409

CURCUMA
aromatica.

Wild Turmeric.

Mysore
2410
Travancore.
2411
HISTORY.
2412

almost disappears when they attain their full size" Drury remarks that it is abundant in the Travancore forests Of Mysore Mr D E Hutchins says *C. aromatica*, the *Kad arasina*, is collected from the forests all over the province

History of *Jadvar* and *Zedoary* —The reader is referred to *Aconitum heterophyllum*, (A 401 & 403), for further particulars regarding the use of the Arabic word *Jadvar* According to certain writers (including Roxburgh) this is applied to a species of *Curcuma*, presumably the present species To Dr Moodeen Sheriff we are indebted for the results of much careful

that
of th
ject
or Za

sian works also used as synonymous, but the former is more correctly the name for the medicinal wood obtained from a species of *Berberis*

2413

of the Bhotias, who

s
p
s

2414

is used by

ahlo bikh

2415

kh of the

hobikh is

given as a tonic in dyspepsia, fevers, and asthma Lastly, a plant never before recorded as used medicinally, namely, *Caragana crassicaulis*, is

2416

Wild Turmeric

CURCUMA
aromatica.

Nepal, "must not be confounded with the word *Nirbisi*, which is the Sanskrit for *Curcuma Zedoaria*" To the hill tribes around Simla and Kulu, at

often more than a palm and diameter to the root is strongly camphoraceous" Dalzell and Gibson say. "The tubers of the root are palmate"

DESCRIP-
TION
2417

DYE
2418

Cosmetic.
2419
MEDICINE
Rhizomes
2420

to promote eruptions" Ainslie says the Muhammadans suppose it to be a valuable medicine in certain cases of snake-bites, administered in small doses, and in conjunction with golden-coloured orpiment, *kust* (*Costus arabicus*), and *ajwan*"

Special Opinions.—§ "Used externally in scabies and the eruption of small pox" (*Surgeon-Major Henry David Cook, Calicut, Malabar*). "Rubbed into a paste with benzoin is a common domestic application to the forehead for headache" (*Surgeon-Major John North, I M S, Bangalore*) "Applied to the forehead in cephalalgia, and a cosmetic."

CURCUMA
caulina.

Black Zedoary.

(T. Ruthnam Moodelliar, Native Surgeon, Chingleput, Madras Presidency.)

TRADE.
2421

Trade.—"The Bombay market is supplied from the Malabar coast. Value, unpeeled R24 to R25 per candy of 5½ cwt; peeled R27 per candy" (*Dymock*).

2422

Curcuma cæsia, Roxb.; *Fl. Ind.*, Ed. C.B.C., 9.

BLACK ZEDOARY.

Vern.—*Lila holdi* or *nīl-kāñka*, BENG.; *Kālī halada*, MAR.; *Nar-kachāra*,
ROMA. *Nar-kach'ra* 10 1-12 11 - 12

3

Bengal.
2423
Dinapore
2424

Habitat.—Roxburgh remarks: "This elegant strongly-marked species is a native of Bengal, where it blossoms in May," and just before the rains. "In the deep ferrugineous middle of the leaves it resembles *C. Zerun*," "the flour of the roots" the Indian market. He adds "through Peters I have been supplied with living" inapore; he informs me that it is common in gardens in Bengal, and is used as a domestic

MEDICINE
Rhizomes,
2425

Per
mo^o
andCosmetic.
2426

cool under
same medi-

TRADE.
2427

Trade.—Dymock says the tubers are internally very hard and horny, of a greyish black, but when cut in thin slices of a greyish-orange. The odour and taste are camphoraceous. "The drug comes overland from

2428

C. caulina. Graham: Dalz. and Gibs., Bomb. Fl., 275.

Vern — *Chavara, chowar*, BOMB.

Habitat.—A plant common at Mahábaleshvar, Bombay, and described by the late Mr. Graham.

Food.—A form of ARROWROOT is said to be prepared from this plant.
It is embedded in St. Croix mud and other things, the last being

FOOD.
Rhizomes,
2429
Arrowroot,
2430

the bazaars at Bombay. In 1878, a European prepared a few hundred pounds of it, and sent samples to be tried by Messrs. Treacher & Co, Phillips & Co, and Kemp & Co, but it was found wanting in nutritive

C. 2430

The Tikor; Turmeric.

CURCUMA
longa.

"The preparation of Arrowroot at Mahābaleshvar is simple. The root (of which a cooly will gather 4 or 5 large basketsful a day for as many annas) is scraped, washed, and rubbed to pulp on a grater, as mortars are found to crush the globules. The pulp must then be washed no less than a dozen times at least, the sediment being stirred at each washing. The dark scum on the sediment and the muddiness of the water of the first washing slowly disappear, till when the sediment is pure-white it is allowed to harden into a cake, which is afterwards reduced to powder. A basketful of roots yields 3-4 lb of pure arrowroot."

2431

Curcuma leucorrhiza, Roxb., Fl. Ind., Ed. C.B.C., 10.

Vern.—Tikor, BENG

Ti

Habitat.—Roxburgh says this is a native of Behar. Mr. J. Glass,

paration of arrowroot from this plant, "the process for obtaining the starchy substance called *Tikor* is as follows: the root is dug up, and rubbed on a stone, or beat in a mortar, and afterwards rubbed in water with the hand, and strained through a cloth, the fecula having subsided, the water is poured off, and the *Tikor* (fecula) dried for use." Dr. Irvine (*Mat Med*, Patna) alluding to this species says its "fine amylaceous farina is equal to arrowroot."

FOOD.
Arrowroot.
2432

C. longa, Roxb., Fl. Ind., Ed. C.B.C., 11.

TURNERIC.

Vern.—Haldi, HIND; Halud, BENG; Haldar, halja, PB; Haridrā,

2433

D

sabā ghīn and the Persian *Zard chubak*. This is probably the *Kurkum* of Dioscorides. U. C. Dutt writes that the Sanskrit *haridrādya* or the two turmeric, signifies turmeric and the wood of *Berberis asiatica*, Moodeen Sheriff says that in many books *Kurkum* is incorrectly given to saffron, and that *haridrā* is also wrongly given to yellow orpiment, that substance being in Sanskrit *Haritā lakam*.

References.—Toigt Hort Sub Cal, 565, Thwaites, En Ceylon Pl, 316; Dals & Gibs, Bomb Fl, 87; Stewart, Pb Pl, 239, Manjella kua Rheed. Hort Ind VI 79; Rumph Am Gmelin and Mat Ind, Sheriff Suppl, 311; Dymock, Mat Med W Ind, 764; Fleming Med Pl and Drugs, as in As Res, Vol XI, 165; Flück & Hanb, Pharmacog, 638; U.S Dispens, 15th Ed 1629; Bent & Trim Med Pl, 4 269; S Arjun, Bomb Drugs, 140; L. L. Dey, Beng Drugs, 45; Murray, Pl and

C. 2433

CURCUMA
longa.

Turmeric

CULTIVATION

Drugs, Sind 21, Waring, Basar Med, 140; Year Book Pharm, 1873, p 113 Medical Topog, Ajmir, 135, Mason, Burma 513 863; Man Coimbatore Dist, 228, 229, and 230, Baden Powell, Pb Pr, 299 380

Habit

Condiment
Form
2434
Dye Form.
2435

rhizomes It is the well-known *haldi* universally used as a condiment with curry-stuffs and also as a dye, and is one of the most profitable of crops The dye-yielding rhizome is harder and much richer in colour than the edible These conditions are thus special adaptations which possibly point to an ancient cultivation At the same time, though several species of *Curcuma* are undoubtedly natives of India, some of which appear to have been mistaken for the true turmeric, there is little of a positive character that would justify the supposition that *Curcuma longa* itself is a native of India Simmonds (*Tropical Agriculture*, p 383) says "The *Curcuma longa* grows wild in the province of Mysore, and is probably indigenous to various other parts" On the other hand, Roxburgh and all botanical writers speak of it only as cultivated, and Anslie even remarks that "The *Curcuma longa* grows wild in Cochin-China, and is there called *Kuong huynh* Loureiro gives us a long list of its medicinal virtues in lepra, jaundice, and other disorders"

it in
at it is
have
superseded some of the indigenous *Curcumas* formerly in use and which bore the names now given to this plant, just as the true arrowroot plant is rapidly displacing the indigenous or East Indian species Dalzell and

Turmeric, see page 664

CULTIVATION

2436
Bengal
2437

CULTIVATION, YIELD, AND SOIL.

Bengal.—The earliest and to this day one of the most complete accounts of the cultivation of turmeric in India is given by Mr. J. H. B. This may be systems "The reflow during the all of them It is

Deshl
2438

ember and
oot The
publishes
erie culti-
ne known
as the *deshi* or country, and the other as the Patna variety The latter is of a richer colour and gives a better outturn Loamy soil, even of a very inferior quality, will grow turmeric It can be grown in shady

Turmeric.

CURCUMA
longa.

held being twice as large and 27 inches apart. Sugar-cane cuttings are very lightly covered with earth; over 6 inches of earth is placed on the turmeric cuttings. The usual planting time is the first week of Jaistya," that is, about the 20th of May. "The plants spring up in about a fortnight. One or two weedings are necessary, and care must be taken that the fields are not inundated. In some parts of Bengal it is not considered good practice to lift the plants the first year. On the setting in of the following rains new shoots appear and the plants are tended exactly as in the first year. 'After about a year and nine months turmeric is lifted' When it is raised the first year, as is the practice in some places, the produce is less in quantity and inferior in quality." The Director of Agriculture, Bengal, has the following estimate of the cost of cultivation:—

	R	a.	p.
6 Ploughings	2	4	0
3 maunds of seed at R ₃	9	0	0
Planting, 8 men at 4 annas a day	2	0	0
To earth up four times	4	0	0
Four weedings, 3 men at a time	3	0	0
Repairing the furrows, 4 men	1	0	0
To dig out, 6 men	1	8	0
To clean, 3 men	0	12	0
To boil, 6 men	1	8	0
To dry, 8 men	2	0	0
Earthen pots	1	0	0
Rent	4	0	0
TOTAL	32	0	0

Mature.
2440

It is not stated whether the cost of the seed is included in the latter being a thimble. Dr. McCann's extensive series of statements are made. Turmeric is planted in Rājshāhī in March to April and dug up a year later. In Sargān not at all. In August, varies from 1 to 2. In Hugli, R₂.

BENGAL
2441Season of
planting

general
Kumao
importa
grown

N. W. P.
2445

URCUMA
longa.

Turmeric

CULTIVATION

N W P

Cost,

2446

Print

2447

BOMBAY.

2448

Yield,

2440

Lokhandi

2450

Aromatic.

245I

PANJAB

2452

-45-

2453

2454

MADRAS.

2455

January"

Panjab —It is not apparently very extensively grown in the Panjab, at

turmeric for the consumption of the whole district" The Gazetteer further states that in the Kangra District there were, in 1880-81, 1,621 acres under this crop and in 1881-82, 1,520 acres

acres under this crop and in 1881-82, 1,520 acres.

Madras—Turmeric cultivation is alluded to in various publications regarding South India, but no article has been found that deals with the Presidency collectively. Of Coimbatore it is stated that it is usually grown as a mixed crop with yams, maize, castor, brinjal, onions, &c. "The soil is thoroughly prepared by repeated ploughing and heavy manuring, municipal sweepings and ashes being a favourite manure." "In June or July, the soil having been ridged up about 2 feet apart, the rhizomes are planted, a cubit or less from one another, on the ridges and thereafter watered every three or four days until the end of December, thenceforward somewhat less often till March and April, when they are dug up. The crop is hoed and weeded several times in the first four months. The other crops are variously planted, the onions on the

C. 2455

CURCUMA
longa.

Turmeric.

PREPARA-
TION.
PANJAB.
2462
MADRAS.
2463

the roots are soaked in lime juice and borax before being powdered instead of being boiled." Of the Panjáb, Mr. Baden Powell says the tubers are taken up in November and dried partly by the action of fire and partly by exposure to the sun. Of Coimbatore it is reported: The roots are carefully sized and separately boiled in a mixture of cow-dung and water, dried and sent to market."

AREA.
2464

AREA UNDER TURMERIC.

	Acres.
Bengal (according to Dr. McOann) perhaps	30,000
Madras	15,000
Bombay	6,000
Berar	2,000
Panjáb	3,500
TOTAL	56,500

TRADE.
2465

TRADE IN TURMERIC.

Regarding the Indian Foreign trade in this article Mr. O'Connor, in his Review of the Trade in 1875-77, wrote "Turmeric was exported to the value of 10½ lakhs of rupees, the quantity being 123,824 cwt. This article has hitherto been recorded in the returns under the heading 'Spices,' but it is more appropriately classed as a dyeing material. It is not really a spice but rather a condiment, and for this purpose

Foreign.
2466

portance. In 1881-82 the exports were 70,783 cwt, valued at Rs. 66,047, as compared with 1877-78, when they amounted to Rs. 12,40,189. In 1885-86 the trade had so far recovered itself that the exports amounted to 156,287 cwt, valued at close on 14 lakhs of rupees. Last year they amounted to 140,994 cwt., valued at Rs. 10,32,025.

Internal.
2467

Full particulars cannot be learned as to the extent of the internal trade, but it must be very extensive, and even a trans-frontier trade exists. Kashmir receives a considerable amount. The various Indian ports last year exchanged 281,117 cwt of turmeric valued at Rs. 24,38,260.

HISTORY
2468

HISTORY OF TURMERIC

Turmeric yields a yellow dye of a fleeting character, which formerly was far more extensively employed by the natives of India than at the present day. Its chief features that recommended it for decorative purposes at marriage ceremonies, &c., were cheapness, ease of preparation, and facility of being removed. But these are conditions even more readily attained by aniline colours, while glaringly brilliant results are obtained, and, consequently, even religious injunctions have

Turmeric.

CURCUMA
longa.

to a certain extent given place to the encroachments of the tar dyes. Writing of this subject Dr. McCann (*in his Dyes and Tans of Bengal*, p. 85), says: "Formerly on festive occasions an infusion of turmeric

HISTORY.

Wedding
Garments.
2469

ing off the evil eye.
all the body with it as

Cosmetic.
2470

in the sect of Vishnu
make the peculiar

Markings on
Foreheads.
2471

st Provinces, says:

Dye Fleeting.
2472

be rendered permanent as a dye." It is somewhat remarkable that John Huyghen Van Linschoten, who spent several years on the Malabar coast from about the date of 1596, should describe the races of people he met with, going into every detail as to their social habits, domestic and

He describes Car-
Tamarind, Ginger,
ration of curry and
chutney makes no mention of the habit of eating turmeric or of dyeing

nizable by mere verbal descriptions. The principal sorts now in commerce

C. 2472

CURCUMA
longa.

Turmeric.

HISTORY.

Cochin
Doubtfully
True Tur-
meric
2473

the trade of Cochin, makes no mention of Turmeric, but at the same time references occur, of turmeric as employed in Europe about the time of which

is horny and of a deep orange-brown, or when in thin shavings of a brilliant yellow Mr A Forbes Scaly of Cochin has been good enough to send us (1873) living rhizomes of this Curcuma, which he states is mostly grown at Alwaye, north east of Cochin, and is never used in the country as *turmeric*, though its starchy tubers are employed for making arrowroot" (Conf with *C. angustifolia* and other sources of East India arrowroot)

DYE

2474
Dye-Yielding
Rhizomes.
2475

TURMERIC DYE

Dye—It has already been stated that a special form of turmeric is grown for this purpose, namely, a harder root, much richer in the dye principle than in the ordinary condiment form. This dye rhizome receives separate names in the various provinces of India, but is most generally known by the name *lok handi*, *haladi*, other dye forms are as *mala-haldi*, *jowala-haldi*, and *ambi-haldi*. Under the paragraph, above devoted to an account of the preparation of the tuber, mention has also been made of the further process which the dyer has to adopt in

Yellow.
2476

The colour is only deposited in the rhizome with age, and hence, in all probability, the above mentioned forms have been obtained by a process of careful selection of stock observed to produce the colour freely. It is of importance, however, that the European merchant, in purchasing for dye purposes, should see that he gets the hard dye-yielding form and not the softer aromatic condition which is used as a condiment. Although, of course, turmeric is still employed by itself as a simple and cheap dye, its more general use at the present day in India, is as an auxiliary to other dyes and in Calico printing. It is also used to some extent to impart a colour to native-made paper. Mordants are but rarely required with the dye, as it is found to attach itself readily enough to wool, silk, or cotton. Alkalies deepen the colour, making it almost red. Alum is said to purify the colour and to destroy all shades of red. The dyers of Calcutta produce a brilliant yellow, known

(Carbonate of Soda)
this process "Here
int, produced always

are sometimes em-

ployed with turmeric, but the chief compound colour in which turmeric plays an important part is the green shades formed along with indigo. The fabric is first dyed with indigo and then dipped in a solution of *haldi*. Turmeric is also often added to sharpen or brighten other colours, as, for example, *Singrahar* (*Nyctanthes arbor-tristis*), lac dye, *al* (*Morinda tinctoria*), safflower (*Carthamus tinctorius*), and *toon* (*Cedrela Toona*).

Green.
2477

C. 2477

Turmeric.

CURCUMA
longa.

The Indian Calico-printers use turmeric by preparing a mixture of 4 gallons of water containing pomegranate rind and alum in the following proportions:—Turmeric 5lb, pomegranate rind 2lb, and alum 1½lb. The compound is left to stand for a night, the surface water strained off, and ½lb of indigo added. It is then prepared for use by being thickened with gum, clarified butter, and flour in the usual way. The colour is greenish yellow and is fleeting." (*Buck, Dyes and Tans of N.-W. P.*, 55)

The rhizome is used by the European and the Chinese in the aniline industry. The colour is very fugitive and is fleeting.

CALIC
PRINTING.
2478

EUROPEAN
USES.
2479

Cotton,
2480

Wool.
2481

Silk,
2482

Curcumin.
2483

Curcumin, and the
a new body
n. The sub-
st with pure
is dried, and
and 1 part of
s on cooling
By pseudo-
C. 2485

Action of
Boracic Acid.
Red color.
2484

Rosocyanin.
2485

CURCUMA
longa.

Turmeric.

EUROPEAN
USES

curcumin is understood the organic resinoid substance resulting from the prolonged action of water upon boro-curcumin, just above-mentioned

Blue Color.
2486

nia turns the alcoholouration changes niacal solution red the alcoholic solu-

Colouration of
Flowers.
Cyanin.
2487

rosococyanin) and pseudo-curcumin are unknown, neither was, until July,

Printing
Silks
2488
Sour Browns
2489
MEDICINE.
2490

alkalies If this suggestion proves correct, on more precise investigation turmeric could become a useful source of preparation of the red colouring matter of flowers, which it is very difficult to obtain by direct extraction

is now employed to a vast extent in stann-dyeing, forming an important constituent in certain compound colours, especially the so-called "sour browns."

decoction of turmeric in purulent conjunctivitis, he says it is very effectual in relieving the pain In coryza he states that the fumes of burnt

Special Opinions — § "The root, parched and powdered, is given in bronchitis in doses of grs xxx to xl" (Civil Surgeon F. Anderson, M.B., Bijnor) "The smoke produced by sprinkling powdered haldi over burnt charcoal will relieve scorpion sting when the part affected is exposed to the smoke for a few minutes A paste made of fresh rhizome is applied on the head in cases of vertigo Fresh juice is cooling Fumes of burning root is employed during hysteric fits" (Assistant

Turmeric; Long and Round Zedoary

CURCUMA
Zedoaria.

MEDICINE.

powdered root is used as a fumigation in commencing catarrhs. The inhalation is generally taken at night and no fluid is allowed for some hours afterwards. The effect is said to be in many cases a complete cure.

and is used for colouring confections, &c.

Chemistry of Turmeric.—Dr. Dymock gives a brief sketch of the chemical history of this subject which should be consulted. "*Curcumin*, the yellow colouring matter of turmeric, has been examined by several chemists, whose experiments have led to the conclusion that its formula is either $C_{10}H_{10}O_2$ or $C_{18}H_{18}O_4$, that it melts at 172° , forms red brown

FOOD.
Condiment.

2491

Curry

Powder.

2492

CHEMISTRY.

2493

Curcuma pseudo-montana, Graham

2494

Vern.—*Sinderwans, sinderbur, sindelwan, hellounda*, BOMB

Habitat.—Said to be a native of the Konkan, springing up at the beginning of the rains.

Food.—"The tubers, which are perfectly white inside, are boiled and eaten by the people during seasons of scarcity. Perhaps this plant, too, yields a part of East India arrowroot, that which comes from Ratnágiri is manufactured from its tubers" (*Lisboa, Dals and Gibs*).

FOOD
Rhizomes.

2495

Arrowroot.

2496

2497

C. rubescens, Roxb

Habitat.—"A native of Bengal, flowering time in the months of April and May, soon after which the leaves appear, and decay about the beginning of the cool season, in November. Every part has a strong but pleasant aromatic smell when bruised, particularly the root." (*Roxb*)

Food.—Roxburgh and Voigt say the pendulous tubers of this species yield a form of arrowroot.

FOOD.
Arrowroot.

2498

2499

C. Zedoaria, Roscoe (non-Roxb); Wight, *lc*, t. 2005.

THE LONG AND THE ROUND ZEDOARY.

Syn.—*C. ZERUMBET, Roxb*

C. 2499

CURCUMA
Zedoaria.

Long and Round Zedoary.

Vern — *Kachura*, HIND; *Sati, sheri, kachura*, BENG.; *Sati, karch*
Sans; *Zurumbdd*, ARAB.; *Kashur, urak-el-kisur*, PERS; *Kach*
Bomb; *k*
gaddala, *k*
kishanna, *k*
Fleming, *k*

References —

274; *Rhede*,

232;

*Dispen**Mot*

771,

*U S**Pl anc**I*, 159*Ind*,*Birdwood*, *Bomb Pr*, 87; *Balfour*, *Cyclop*, 859; *New Off Guide to**Mus of Ec Bot*, 62

Habitat — *Roxburgh* says it is a native of Chittagong, from which pl

ABIR.
2500

2501

solc

The

to c

composition. In Bengal the root-stocks of *C. Zedoaria* *Roscoe* are used

Zedoary.
2502MEDICINE
Rhizomes
2503

C. 2503

Long and Round Zedoary; the Dodder

CUSCUTA
reflexa.

properties Employed in native practice as a stomachic, and also applied to bruises and sprains "The natives chew the root to correct a sticky taste

MEDICINE.

Special Opinions — § "The rhizome of this plant is the *Amba-haldi* of the Bombay bazar Bruised with alum in water, it is applied to bruised joints and other parts to remove echimoses" (*Assistant Surgeon Sakharam Arjun Ravat, L M, Girgaum, Bombay*) "Small bits of the rhizomes are put in the mouth and chewed to allay cough" (*Assistant Surgeon Anund Chunder Mukerji, Noakhally*) "Demulcent, expectorant, and aromatic, dose about 1 drachm" (*Civil Surgeon John McConaghey, M D, Shahjahanpore*) "The rhizome is considered to be a cooling medicine, also tonic and expectorant" (*Surgeon-Major J M. Houston, Durbar Physn, Travancore, and Civil Apoth. John Gomes, Medical Storekeeper, Trevandrum*) "This is the *Kochora* of the bazar It is used as an odoriferous ingredient of the cosmetics used for the cure of chronic skin diseases and internally as a mild aromatic stimulant in fever and colds" (*Assistant Surgeon Sakharam Arjun Ravat, L M, Girgaum, Bombay*) "The roots imported into Leh as *kachür*, *judwar*, and called by the Bhotas '*Bozbrga*' employed in Yarkand for washing the body, acting as a rubefacient (*Surgeon Major J. E T. Aitchison, Simla*) "The rhizomes are used by singers as a masticatory for clearing the throat of tenacious mucus, they are also used in cases of irritation

Judwar of
Yarkand.
2504

Linn, see A 430

Perfumery — The rhizomes of this plant constitute one of the most important articles of native perfumery

PERFUMERY.

Trade — Dymock says the Bombay supply comes from Ceylon, value Rs 20 to Rs 30 per candy of 7 cwt: as already stated, Roxburgh affirms that Bengal gets its supply from Chittagong

2505
TRADE
2506

Curcuma Zerumbet, *Roscoe* (non *Roxb*)

2507

The writer is unable to isolate the economic facts recorded by certain authors under this name from those given for *Curcuma Zedoaria*, and he suspects that all refer to one and the same plant, or to Roxburgh's *Zingiber Zerumbet*

CUSCUTA, Linn, *Gen Pl.*, II, 881.Cuscuta reflexa, *Roxb*; *Fl Br. Ind.*, IV., 225, CONVOLVULACEÆ

2508

THE DODDER

Syn — C. GRANDIFLORA, Wall; C. VERUCOSA, Sweet, C. MACRANTHA, Don
Ve-

CUSCUTA
reflexa.

The Dodder or Cuscuta.

Some confusion exists regarding the vernacular names given to the species of *Cuscuta*. Dymock describes three species, two of which he has not determined botanically: he gives *Adzraeli* as the local Bombay name for *C.*

in the Western Himalaya, growing on the spiny plant—*Prinsepia utilis*. Roxburgh, who first described that species, states that it was found growing on *Crotalaria juncea*. The *Flora of British India* justly remarks that it is a puzzle to know where Roxburgh found it since the species, as known to modern botanists, does not occur much below 6,000 feet. It is distributed from Simla to Kashmir, Beluchistan and Afghanistan. Roxburgh gives it the name of *algusi*, and calls *C. reflexa kudi-algusi lata*, a name doubtless given in allusion to the yellow colour of the whole plant when mature. Stewart distinguishes four species, *C. macrantha*, *Don*—the *niltadhari*; *C. pedicellata*, *Led* (the *kwiklapot*, *brand* or *amla*), *C. planiflora*, *Fenore*, and *C. reflexa*, the *dhdsbel*, *astimun*, *kasús*.

Habitat.—An extensive parasitic climber, making the trees quite hoary upon which it occurs, often growing to such an extent as to completely cover every bough and leaf. It occurs throughout the plains of India and ascends the Himalaya to about 8,000 feet.

Dye.—Mr Baden Powell states that at Jhelam this plant is sometimes used as a dye. It would be a great matter if it could be utilised in this manner, as many trees are often completely covered and often killed by the plant. The dye is apparently unknown in Bengal. Mr. Baden Powell does not mention the colour, it is probably a yellow. Drury says it is

DYE.
2509

MEDICINE
Plant
2510

Seeds.
2511

Stems.
2512

FODDER.
2513

Dymock says of the Persian dodder—*Astimun*—that it “has a bitter taste, in Arabic and Persian works it is described as the *Astimun* of the Greeks, which had so great a reputation as a remedy in melancholy madness; it is still a medicine of importance with the hakims of India,

dis-
tant
ation

“Edgeworth mentions that the mountaineers believe that crows pluck sprigs of this to drop into water, when they become snakes, and so furnish food for them.”

C. 2513

The Guar.

CYAMOPSIS
psoraloides.

Cus-cus (khus-khus), see *Andropogon muricatus*, A. 1097.

Cuscus seeds, see *Papaver somniferum*

Cusparia or Angustura bark, see *Galipea Cusparia*, St. Hil., RUBIACEÆ.

Custard Apple, see *Anona squamosa*, A. 1166.

Cutch, see *Acacia Catechu*, A. 135.

Cuttle-fish, see *Molusca*.

CYAMOPSIS, DC.; *Gen Pl.*, I, 493.

Cyamopsis psoraloides, DC ; *Fl Br. Ind.*, II, 92, LEGUMINOSÆ

2514

Vern.—*Guar*, *dararhi*, *kuwara*, *kauri*, *syansundari*, *phaligawar*, *kachhur*, *khurli*, *khulti*, N-W P. and OUDH, *Guwar*, GUJ, *Gauri*, *mutki*, *gawar*, BOMB., *Buru raser*, SANTAL, *Pai pdaon*, BURM. (*Kuru*, *Pegu Rept.*)

Habitat.—Cultivated in many parts of India from the Himalayas to the Western Peninsula. It is a robust erect annual, 2 to 3 feet high, grown as a rainy season crop.

Cultivation.—In the Bombay Gazetteer (Gujarat) it is said to be grown as

CULTIVA-
TION.
2515

different purposes,—as a vegetable for human consumption, and as a pulse for horses and cattle. For the former purpose it is invariably grown on highly manured land near villages, and assumes a much more luxuriant habit of growth than when grown for cattle. The portion eaten as a vegetable is the pod, which is plucked while green, after the fashion followed with the French beans of English gardens. As a cattle fodder it is grown for its grain and is then sown on light sandy soil, side by side

FOOD
Vegetable.
2516
Horse-food.
2517
Pulse
2518
FODDER.
2519

stri
catt
for

in Fatehpur and Allahabad. The value of a purchased animal is

CYANOTIS
tuberosa.

The Spider worts

noticed It occupies there more than ten times as large an area as in any other Division The cultivation of guar also reaches its maximum in the same tract and is an indication of the care of agricultural stock which one

the Panyab proper which exhibits a sample, the pulse is stated by the

CYANANTHUS, Wall ; Gen Pl, II 557

2520

Cyananthus, sp. (? C limfolius, Wall), Fl Br. Ind, III, 434, [CAMPANULACEÆ

Vern — Murra, Pb

Habitat — A plant with pretty blue flowers, growing at 10 000 to 12 000 feet in Chumba "

MEDICINE
2521

Medicine — The calyces are eaten, being mawkish sweet, and are said to be good for asthma " (Stewart, Pb Pl)

CYANOTIS, Don ; Gen Pl, III, 831, Wight, Ic, t. 2082 & 2089

2522

Cyanotis axillaris, Ræm et Schultes, DC, Monogr Phan, III, 244, Clarke's Commelinaceæ, table 35, COMMELINACEÆ

ONE OF THE SPIDER WORTS

Vern — Nirpi li (Rheede), TAM, Saltraj, bagha-nulla (Ainslie), HIND Itsaka (Lisboa), BOMB

MEDICINE
2523

Habitat — A herbaceous annual, met with in many parts of India, distributed to the Malay, China and Australia

Medicine — Rheede says that on the Malabar coast this is viewed as a

FAMINE
FOOD
Seeds
2524
2525

of this as also of Commelina communis, were eagerly sought for during the Bombay famine, they are wholesome and nutritious

C. tuberosa, Ræm & Schultes, DC, Monogr Phan, III 249

Syn — TRADESCANTIA TUBEROSA Roxb C ASCENDENS Dals in Hook Jour Bot p 343 (1852), C SARMENTOSA, Wight Ic, 2087

MEDICINE
Root
2526
FOOD
Leaves
2527

Vern — Mero n chunchi (a name given from the resemblance of the roots to the papillæ of the goat) Hodo jereng arak (the vegetable) SANTAL

Medicine — The Rev A Campbell says the ROOT is given in long continued fevers and also for worms in cattle

Food — The LEAVES are eaten by the Santals as a pot herb

C. 2527

SEIR FISH, Cycas or Sago Plant

CYCAS
RumphiiCYBIUM, Cu., *Dis*, *Fishes of Ind*, 254Cybium Commersonii, *Cuv. & Val*

SEIR FISH

Vern.—*Sermi*, HIND ; *unjurum* (male) *koram* (female), TEL ;
Konam, *mah-wu luachi* or *ah ku lah*, TAM ; *Chumbum*, MAL

Habit.

Medic

mended.

Quart A

taste of

to putrify

2528

MEDICINE,
Oil

2529

CYCAS, Linn., *Gen Pl*, III, 444

2530

The brief notes here given of the species of CYCAS will be found supplemented under SAGO. This has been rendered necessary, from its being often difficult to discover to which plant the earlier writers refer

Cycas circinalis, Linn., *DC Prod XVI, II*, 526, CYCADACEÆ

2531

Syn.—*C SPHERICA*, Roxb., *Fl Ind Ed C B C*, 709, *C CIRCINALIS*, Linn.
in Thwaites En Ceylon Pl, 294, *TODDER PANNA*, *Rheede, Hort*
Mal III 9

Vern.—*Orasmaro URIVA* *Maddé*, SING Under *Cycas circinalis*, Linn.,
 Ainslie gives the following names which all appear to refer to Sago and
 not necessarily to CYCAS.—*Show arisi*, TAM ; *Saukk chawal*, DUK ;
Sabudana HIND ; *Zombium*, TEL ; *Sagu*, MAL ; *Sekuhme*, SING ;
Sagu, JAVA ; *Sagu*, BALI (*Mat Ind I*, 361)

Habitat.—A palm-like tree met with on the mountains of the Malabar
 coast and in Ceylon

FOOD,
Seeds.

2532

Flour.

2533

2534

C. pectinata, Griff. as in Kurz, *For Fl Burm*, 503

Vern.—*Tlatat* NEPAL

Habitat.—An evergreen simple-stemmed palm like tree, found in
 Sikkim, Eastern Bengal, and Burma, often in *sal* or *eng* or pine forests
 (Gamble)

FOOD

Sago

2535

TIMBER

2536

C. revoluta, Thunb

2537

Often called the SAGO PALM OF JAPAN AND CHINA

Habitat.—A Japanese species often cultivated in India, has a short
 thick stem

C. Rumphii, Miq., *Gamble, Min Timb*, 415

2538

Syn.—*C CIRCINALIS* Roxb., *Fl C B C*, 709

Vern.—*Baragudu*, TEL ; *Tolima marani*, MAL ; *Moudang*, BURM

2539

C 2538

CYDONIA
vulgaris

Cycas, Quince

Habitat — A palm like tree with a simple or branched stem, abundant in the island of Andaman

RESIN

2530

MEDICINE

2540

Scales

2541

FOOD

Sago

2542

Seeds

2543

2544

RESIN

2545

that it excites suppuration in an increased short time

Special Opinion — "The scales of the cone of the male tree anodyne, dose 30 to 60 grains or more" (*Apothecary Thomas Wirt, Madanapalle, Cuddapah*)

Food — The interior of the stem yields a good quality of sago or starch, the nutty seeds are in Ceylon made into flour, but they are also eaten by the hill tribes of India

Cycas siamensis, Miq, Kurz, Burm For. Fl, II, 503

Habitat — An evergreen, low, stemless palm-like tree frequent in the eng and dry forests of the Prome district Burma

Resin — Exudes a peculiar whitish gum, like tragacanth (*Kurz*)

CYDONIA, Tourn (PYRUS, Linn), Gen Pl, I, 626

2546

Cydonia vulgaris, Pers, Fl Br Ind, II, 368, ROSACEÆ.

THE QUINCE

Syn — PYRUS CYDONIA, Linn

Vern — *Bihl* (abi, according to Ainslie), HIND, *Bam tsuntā, bamsutū* KASHMIR, *Shimai madalavirai* TAM, *Bhi tursh safarjal* ARAB, *Moodeen Sheriff* gives the following names for QUINCE seeds — *Habbus safarjal* ARAB, *Bihl-dānah beh-dānah tukhme-ābi* PERS, *Beh dānah* HIND, *Duk, Shimai madalavirai*, TAM, *Shūne-dali ibā bja* SING, *Shime-dlimba-tittulu*, TEL

References — *Brandis For Fl* *Pb Pl 80, DC, Origini Cu* *Pharm Ind, 324 Ainslie 3* *Pharm Ind 211; Dymock 3* *& Trim Mel Pl I, 106* *Mat Med Patna 10 106* *U. Pl Bomb 119 Birlinoo* *XI (Quince used in his day was brought from Crete)*

Habitat — Cultivated in Afghanistan and the North West Himalaya up to 5000 feet DeCandolle says it grows wild in the woods in the north of Persia, near

sus and in Anatolia

Sanskrit name is known

but its Persian name

quince, and for the wild plant *armud*. The names in use in Europe point to an ancient knowledge of the species to the west of its original country. DeCandolle adds that it may have been naturalized in Europe before the time of the Crusades (*Cult Pl, 237*)

OIL.

2547

Hills the ground at certain seasons

rotting under the trees. This might

use and probably is a substitute for quince

Medicine — Ainslie says — The fruit of this article which is found in India and bazars is chiefly in use amongst the Muhammadan practitioners,

MEDICINE

Seed

2548

C 2548

Quince

CYDONIA
vulgaris.

MEDICINE

who occasionally order an infusion or decoction of the SEEDS as a demul-

tural, cephalic, and
 eup, and BARK of
 account of their astri
 most, and slightly a
 native practice, the
 plants as a demul
 blisters" (*D. mock*)
 author:—"The seed

Fruit
2549Bark.
2550Mucilage
2551

corresponds in composition with that of linseed "

The seeds coagulate 40 times their weight of water (*Pharmacographia*)Special Opinions—§ "A cold infusion of the seeds forms a pleasant demulcent drink, which is much used in native practice in cases of irritation of the urinary organs" (*Pharmacographia*)

"I use it as a de
 about one drachm
 are known here as
 plants and semir
 Ahmedabad). Qu

t demulcent
of diarrhoea

ly juicy and
 odour, is also made into preserve, and, as having a powerful
 sometimes made from it It is supposed by some to have been the
 Golden Fruit of the Hesperides
 cially near Naggar), and the fruit is

FOOD,
Fruit.
2552

seed are largely imported into
 the fruit is eaten fresh, candied
 istan excels all other quinces
 no other fruit of remarkable g
 Flora makes no mention of

the
 of 5, 3, 2, according to quality. Moodeen Sherif points out that *Beh-
 dānah* and *Bé-dānah* are so much alike in sound that mistakes are likely
 to be made The latter is the name for a peculiar seedless raisin but
 is often loosely applied to all raisins.

TRADE.
2553

Cymbopogon, see Andropogon; GRAMINEÆ.

C. citratum, DC., see Andropogon citratus, DC., A. 1079

C. laniger, Desf., see Andropogon laniger, Desf.; A. 1093

C. Martini, Roxb.; Munro, see Andropogon Schoenanthus, Linn.; A.

1117

C. 2553

CYNODON
Dactylon

Artichoke, Doorwa Grass

Cymbopogon Nardus, Linn, see *Andropogon Nardus*, Linn, A. 1107

2554

CYNANCHUM, Linn, Gen Pl 762

[354, ASCLEPIADEÆ

Cynanchum pauciflorum, Br, Fl Br Ind IV, 23, Wight, Ic,

Syn — *ASCLEPIAS TUNICATA* Roxb Fl Ind Ed C-B C, 253 CYNANCHUM PAUCIFLORUM R Br in Dalz & Gibs Bomb Fl 148 CYNOCOTONUM PAUCIFLORUM Deca 310 Thwaites E Ceylon Pl 195

Vern — *Chagul pati* BENG Kan kumbala SING

Habitat — A large twining shrub met with in the Deccan Peninsula from the Concan southwards to Travancore and Ceylon This is the region given in the Flora of British India, but according to Roxburgh (*Asclepias tunicata*), it is found in Bengal also.

FOOD
Leaves
2555

Food — The Cinghalese eat the young leaves of this and of many other plants of this natural family, in their curries (*Enumeratio Plantarum Zeylanicæ*, 195)

This does not appear to be the case in Bengal, Roxburgh simply remarking that its milky juice is particularly gummy

CYNARA, Linn, Gen Pl, II, 469

2556

Cynara Scolymus, Linn, COMPOSITÆ

ARTICHOKE

" " " " " " " " " "

"
"
6,

Habitat — Cultivated to a limited extent over most parts of India for the European market

FOOD
2557

Food — The lower parts of the thick imbricated scales of the flower-fleshy are eaten the artichoke in minger says it is than in England of September 15

suitable
days
be plac
art chok
from the
the begi

CYNODON, Pers, Gen Pl, III 1164

[GRAMINEÆ

2558

Cynodon Dactylon, Pers Duthie Fodder Grasses N Ind, 52,
CREEPING PANIC GRASS OR DOORWA, COUCH GRASS

Syn — *C STELLATUS* Willd; *PANICUM DACTYLON* Linn PASPALUM DACTYLON DC DIGITARIA DACTYLON Scop

Vern — *D b dauria d bra kabbar khabbal talla tulla* Ps B rava TRANS INDUS, *Dob null dub*, RAJ, Chibbur SIND, *Dub, darbd, dābla*,

C. 2558

CYNODON
Dactylon.

Dub or Doorwa Grass.

MEDICINE.

FOOD.
2563
FODDER.
2564

roots. It is the most common and useful grass in India, and its stems as well as its roots form a large proportion of the food of our horses and cows. Mr. Duthie says it varies considerably both in habit and nutritive qualities, according to the nature of the soil or climate. It makes excellent hay and will keep for years. It is by far "the most useful of all fodder grasses, especially for horses." "It is considered to be a first class fodder grass in Australia, where it is widely distributed,

taxis.
times
oper-
roots
311"

2565

however, must possess considerable nutritive qualities, and poor soils it is able to be crushed out by inferior types of plants, but on those of fair quality it is very persistent and difficult to eradicate, the latter point is detrimental to its use as a crop to be taken in a rotation. When highly cultivated it yields heavily under irrigation and is grown for hay near some large stations. In 1868 there was a plot of this grass on the

The following system is recommended for putting down this grass:—
"The land having been well cleaned should receive a dressing of fold-yard manure: when ploughing in the manure a woman should follow, each

The Cynoglossum or Dog's tongue

CYNOGLOSSUM
micranthum.

Regarding the curing of hay the following remarks with reference to this grass are of value —

"Hariali, like most other meadow grasses, should be cut immediately the flower begins to appear, in this state the juices of the grass are more nutritious, and the hay is far superior than when made from the fully matured plant. Besides when cut before the seed appears, the plant is more vigorous and produces another crop much sooner. Hariali hay is generally considered to be one of the best for cattle and horses."

FODDER.
Hay.
2566

or at the most three days, should suffice for making the hay.

"Cutting should not commence until the dew is off the grass. The grass should remain on the ground for an hour or so after being cut. It should then be turned and tossed until sun-set. It cannot be tossed too much during a hot sun. To preserve the green colour and aroma of the hay it is absolutely necessary to keep it moving. At night, if the dews are heavy, it should be put up in small cocks, each containing from two to three cwt. The cocks should be covered with straw or matting, and be open at the top. The cocks should be turned over at night."

of course putting it again into cock at night.

"Hay thus rapidly made is rich in saccharine matters, and is, therefore, very liable to heat and ferment, this, to a moderate extent, does no harm."

2567

"It is very palatable and is eaten with great abundance, and is of a superior quality. It grows luxuriantly in the rivers in the southern division, and in the district. The juice of the leaves is used for medicinal purposes." (Topography of Dacca by J. J. J.)

2568

lor, 60)

CYNOGLOSSUM, Linn : Gen Pl, II, 848

Cynoglossum micranthum, Desf., Fl Br Ind, IV, 156

Vern — Nilakras, Pa, Oudhuphull, Guj, Adhopushpi, Sakas, J. J. J., henda, SING

4/10

C. 2569

Mats and Matting.

CYPERUS
corymbosus

Cyperus bulbosus, Vahl., see *C. jemicus*, Rottb ; CYPERACEÆ.

2584

C. compressus, Linn ; Clarke in *J Linn. Soc.*, XXI., 97

Vern.—Chuncha, BENG., Salitunga, TEL., Wek-tamyet, BURM.

References.—Roxb., *Fl Ind*, *Ed C B C*, 65; Dalz & Gibs, *Bomb Fl*, 282, *Cyperus* in *Griff Itin Notes* No 167, p 12, and 191, p. 362; Kurz, *Rept*, Pegu

Habitat.—A common species throughout India, ascending the hills to 2,000 feet in altitude. A special form is known as *var. pectiniformis*. This is said to occur in Lucknow, Chutia Nagpur, and Assam. Thwaites says it is very common in the warmer parts of Ceylon. Roxburgh remarks that it "delights in a moist soil."

C. corymbosus, Rottb ; C.B. Clarke in *Jour. Linn Soc*, XXI, 158.

2585

Syn.—*C. SEMINUDUS*, Roxb., *Fl Ind*, *Ed C B C*, 63, *Nees in Wight*, *Contrib*, p 80; *PAPYRUS PANGOREI*, *Nees in Wight, Contrib*, p 88, in part

Vern.—Gola mekhi, BENG.; Godá tunga kúda (Roxb.) and Goddu-tunga kodu (Elliot), TEL., Galchi, SING.

Habitat.—Found throughout India, and on the islands of Ceylon.

It should be observed that the name *C. Pangore* is open to the greatest possible ambiguity. The Madras plant mentioned under that name by Dr Bidie, CIE, is *C. corymbosus*, Rottb., *var. Pangore*, Rottb., but *C. Pangore*, Roxb., is *C. malaccenses*, Lam ; *C. Pangorei*, Thw., is *C. ...*

FIBRE.
2586MATS,
2587
Tinnevely,
2588Palghat,
2589

CYPERUS
Haspan.

Sedges used for

Tinnevely, and the article is therefore heavier, coarser in texture, and not so flexible."

FODDER.
2590

Fodder.—"Cattle are not fond of it, and it is only eaten occasionally by buffaloes" (Roxb)

2591

Cyperus elegans, Linn, C B. Clarke, Linn Soc Jour, XXI, 125

Syn —C MÆSTUS, Kunth; C NIGROVIRIDIS, Thw, En Ceylon Pl, 344

Vern —*Wek chan*, BURM (*Kura*, Pegu Rept)

Habitat —A native of Bengal and the Malay Peninsula, Sikkim 1,500 feet, Assam, Khasia hills 1,200 feet Sylhet, Yunan, Chittagong, Mergui, Tenasserim, and the Andaman Islands

2592

C. esculentus, Linn, C B Clarke, Jour Linn Soc, XXI, 178

Syn —C TUBEROSUS, Rottb

Vern —*Kaserá dila*, PB, *Sha ts'au* CHINESE

Habitat —There are five or six distinct forms of this plant, of which two occur in India viz, *forma tuberosa* (sp Rottb) in Madras and *forma hindustanica* in Northern India

MEDICINE
Root

2593

FOOD

Root

2594

Medicine and Food —Stewart says "In N-W. Provinces the root is used as food, and is officinal as *kaserá* The *dila* root, mentioned by Bellew as eaten in the Peshawar valley, may be the same *Dila*, however, appears to be a generic name for the CYPERACEÆ, the roots of several of which are eaten by pigs, and their stems, &c, browsed on by cattle, as is

FIBRE

2595

Coffee Sub-

stitute

2596

2597

C. exaltatus, Retz; C B Clarke, Linn Soc Jour, XXI, 186

Syn —C UMBELLATUS Vahl according to Roxb, Fl Ind, Ed C B C, 69, C VENUSTUS R Br, Thwaites En Ceylon Pl, 432 (nec Nees nec Kunth), C ALTUS Nees, in Wight, Contrib, 84

Vern —*Pedda shaka*, TEL

Habitat —Commonly found in Bengal (Chuttia Nagpur, Rajmahal, Mysore, Madras, Central

on "A large species, grow-

FIBRE.

2598

Mats

2599

note —This sedge is often used for matting, Mr C B Clarke de-

fore be held a distinct species but whatever it is considered, " "

2600

C. Haspan, Linn, Clarke, Linn Soc Jour, XXI, 119

Syn —C UMBELLATUS, Vahl, is the *Pedda sika* of the Telugus.

C. 2600

CYPERUS
rotundus

Sedges used for

Habitat.—Roxburgh says of his *C. Pangorei* that it is a native "of the banks of the Ganges, and serves, with *C. inundatus*, the same useful marks that ng the cold he Sunder-Japan, and

2610

Cyperus niveus, Retz, C. B. Clarke, Linn Soc Jour, XXI., 108

Vern.—Birmutha, SANTAL

Habitat.—Throughout India and Burma (Beluchistan, Kashmir, Panjáb, Kumaon, Simla, Kulu, Nepal, Sikkim, Assam, Bengal, Chutia Nagpur, Rajmahal, &c.), Madras, &c., &c. A native of shady moist pasture land (Roxb)

C. pertenuis, Roxb, see *C. scariosus*, R Br

2611

C. Pongarei, Rolfb, as in Roxburgh, see *C. malaccensis*; and for other plants named by different authors as *Cyperus Pangorei*, see *Cyperus corymbosus*

2612

C. rotundus, Linn, C. B. Clarke, Linn. Soc Jour, XXI, 167.

Syn.—*C. HEXASTACHYOS*, Roxb

Vern.—Muthá, motha, BFNQ, Batha bijir, MUNDARI, Utru banda, U S m B SING

References.—Roxb, Fl Ind, Ed C B C, 66, Jour As Soc, Pt II (1867), p 82, Home Dept Off of Crops and Forests, Di. Hore's Tour: Andhra pp Ind 128; U W Ind, 2nd Pb Pr 382,

Habitat.—A plentiful species in India occurring from Kuram Valley, Afghanistan, Gilgit, and Kashmir to Simla, Garhwal, and the Kharia scending the nd Poonr to of the plant

DYE

2613

OIL,

2614

Dye.—Used in certain dye preparations to impart a perfume to the fabric.

MEDICINE,

Roots

2615

gent Stimulant and diuretic properties are also attributed to them. They are further described as vermifuge. In native practice, they are held in great esteem as a cure for disorders of the stomach and irritation of the bowels. The bulbous roots are scraped and pounded with green ginger, and in this form mixed with honey they are given in cases of

C. 2615

Mats and Matting.

CYPERUS
SCARIOSUS.

MEDICINE.

dysentery in doses of about a scruple (*Med. Top of Dacca* by J. Taylor, p 54) "In the Concan the fresh tubers are applied to the breast in the form of *lep* (maligma) as a galactagogue C. rotundus is the *κύτερος* of the Greeks and is mentioned by Dioscorides, who says it is the *Juncus* or *Radix Junci* of the
gue, and applied to
it is also an ingredie
as an aromatic pla

is mentioned in the Iliad (21, 351), and Odyssey (4, 603), and by Theophrastus in his fourth book, it appears to have been a favourite food of horses Pliny (21, 18) calls it *Juncus triangularis* or *angulosus*; it is probably the *Juncus* of Celsus (3, 21) mentioned as an ingredient in a diuretic medicine for dropsy, although he calls it *Juncus quadratus*" (*Dymock*, p 844) Arabian and Persian writers describe the drug as
te that it is
doses as an
ingredient

cine) "The roots are in Chutia Nagpur used in fever" (*Rev A Campbell*) "The fresh roots are stimulant and diaphoretic" (*Bombay Gazette* V I, p 14)

Fodder.—Cattle eat this so-called grass, and hogs are remarkably fond of the roots

FODDER,
2616

Cyperus scariosus, R. Br.; C B C, Linn Soc. Jour, XXI, 159

2617

Syn.—CYPERUS PERTENUIS Roxb, Fl Ind, Ed C B C, 66

Vern.—*Nagar mothā*, HIND, *Nagar mutha* BENG, *Lamdā* MAR, *Soade kaji*, soad, ARAB, *Mushke-namin*, PERS, *Nagar-mustaka*, SANS, *Nagar motah*, DEC; *Muttah k ch*, *kōraik kishangu*, TAM, *Tinga gaddala veru*, *kēlatunga muste* TEL, *Kēra kishanna*, MAL, *Konnari gadde*, KAN, *Vomon nuu*, BURM

References —Roxb, Fl Ind, Ed C B C, 66 *Med Top Ajmir* 147, *Dymock Mat Med W Ind*, 2nd Ed, 815, *Irvine, Mat Med Patna*, 75 *Birdwood Bomb Pr*, 94; *Liottard, Dyes, Supp*, IV

Habitat —A delicate, slender grass, met with in damp places in Bengal, Oudh, and rare in the Panjāb, by no means so common a plant as C rotundus

"Nagar mothā, DUK, id II, 162) under the

Dye —The rhizomes are used in dyeing to give a scent to the fabric, and as a perfume for the hair Roxburgh describes them as "tuberous with many dark coloured villous fibres" "Its naked delicate form, small

DYE,
2618

MEDICINE,
Root,
2619

C 2619

CYPERUS
tegetum

Sedges used for

MEDICINE

Cyperus, but consider it to be inferior to *C. rotundus*. "Two kinds of *Nagarmoth* are met with in the Bombay market—Surat and Kathiawar, the first is heavier and more aromatic than the second. Value, Surat, Rs 2 per maund of 37½ lb, Kathiawar Rs 1½. The Surat *Nagarmoth* is probably obtained from Rājputana, where the plant is common in tanks (Dymock). U O Dutt says "The root of *C. pectenatus* is somewhat tuberous with many dark coloured villous hairs. It grows in low wet places, and is chiefly used in the preparation of medicated oils."

Special Opinions.—§ Roots, when bruised have a fragrant smell, and for this reason native females keep a stock of the powdered root to wash their bodies with' (*Honorary Surgeon P. Kinsley, Chicacole Ganjam Madras Presidency*). 'Is given in conjunction with Valerian in cases of epilepsy' (*Surgeon-Major C. W. Calthron, M.D., 4th Bengal Cavalry, Morir*). "The root is astringent, useful in diarrhoea" (*Surgeon-Major J. M. Houston, Durbar Physn, Travancore and Civil Apothecary*). The decoction is used in C. H. Peacocke, Surgeon Major Robb,

2620 *Cyperus stoloniferus*, Retz, C. B. Clarke, Linn Soc Jour, XXI, 172

Syn.—*C. LITTORALIS*, R. Br., *C. TUBEROSUS* Baker

Vern.—*Jatama* is a name given in South India to this plant

Part used A L C S J L C

FRUMERY,
2621

called *Sanbal-Hind* and *Sanbal-ul-tarb* and in Upper India *Jatamansi* and *Balchhar*. But as the true plant is only found at great elevations beyond the tropics, the long tubers of various species of grass (*Schænanthus*) are also under the names of

2622 *C. tegetiformis*, Roxb., C. B. C., Linn Soc Jour, XXI, 157

Syn.—*C. NUDUS* Roxb., Fl Ind, Ed C B C pp 63 and 70, *C. BENGALENSIS* Spreng

Vern.—*Gula-methi* BENG., *Sura* SINTAL

Habitat.—A native of low wet places over Bengal, flowering during the rains' (*Roxb*). Clarke mentions as localities—Calcutta, Chittagong, Noakhali, Burisal, Mymensing, Pandua, and Assam. He also states that the plant occurs in China and Japan.

Fibre.—Roxburgh writes "This species is very like *C. tegetum*, and about the same size, though I am informed it is never used for mats. To know it from *C. tegetum* attend to the involucre, which in this is only about one-fourth the length of the umbel, but in that as long or longer."

FIBRE,
Mats
2623

2624 *C. tegetum*, Roxb., C. B. Clarke, Linn Soc Jour, XXI, 160

Syn.—*C. CORYMBOSUS* Koenig in part *C. SCHIMPERIANUS* Steud
C. DENISCENS Steud *C. PANGOREI* Thwaites (non Retz) Enum
Pl Zeyl 344 *PAPYRUS DENISCENS* Nees in Richt Contr 6, 89; *C. PANGOREI*, Nees (the greater part) and *C. CORYMBOSUS*, Nees

C. 2624

5

Mats and Matting

CYPERUS
tegetum

Note by Mr. Clarke. This plant abundant in India is the authentic
C. TEGETUM Roxb. It differs decidedly from C. CORYMBOSUS in the

Vern.—*Mudar ka* BENG. *Wella* BURN.

Habitat.—A common species in India, Abyssinia and Egypt. Mr.
Clarke mentions the following localities: Almora (1200 feet), Chumba

FIBRE

Mats
2625

maker passes the culms with the hand alternately over and under the
successive threads of the warp and presses them home.

In different districts of India it is believed that two or three allied
species are used for this purpose. In Madras the form *C. corymbosus*
seems to be chiefly used. Royle repeating Roxburgh states that
the culms or stalks of the plant when green are split into three or four
pieces which on drying contract so much as to bring the margins in

TRADE
2626

